



HAL
open science

An asymptotic criterion in an explicit sequence

André Voros

► **To cite this version:**

| André Voros. An asymptotic criterion in an explicit sequence. 2015. cea-01166324

HAL Id: cea-01166324

<https://hal-cea.archives-ouvertes.fr/cea-01166324>

Preprint submitted on 30 Jul 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An asymptotic criterion in an explicit sequence

André Voros

CEA-Institut de Physique Théorique de Saclay (CNRS UMR 3681)

F-91191 Gif-sur-Yvette Cedex (France)

E-mail: `andre.voros@cea.fr`

June 28, 2015

Abstract

We report a novel asymptotic (large-order) behavior in an explicit sequence built out of the Bernoulli numbers and analyzed by a variant of instanton calculus or Darboux's theorem.

We will use: B_{2m} : the Bernoulli numbers; γ : Euler's constant; $k!! = k(k-2)(k-4)\dots$: double factorial (with $0!! = (-1)!! = 1$ as usual).

The real sequence explicitly spelled out for $n = 1, 2, \dots$ as

$$u_n = (-1)^n \left[2^{-2n} \sum_{m=1}^n \frac{(-1)^m}{2m-1} \binom{2(n+m)}{n+m} \binom{n+m}{2m} \log \frac{|B_{2m}|}{(2m-3)!!} - \frac{(2n)!!}{2(2n-1)!!} \log 2\pi \right] \quad (1)$$

can be thus numerically computed (trivially to thousands of terms), and very early it satisfies (figs.)

$$u_n \approx \log n - 1.703 . \quad (2)$$

This can be validated *assuming the Riemann Hypothesis* (= RH), as

$$u_n \sim \log n + K, \quad K = \frac{1}{2}(\gamma - \log(2\pi^2) - 1) \approx -1.70269564368 . \quad (3)$$

With RH verified up to an ordinate $T_0 \gtrsim 2 \cdot 10^{12}$ currently, [5] it is plausible indeed to witness the behavior (3).




Figure 1: The coefficients u_n computed by (1) up to $n = 3500$, on a logarithmic n -scale, vs the function $(\log n + K)$ of (3) (straight line); the first values are $u_1 = \log \pi - \frac{1}{2} \log 54 \approx -0.84976213743$, $u_2 = -\frac{4}{3} \log \pi + \frac{23}{24} \log 2 + \frac{55}{24} \log 3 - \frac{35}{24} \log 5 \approx -0.69148426053$, $u_3 \approx -0.46222439972$.




Figure 2: As fig. 1 but for the remainders $\delta u_n = u_n - (\log n + K)$, on a very dilated vertical scale. (The connecting segments between data points are only drawn for clarity.)

Now by large-order analysis through exponential asymptotics, [3][1] we found that *if RH is false*, u_n will also admit a clear-cut asymptotic form but of a wholly different nature, dominated by individual terms $F_n(\rho)$ contributed by every zero $\rho = \frac{1}{2} + t + iT$ off the critical line with $0 < t (< \frac{1}{2})$:

$$F_n(\rho) \sim f(\rho)(-1)^n \frac{(2n)^{\rho-1/2}}{\log n} \quad \text{for } n \rightarrow \infty, \quad (4)$$

where f is an explicit function independent of n with the main property

$$|f(\rho)| \approx |T|^{-t-2} \text{ for } |T| \gg 1 \implies |F_n(\rho)| \approx |T|^{-t-2}(2n)^t / \log n. \quad (5)$$

Each such $F_n(\rho)$ will ultimately dominate (3) in the $n \rightarrow \infty$ limit, but starts out exceedingly tiny at low n , and does not approach unity until

$$n \gtrsim \frac{1}{2}|T|^{1+2/t} \quad (\text{at best } O(|T|^{5+\epsilon}) \text{ for } t \rightarrow \frac{1}{2}^-); \quad (6)$$

yet we think that, with efficient signal-processing, the “signal” $F_n(\rho)$ of ρ within u_n ought to be detectable much sooner than at (6) (at $n \gtrsim O(|T|^{1+1/t})$ or even less, but within $n \gg |T|$). Still, to seek a violation of RH and verify the form (4), $|T| > T_0$ is necessary, and $T_0 \gtrsim 2 \cdot 10^{12}$ implies very large n -values.

The major issue is then that u_n is an alternating sum of terms which turn out to be exponentially larger by an order of $(3 + 2\sqrt{2})^n$. Increasingly delicate cancellations thus take place, requiring a precision beyond $\approx 0.7656 n$ decimal digits to evaluate u_n by (1). On the other hand, this purely technical demand seems to be the *sole* obstacle raised by the use of (1) at unlimited n .

While other sequences sensitive to RH for large n are known, [6][2][7][4] we are unaware of any previous case combining a fully *closed form* like (1) with a practical sensitivity threshold of *tempered growth* $n = O(T^\nu)$.

Details and derivations are currently under completion. [8]

References

- [1] R. Balian, G. Parisi and A. Voros, *Quartic oscillator*, in: *Feynman Path Integrals* (Proceedings, Marseille 1978), eds. S. Albeverio *et al.*, Lecture Notes in Physics **106**, Springer, Berlin (1979) 337–360.

- [2] L. Báez-Duarte, *A sequential Riesz-like criterion for the Riemann Hypothesis*, Int. J. Math. Math. Sci. **21** (2005) 3527–3537.
- [3] R.B. Dingle, *Asymptotic Expansions: their Derivation and Interpretation*, Academic Press (1973).
- [4] Ph. Flajolet and L. Vepstas, *On differences of zeta values*, J. Comput. Appl. Math. **220** (2008) 58–73, and refs. therein.
- [5] X. Gourdon, *The 10^{13} first zeros of the Riemann Zeta function, and zeros computation at very large height*, preprint (Oct. 2004), <http://numbers.computation.free.fr/Constants/Miscellaneous/zetazeros1e13-1e24.pdf>
- [6] X.-J. Li, *The positivity of a sequence of numbers and the Riemann Hypothesis*, J. Number Theory **65** (1997) 325–333.
- [7] K. Maślanka, *Báez-Duarte's criterion for the Riemann Hypothesis and Rice's integrals*, preprint (2006, revised 2008) arXiv:math.NT/0603713v2.
- [8] A. Voros, to appear shortly (2015).