

Synergistic effects in ambiphilic phosphino-borane catalysts for the hydroboration of CO₂

Anis Tlili, Arnaud Voituriez, Angela Marinetti, Pierre Thuéry, Thibault Cantat

► **To cite this version:**

Anis Tlili, Arnaud Voituriez, Angela Marinetti, Pierre Thuéry, Thibault Cantat. Synergistic effects in ambiphilic phosphino-borane catalysts for the hydroboration of CO₂. *Chemical Communications*, Royal Society of Chemistry, 2016, 52, pp.7553-7555. 10.1039/c6cc02809h . cea-01329300

HAL Id: cea-01329300

<https://hal-cea.archives-ouvertes.fr/cea-01329300>

Submitted on 15 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Synergistic Effects in Ambiphilic Phosphino-Borane Catalysts for the Hydroboration of CO₂

Anis Tlili,^{a,b} Arnaud Voituriez,^{b*} Angela Marinetti,^{b*} Pierre Thuéry^a and Thibault Cantat^{a*}

The benefit of combining both a Lewis acid and a Lewis base in a catalytic system has been established for the hydroboration of CO₂, using ferrocene-based phosphine, borane and phosphino-borane derivatives.

Featuring a carbon atom at the +IV oxidation state, CO₂ is a stable molecule and its conversion to either fuels or chemicals necessitates the use of a mild reductant and an efficient catalyst, to promote the cleavage of C–O bonds and the formation of novel C–H and/or C–C bonds.¹ In this respect, hydroboranes (R₂BH) have a mild reduction potential (ca. –0.5 V vs NHE) well-suited for the reduction of CO₂ to the methanol level.² In contrast to borohydrides, the reduction of CO₂ with hydroboranes requires a catalyst.³ In fact, the catalytic hydroboration of CO₂ has first been reported in 2010 by Guan and coworkers who showed that nickel(II) pincer complexes were able to catalyze the reduction of CO₂ to methoxyboranes in the presence of 9-BBN, catBH and pinBH.⁴ Shortly after, Sabo-Etienne and Bontemps described efficient ruthenium catalysts for this transformation and a variety of inorganic and organometallic catalysts have been reported since then.⁵ Notably, the polarized B–H bond in hydroboranes can also be activated by simple organic compounds: the organocatalyzed hydroboration of CO₂ has been unveiled in 2013 by Fontaine and coworkers with P/B Frustrated Lewis Pairs (FLPs), and by our group with guanidines and amidines bases as well as using an isolated FLP-CO₂ adduct.⁶ This collection of catalysts has been completed with several bases among which phosphines proved to be potent catalysts for this reduction process.⁷ In this context, we have also shown recently that proazaphosphatranes enable the first catalytic methylation of amines involving CO₂ reduction with hydroboranes.^{7a} Thus, since the hydroboration of CO₂ is a reaction of fundamental importance, it becomes crucial now to better apprehend the role of the organocatalysts in these reduction processes. In particular, the possible positive influence of combining a Lewis base with a Lewis acid in the same catalytic system is still to be established.


To address this question, we report herein the synthesis of simple phosphine, borane and phosphino-borane compounds based on the same ferrocene backbone (Chart 1). Comparing the catalytic performances of the separated Lewis acids and bases and their combination as intermolecular or intramolecular systems demonstrates the benefits of the ambiphilic systems, for the first time.



The ferrocene backbone is a well-suited platform to assemble Lewis acids and bases by substituting the cyclopentadienyl ligands on iron. This approach has been exemplified over the last years by the groups of Bourissou, Aldridge and Erker to access novel FLPs and ambiphilic


ligands.⁸ Additionally, rotation around the ferrocene axis provides some flexibility that can adapt the catalyst to the geometric constraints associated with the catalytic pathway. In this respect, ferrocene based FLPs differ from the system used by Fontaine *et al.* for the hydroboration of CO₂, where a benzene ring maintains a rigid structure between the P and B centres.^{6a}

In this study, diphenylphosphinoferrocene, **1**,^{9a} was isolated by quenching the lithiated ferrocene^{9b} with chlorodiphenylphosphine (Scheme 1). Similarly, the corresponding ferrocenyl-pinacolborane **2-Bpin** was obtained in a moderate 25% yield by reacting bis(pinacolato)diboron with lithiated ferrocene. Crystals of **2-Bpin** were obtained by recrystallization from dichloromethane (see ESI). When dimethylboron fluoride was used as an electrophile, **2-BMes₂** was obtained in a significantly better yield of 71% (Scheme 1).



Scheme 1. Synthesis of **1**, **2-Bpin** and **2-BMes₂**

The phosphino-borane derivatives **3-Bpin** and **3-BMes₂** were prepared using an established procedure based on a sequential halogen/lithium exchange starting from 1,1'-dibromoferrocene (Scheme 2).^{8a} Thus, 1-bromo-1'-(diphenylphosphino)ferrocene **4** was obtained in 87% isolated yield. Subsequently, a second bromine/lithium exchange was conducted and the lithiated ferrocene was quenched with bis(pinacolato)diboron, yielding the desired product **3-Bpin** in a low 18% yield. Other pinacol-boron electrophiles including 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane, were also tested, without success. Alternative routes were explored to improve the yields of **3-Bpin**. Unfortunately, inverting the reaction sequence by quenching 1-bromo-1'-lithioferrocene with bis(pinacolato)diboron first proved unproductive. Also, quenching **4** with trimethyl borate and subsequent reaction with pinacol failed.

Scheme 2. Synthesis of **3-Bpin** and **3-BMes₂**

As reported by Bourissou *et al.*, replacing bis(pinacolato)diboron with dimesitylboron fluoride yielded **3-BMes₂** in 82% yield.^{8a} It is noteworthy that the P/B derivatives **3-BMes₂** and **3-Bpin** display ³¹P NMR chemical shifts similar to those of (diphenylphosphino)ferrocene (**1**: $\delta = -16.2$; **3-Bpin**: $\delta = -17.4$; **3-BMes₂**: $\delta = -19.3$ ppm). This suggests the absence of any interaction between the two heteroatoms, as noted previously by Bourissou *et al.* Moreover, no interaction with CO₂ has been observed either by ³¹P or ¹H NMR, when a solution of **3-BMes₂** is exposed to an atmosphere of CO₂.¹⁰

Having in hand a homogeneous series of P-, B- and P/B-based catalysts, their reactivity has been investigated in the catalytic hydroboration of CO₂. At room temperature, phosphine **1** (1 mol%) exhibits a negligible catalytic activity in the conversion of CO₂ and 9-borabicyclo[3.3.1]nonane (9-BBN) to methoxyborane MeO-BBN (Table 1, Entry 1). Indeed a Turn Over Number (TON) lower than 5 was observed after 28 h at 25 °C. Increasing the reaction temperature to 70 °C did not improve the catalytic performances of **1** (Entry 11 in Table 1). These results are in line with the results of Stephan *et al.* on the phosphine catalysed hydroboration of CO₂.^{7b} Similarly, the borane **2-BMes₂** is inefficient both at 25 and 70 °C (Entries 2 and 12 in Table 1). In stark contrast, a mixture of **1** and **2-BMes₂** provides a competent catalytic system, enabling the reduction of CO₂ to MeOBBN with a TON of 41 after 28 h at 25 °C (TOF = 1.5 h⁻¹). This result demonstrates for the first time the synergistic and positive influence of combining a Lewis base and a Lewis acid in the reduction of CO₂ with hydroboranes. Replacing **2-BMes₂** with the less acidic **2-Bpin** borane somewhat deactivates the catalytic system and a TON of 36 (TOF = 1.3 h⁻¹) is measured under the same conditions. To further establish the benefit of an ambiphilic structure, the catalytic potential of the intramolecular Lewis pairs **3-BMes₂** and **3-Bpin** has been determined. **3-BMes₂** presents a better activity than the mixture of **1** and **2-BMes₂** and a TON of 100 was reached after 28 hours at 25 °C (TOF = 3.6 h⁻¹).

It is worth noting here, that the reaction course follows a profile similar to the one observed previously for guanidines and N/B FLPs.^{6f} Indeed, monitoring the product distribution over time, using ¹H and ¹³C NMR spectroscopy, reveals that CO₂ is first reduced to the boryl formate HCOO-BBN which is present in only a very low concentration and does not accumulate (Fig. S1). This intermediate is subsequently reduced to the acetal form H₂C(OBBN)₂, with a conversion rate of 90% after just 1 hour. Finally, reduction of the acetal provides MeO-BBN. This step is slow and requires 27 hours at 25 °C to yield the methoxyborane species quantitatively. The kinetics of the reaction improved at 70 °C. These conditions revealed a remarkable catalytic activity of **3-BMes₂**, allowing the full conversion of CO₂ to the corresponding methoxyborane with a TON of 100, in only 2 hours (Table 1, entry 7). Reducing the loading of catalyst **3-BMes₂** to 0.1 and 0.05 mol%, afforded an excellent TON of 1980, after 20 h at 70 °C, and a TOF of 250 h⁻¹, while the isolated components of the catalytic system, namely **1** and **2-BMes₂**, are unreactive (Table 1, Entries 9-12).

Table 1. Catalytic reduction of CO₂ to methoxyborane CH₃OBBN

catalysts:

	1	2-BMes₂	2-Bpin	3-BMes₂	3-Bpin
X	PPh ₂	H	H	PPh ₂	PPh ₂
Y	H	BMes ₂	Bpin	BMes ₂	Bpin

Entry ^a	Catalyst (mol%)	T °C	Time (h)	TON ^b	TOF (h ⁻¹) ^b
1	1 (1 mol%)	25 °C	28h	<5	<0.2
2	2-BMes₂ (1 mol%)	25 °C	28h	0	0
3	3-Bpin (1 mol%)	25 °C	28h	84	3.0
4	3-BMes₂ (1 mol%)	25 °C	28h	100	3.6
5	1 (1 mol%) + 2-Bpin (1 mol%)	25 °C	28h	36	1.3
6	1 (1 mol%) + 2-BMes₂ (1 mol%)	25 °C	28h	41	1.5
7	3-BMes₂ (1 mol%)	70 °C	2h	100	50
8	3-Bpin (1 mol%)	70 °C	2h	57	29
9	3-BMes₂ (0.1 mol%)	70 °C	4h	1000	250
10	3-BMes₂ (0.05 mol%)	70 °C	20h	1980	99
11	1 (1 mol%)	70 °C	2h	<2	<1
12	2-BMes₂ (1 mol%)	70 °C	2h	0	0


[a] reactions performed in J-Young tubes; monitored by NMR spectroscopy with 0.0048 mmol of catalyst, 0.48 mmol of 9-BBN in *d₈*-THF (0.40 mL) and exposed to an atmosphere of CO₂ (1 bar), [b] The TON is obtained by ¹H NMR monitoring (CH₃O-BBN formation), with mesitylene as internal standard.

Dialkylboranes, other than 9-BBN, were also tested as potential reductants. As expected, the hydroboration of CO₂ with catecholborane is slower than with 9-BBN because of the lower Lewis acidity of the borane. In fact, under the optimized reaction conditions (48 h at 70 °C), **3-BMes₂** promotes the reduction of CO₂ to MeOBBN with a modest TOF of 23. Replacing catecholborane with BH₃SMe₂ did not lead to any significant product formation.

Again, replacing the BMes₂ unit with a Bpin fragment somewhat slows down the catalytic reaction and a TOF of 3.0 h⁻¹ is measured with **3-Bpin**, at 25 °C (vs 3.6 h⁻¹ for BMes₂). The difference of reactivity between **3-BMes₂** and **3-Bpin** is more pronounced at 70 °C, with TOFs values of 50 and 29 h⁻¹, respectively (Entries 7-8 in Table 1). This behaviour directly follows from the Arrhenius rate law.

Previous studies by the groups of Stephan and Cantat have established that basic guanidines, N-heterocyclic carbenes and phosphines can promote the hydroboration of CO₂.^{6f, 11} Mechanistic and computational investigations have shown that basic catalysts proceed through the activation of the B-H bond, by coordination of the Lewis base to the hydroborane.^{6f} Therefore we propose here that the benefit of combining a trivalent phosphorus function with a remote Lewis acid, as in **3-BMes₂**, results from a concomitant H-transfer / borane-formate adduct formation, in the first step of the CO₂ reduction process (Scheme 3). In this approach, linking together the Lewis acid and the Lewis base functions should reduce the volume of the non-productive region of the conformational space between the catalysts

and the substrates and thus increase further the reaction rates.¹² In fact, experimentally, catalysts **3-BMes₂** and **3-Bpin** are about twice as reactive as the corresponding bimolecular systems at 25 °C. This difference of reactivity corresponds roughly to a decrease of ca. 1 kcal/mol in activation energy, consistent with an entropic effect.



Scheme 3. Proposed synergistic effect of phosphino-borane catalysts in the hydroboration of CO₂

Overall, we have demonstrated for the first time that the combination of phosphines and boranes offers a synergistic effect in the catalytic reduction of CO₂ with hydroboranes. While the phosphine and the borane fragments, taken separately, do not catalyse the reduction of CO₂, the P/B derivatives **3**, combining the two functions, display high catalytic activity.

Notes and references

^a NIMBE, CEA, CNRS, Université Paris-Saclay, Gif-sur-Yvette, France

E-mail: thibault.cantat@cea.fr.

Fax: (+): (+33) 1 6908 6640

Homepage: <http://iramis.cea.fr/Pisp/thibault.cantat/index.html>.

^b Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-Sud, Université Paris-Saclay, 1 av. de la Terrasse, 91198 Gif-sur-Yvette (France)

E-mail: arnaud.voituriez@cnrs.fr.

angela.marinetti@cnrs.fr.

Acknowledgements

We thank the CHARMMMAT Laboratory of Excellence, CEA, CNRS, Institut de Chimie des Substances Naturelles, and the ANR for financial support. The European Research Council is acknowledged for an ERC Starting Grant to T.C (Grant Agreement no. 336467).

† Footnotes should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.

Electronic Supplementary Information (ESI) available: Fig. S1 and experimental procedures. See DOI: 10.1039/c000000x/

1 a) M. Aresta, *Carbon Dioxide as Chemical Feedstock*, Wiley-VCH, Weinheim, 2010; b) P. G. Jessop, T. Ikariya and R. Noyori, *Chem. Rev.*, 1995, **95**, 259; c) K. Huang, C. L. Sun and Z. J. Shi, *Chem. Soc. Rev.*, 2011, **40**, 2435; d) C. Federsel, A. Boddien, R. Jackstell, R. Jennerjahn, P. J. Dyson, R. Scopelliti, G. Laurenczy and M. Beller, *Angew. Chem. Int. Ed.*, 2010, **49**, 9777; e) C. Federsel, R. Jackstell and M. Beller, *Angew. Chem. Int. Ed.*, 2010, **49**, 6254; f) A. Tlili, X. Frogneux, E. Blondiaux and T. Cantat, *Angew. Chem. Int. Ed.*, 2014, **53**,

- 2543; g) A. Tlili, E. Blondiaux, X. Frogneux, and T. Cantat, *Green Chem.*, 2015, **17**, 157; h) C. A. Huff and M. S. Sanford, *J. Am. Chem. Soc.*, 2011, **133**, 18122; i) K. Tominaga, Y. Sasaki, M. Kawai, T. Watanabe and M. Saito, *Chem. Commun.*, 1993, 629; j) S. Wesselbaum, T. V. Stein, J. Klankermayer and W. Leitner, *Angew. Chem. Int. Ed.*, 2012, **51**, 7499; k) S. Savourey, G. Lefèvre, J.-C. Berthet, P. Thuéry, C. Genre and T. Cantat, *Angew. Chem. Int. Ed.*, 2014, **53**, 10466; l) S. Wesselbaum, V. Moha, M. Meuresch, S. Brosinski, K. M. Thenert, J. Kothe, T. vom Stein, U. Englert, M. Hölscher, J. Klankermayer and W. Leitner, *Chem. Sci.*, 2015, **6**, 693; m) S. Bontemps, *Coord. Chem. Rev.*, 2016, **308**, 117.
- 2 *CRC Handbook of Chemistry and Physics*, 92th ed., Ed. W. M. Haynes, D. R. Lide, The Chemical Rubber Co., 2011.
- 3 For a general recent review see: C. C. Chong and R. Kinjo, *ACS Catal.*, 2015, **5**, 3238.
- 4 S. Chakraborty, J. Zhang, J. A. Krause, and H. Guan, *J. Am. Chem. Soc.*, 2010, **132**, 8872.
- 5 a) S. Bontemps, L. Vendier and S. Sabo-Etienne, *Angew. Chem. Int. Ed.*, 2012, **51**, 1671; b) S. Bontemps, and S. Sabo-Etienne, *Angew. Chem. Int. Ed.*, 2013, **52**, 10253.
- 6 a) M.-A. Courtemanche, M.-A. Légaré, L. Maron, and F.-G. Fontaine, *J. Am. Chem. Soc.*, 2013, **135**, 9326; b) M.-A. Courtemanche, M.-A. Légaré, L. Maron, and F.-G. Fontaine, *J. Am. Chem. Soc.*, 2014, **136**, 10708; c) F.-G. Fontaine, M.-A. Courtemanche, and M.-A. Légaré, *Chem. Eur. J.*, 2014, **11**, 2990; d) M.-A. Légaré, M.-A. Courtemanche and F.-G. Fontaine, *Chem. Commun.*, 2014, **50**, 11362; e) R. Declercq, G. Bouhadir, D. Bourissou, M.-A. Légaré, M.-A. Courtemanche, K. S. Nahi, N. Bouchard, F.-G. Fontaine, and L. Maron, *ACS Catal.*, 2015, **5**, 2513; f) C. Das Neves Gomes, E. Blondiaux, P. Thuéry and T. Cantat, *Chem. Eur. J.*, 2014, **20**, 7098; g) T. Cantat, C. Gomes, E. Blondiaux, O. Jacquet, patent WO2014162266 filed April 3, 2013.
- 7 a) E. Blondiaux, J. Pouessel and T. Cantat, *Angew. Chem. Int. Ed.* 2014, **53**, 12186; b) T. Wang and D. W. Stephan, *Chem. Commun.*, 2014, **50**, 7007; c) S. Y.-F. Ho, C.-W. So, N. Saffon-Merceron and N. Mézailles, *Chem. Commun.*, 2015, **51**, 2107.
- 8 a) M. W. P. Bebbington, S. Bontemps, G. Bouhadir, M. J. Hanton, R. P. Tooze, H. Van Rensburg and D. Bourissou, *New J. Chem.*, 2010, **34**, 1556. For some applications of phosphino-borane compounds in catalysis, see: b) S. Bontemps, G. Bouhadir, K. Miqueu, D. Bourissou *J. Am. Chem. Soc.* 2006, **128**, 12056; c) S. Porcel, G. Bouhadir, N. Saffon, L. Maron, D. Bourissou, *Angew. Chem., Int. Ed.*, 2010, **49**, 6186. For other phosphino-boranes based on a ferrocene backbone, see: a) I. R. Morgan, A. Di Paolo, D. Vidovic, I. A. Fallis and S. Aldridge, *Chem. Commun.*, 2009, 7288. b) X. Wang, G. Kehr, C.G. Daniliuc, and G. Erker, *J. Am. Chem. Soc.*, 2014, **136**, 3293.
- 9 a) D. A. Khobragade, S. G. Mahamulkar, L. Pospíšil, I. Cisařová, L. Rulíšek, U. Jahn, *Chem. Eur. J.*, 2012, **18**, 12267; b) R. Sanders, U. T. Mueller-Westerhoff, *J. Organomet. Chem.*, 1996, **512**, 219.
- 10 Similarly, no adduct forms when **3-BMes₂** is reacted with a stoichiometric amount of 9-BBN.
- 11 T. Wang and D. W. Stephan, *Chem. Eur. J.*, 2014, **20**, 3036.
- 12 S. Milstien and L. Cohen, *J. Am. Chem. Soc.*, 1972, **94**, 9158.