

Local velocity measurements in the Shrek experiment at high reynolds number

Christophe Baudet, Michel Bon-Mardion, Patrick Bonnay, A. Braslau, Bernard Castaing, Francesca Chillà, Laurent Chevillard, François Daviaud, Pantxo Diribarne, Bérengère Dubrulle, et al.

► To cite this version:

Christophe Baudet, Michel Bon-Mardion, Patrick Bonnay, A. Braslau, Bernard Castaing, et al.. Local velocity measurements in the Shrek experiment at high reynolds number. 15th European Turbulence Conference, Delft University of Technology (TU DELFT), Aug 2015, Delft, Netherlands. cea-01370530

HAL Id: cea-01370530

<https://hal-cea.archives-ouvertes.fr/cea-01370530>

Submitted on 22 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LOCAL VELOCITY MEASUREMENTS IN THE SHREK EXPERIMENT AT HIGH REYNOLDS NUMBER

C. Baudet^{1,2}, M. Bon Mardion^{3,4}, P. Bonnay^{3,4}, A. Braslau⁵, B. Castaing⁶, F. Chillà⁶, L. Chevillard⁶, F. Daviaud⁵, P. Diribarne^{3,4}, B. Dubrulle⁵, D. Duri^{1,2,3,4}, D. Faranda⁵, B. Gallet⁵, M. Gibert^{7,8}, A. Girard^{3,4}, B. Hébral^{7,8}, T. Lehner¹⁰, I. Moukharski⁵, J.M. Poncet^{3,4}, J.-P. Moro⁹, P.-E. Roche^{7,8}, B. Rousset^{3,4}, E. Rusaouën^{7,8}, B. Saint-Michel⁵, J. Salort⁶, E-W. Saw⁵, K. Steiros⁵, C. Wiertel-Gasquet⁵ (SHREK collaboration)

¹Univ. Grenoble Alpes, LEGI, F-38041 Grenoble, France

²CNRS, LEGI, F-38041 Grenoble, France

³Univ. Grenoble Alpes, INAC-SBT, F-38000 Grenoble, France

⁴CEA, INAC-SBT, F-38000 Grenoble, France

⁵CEA, IRAMIS, SPEC, CNRS UMR 3680, Laboratoire SPHYNX, 91191 Gif-sur-Yvette, France

⁶Laboratoire de Physique de l'ÉNS de Lyon, CNRS/Université Lyon - F-69364 Lyon cedex 7, France

⁷Univ. Grenoble Alpes, Institut NEEL, F-38042 Grenoble, France

⁸CNRS, Institut NEEL, F-38042 Grenoble, France

⁹CEA, DEN-DANS-DM2S-STMF-LIEFT, F-38000 Grenoble, France

¹⁰LUTH, Observatoire de Paris, CNRS, France

Abstract We report preliminary results obtained using new local velocity probes in the Superfluid Helium high REynold number von Kármán flow (SHREK) experiment for different forcing conditions. The presentation will focus on the validation of the signals obtained from a hot-wire and a total head pressure tube in both normal and superfluid phases of liquid helium.

The SHREK experiment (SuperFluid von Kármán experiment) has been designed to compare properties of superfluid turbulence with ordinary turbulence. The SHREK experiment (see Fig. 1) makes particularly use of the phase transition experienced by helium at $T_\lambda \approx 2.2$ K, thus allowing to make experiments at very high Reynolds values with normal (above T_λ) and superfluid helium (below T_λ) [3, 7]. The experimental setup is described in [4]. It consists of a cylinder of radius $R = 0.39$ m and height $H = 1.2$ m (Fig. 1). The fluid is mechanically stirred by a pair of coaxial impellers. The impellers are disks of radius $0.925R$, fitted with 8 radial blades of height $0.2R$ and curvature radius $0.4625R$. The impellers rotation rate f_1 and f_2 can be varied independently from 0.1 to 2 Hz and the total dissipated power can range from 120W at 1.6 K to 400W above 1.9 K. This experiment benefits from high flexibility of flow conditions, due to the large variation of helium properties over the available temperature range (1.6 K to 5 K). Both superfluid and normal turbulence measurements are possible in the same experiment, with adjustable fraction of the superfluid component (from $\approx 85\%$ at 1.6 K to 0% above T_λ).




Figure 1. Schematic view of the experimental setup and the impellers blade profile

Besides global measurements[6] (torques and calorimetry), the experiment was designed to enable local measurements in the equatorial plane or next to the impellers. In the sequel, we present two new sensors, a Platinum-Rhodium hot wire

and a new design of miniature Pitot anemometers, that were successfully operated in SHREK experiment to obtain local velocity measurements.




Figure 2. Inter-correlation coefficient of a Pitot tube and a hot-wire signals obtained in He I (red diamonds) and He II (blue circles). Acquisitions performed in HeJet experiment [1]

First we bring new insight in the understanding of the signal of hot-wires in He II (see [2]). In particular, we show that the inter-correlation coefficient of the signal of a hot-wire with the signal of a Pitot tube (for which the design can be found in [7]), is the same in He II as in He I where the signal can reliably be interpreted as a velocity (see Fig. 2). We also show preliminary velocity measurements obtained in SHREK using this kind of hot-wire.

Second, we report the first results obtained using novel miniature total head pressure tube (Pitot tube), obtained for a large range of rotation frequencies f_1 and f_2 , in both normal and superfluid regime. The Pitot Helmholtz frequency is ≈ 450 Hz giving useful bandwidth somewhat higher. The sensitivity is in the $10^{-7} m^2/s^2$ range and can be further improved by combination of 2-amplifiers. The results are compared with results obtained in a scale 1:4 experiment, operated in Saclay with water [5].

Acknowledgment

This work was supported by the French National Research Agency grant ANR-09-BLAN-0094-01 and by the European Community Framework Programme 7, EuHIT - European High-performance Infrastructures in Turbulence, grant agreement no. 312778.

References

- [1] D. Duri, C. Baudet, P. Charvin, J. Virone, B. Rousset, J.M. Poncet, and P. Diribarne. Liquid helium inertial jet for comparative study of classical and quantum turbulence. *Review of Scientific Instruments*, **82**, 2011.
- [2] D. Duri, C. Baudet, J.-P. Moro, P.-E. Roche, and P. Diribarne. Heat transfer enhancement on thin wires in superfluid helium forced flows. *arXiv preprint arXiv:1410.0167*, 2014.
- [3] J. Maurer and P. Tabeling. Local investigation of superfluid turbulence. *Europhys. Lett.*, **43** (1):29, 1998.
- [4] B Rousset, P Bonnay, P Diribarne, A Girard, JM Poncet, E Herbert, J Salort, C Baudet, B Castaing, L Chevillard, et al. Superfluid high reynolds von kármán experiment. *Review of Scientific Instruments*, **85**(10):103908, 2014.
- [5] B Saint-Michel, B Dubrulle, L Marié, F Ravelet, and F Daviaud. Influence of reynolds number and forcing type in a turbulent von kármán flow. *New Journal of Physics*, **16**(6):063037, 2014.
- [6] B Saint-Michel, E Herbert, J Salort, C Baudet, M Bon Mardion, P Bonnay, M Bourgoïn, B Castaing, L Chevillard, F Daviaud, et al. Probing quantum and classical turbulence analogy in von kármán liquid helium, nitrogen, and water experiments. *Physics of Fluids (1994-present)*, **26**(12):125109, 2014.
- [7] J Salort, C Baudet, B Castaing, B Chabaud, F Daviaud, T Didelot, P Diribarne, B Dubrulle, Y Gagne, F Gauthier, et al. Turbulent velocity spectra in superfluid flows. *Physics of Fluids*, **22**(12):5102, 2010.