

Resumming large higher-order corrections in non-linear QCD evolution

E. Iancu, J. D. Madrigal, A. H. Mueller, Gregory Soyez, D. N. Triantafyllopoulos

► **To cite this version:**

E. Iancu, J. D. Madrigal, A. H. Mueller, Gregory Soyez, D. N. Triantafyllopoulos. Resumming large higher-order corrections in non-linear QCD evolution. Nuclear Physics B Proceedings Supplement, 2015, 2016, pp.046. <10.1016/j.nuclphysbps.2016.05.046>. <cea-01465826>

HAL Id: cea-01465826

<https://hal-cea.archives-ouvertes.fr/cea-01465826>

Submitted on 13 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Resumming large higher-order corrections in non-linear QCD evolution

E. Iancu^a, J.D. Madrigal^a, A.H. Mueller^b, G. Soyez^a, D.N. Triantafyllopoulos^{c,*}

^a*Institut de Physique Théorique, CEA Saclay, CNRS UMR 3681, F-91191 Gif-sur-Yvette, France*

^b*Department of Physics, Columbia University, New York, NY 10027, USA*

^c*European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT*)
and Fondazione Bruno Kessler, Strada delle Tabarelle 286, I-38123 Villazzano (TN), Italy*

Abstract

Linear and non-linear QCD evolutions at high energy suffer from severe issues related to convergence, due to higher order corrections enhanced by large double and single transverse logarithms. We resum double logarithms to all orders by taking into account successive soft gluon emissions strongly ordered in lifetime. We further resum single logarithms generated by the first non-singular part of the splitting functions and by the one-loop running of the coupling. The resulting collinearly improved BK equation admits stable solutions, which are used to successfully fit the HERA data at small- x for physically acceptable initial conditions and reasonable values of the fit parameters.

1. Introduction

The Color Glass Condensate (CGC) [1] is an effective theory for the soft, small- x , components of a fast moving hadron. It predicts the saturation of gluon occupation numbers to values of order $\mathcal{O}(1/\bar{\alpha})$, where $\bar{\alpha} = \alpha_s N_c / \pi$ with N_c the number of colors. The most convenient degrees of freedom are Wilson lines for the scattering of projectile partons off the hadron, and gauge invariant combinations of these Wilson lines satisfy unitarity constraints as a result of gluon saturation. These correlators are the building blocks for calculating many observables local in rapidity, like total cross sections in deep inelastic scattering, single and multi-particle production in pA and dA collisions, energy density and its fluctuations in the first stage after an AA collision etc. To an excellent accuracy, such correlators can be expressed in terms of the dipole scattering amplitude $T_{12} = 1 - (1/N_c) \text{tr}(V_1^\dagger V_2)$ [2–4]. The indices stand for the dependence on the 2-dimensional vectors z_1 and z_2 , which are the transverse coordinates of the quark and the antiquark composing the dipole. The amplitude T_{12}

is small in the region where the target is dilute and approaches unity for a dense target. The saturation momentum $Q_s(Y)$ controls the separation between these two regimes, and increases with the rapidity difference Y between the projectile dipole and the hadronic target.

2. The BK equation at NLO

The S -matrix $S_{12} = 1 - T_{12}$ evolves with Y according to the Balitsky-Kovchegov (BK) equation [5, 6]. Typical diagrams contributing to the LO and the NLO evolution involve one or two parton emissions respectively, are shown in Fig. 1, and the corresponding equation written in a form suited to our purposes reads [7–9]

$$\begin{aligned} \frac{dS_{12}}{dY} = & \frac{\bar{\alpha}}{2\pi} \int d^2z_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left[1 + \bar{\alpha} \bar{b} \left(\ln z_{12}^2 \mu^2 - \frac{z_{13}^2 - z_{23}^2}{z_{12}^2} \ln \frac{z_{13}^2}{z_{23}^2} \right) \right. \\ & \left. + \bar{\alpha} \left(\frac{67}{36} - \frac{\pi^2}{12} - \frac{5}{18} \frac{N_f}{N_c} - \frac{1}{2} \ln \frac{z_{13}^2}{z_{12}^2} \ln \frac{z_{23}^2}{z_{12}^2} \right) \right] (S_{13} S_{32} - S_{12}) \\ & + \frac{\bar{\alpha}^2}{8\pi^2} \int \frac{d^2z_3 d^2z_4}{z_{34}^4} \left[-2 + \frac{z_{13}^2 z_{24}^2 + z_{14}^2 z_{23}^2 - 4z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2 - z_{14}^2 z_{23}^2} \right. \\ & \left. \ln \frac{z_{13}^2 z_{24}^2}{z_{14}^2 z_{23}^2} + \frac{z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2} \left(1 + \frac{z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2 - z_{14}^2 z_{23}^2} \right) \ln \frac{z_{13}^2 z_{24}^2}{z_{14}^2 z_{23}^2} \right] \\ & (S_{13} S_{34} S_{42} - S_{13} S_{32}), \end{aligned} \quad (1)$$

*Corresponding author

Email address: trianta@ectstar.eu

(D.N. Triantafyllopoulos)




Figure 1: Typical diagrams contributing to the BK equation. The thick vertical line stands for the hadron. Left: LO terms. Middle: NLO terms proportional to N_f . Right: NLO terms proportional to N_c .

where we have neglected $1/N_c^2$ suppressed terms¹. In Eq. (1) we have defined $z_{ij} = z_i - z_j$, with z_3 and z_4 the transverse coordinates of the daughter partons, $\bar{b} = (11N_c - 2N_f)/12N_c$ is the first coefficient of the QCD β -function, and μ is a renormalization scale at which the coupling is evaluated.

3. Large transverse logarithms

In general, a NLO computation is expected to add a small $O(\bar{\alpha})$ correction to the LO result. However, this is not the case here since there are terms in the NLO kernel which get large in certain kinematic domains and invalidate the strict expansion in $\bar{\alpha}$. These are collinear logarithms, i.e. logarithms associated with very disparate transverse dipole sizes between successive emissions. More precisely, let's consider the regime

$$1/Q_s \gg z_{14} \simeq z_{24} \simeq z_{34} \gg z_{13} \simeq z_{23} \gg z_{12}, \quad (2)$$

that is, the parent dipole is the smallest one, a gluon is emitted far away at z_3 and another one even further at z_4 , while all dipoles remain small so that the scattering is weak. Then the dominant NLO contribution involves the double transverse logarithm (DTL) in the single integration term in Eq. (1). Moreover, a careful expansion of the kernel in the double integration term, reveals that there is also a single transverse logarithm (STL). Linearizing in the amplitude T , assuming that the latter depends only on the magnitude of z_{ij} , relabelling some variables and putting everything together, we find that

¹We have further ignored a term proportional to N_f/N_c , with N_f the number of flavors, but which becomes of order $O(N_f/N_c^3)$ in the weak scattering regime.

Eq. (1) in the collinear region becomes [10, 11]

$$\frac{dT_{12}}{dY} = \bar{\alpha} \int_{z_{12}^2}^{1/Q_s^2} dz_{13}^2 \frac{z_{12}^2}{z_{13}^4} \left(1 - \frac{1}{2} \bar{\alpha} \ln^2 \frac{z_{13}^2}{z_{12}^2} - \frac{11}{12} \bar{\alpha} \ln \frac{z_{13}^2}{z_{12}^2} \right) T_{13}. \quad (3)$$

To arrive at the above, we have dropped virtual terms, i.e. terms proportional to T_{12} , since largest dipoles scatter stronger. Now it becomes evident that for large enough daughter dipole size z_{13} , the NLO contribution can be comparable to, or even larger than, the LO one due to the presence of a large DTL and/or STL. E.g. consider the simple, but realistic initial condition

$$T_{12} = \begin{cases} z_{12}^2 Q_s^2 & \text{for } z_{12}^2 Q_s^2 \ll 1 \\ 1 & \text{for } z_{12}^2 Q_s^2 \gg 1. \end{cases} \quad (4)$$

Then the transverse integration in Eq. (3) becomes logarithmic and the one-step in rapidity evolution gives

$$\Delta T_{12} = \bar{\alpha} \Delta Y z_{12}^2 Q_s^2 \ln \frac{1}{z_{12}^2 Q_s^2} \left(1 - \frac{1}{6} \bar{\alpha} \ln^2 \frac{1}{z_{12}^2 Q_s^2} - \frac{11}{24} \bar{\alpha} \ln \frac{1}{z_{12}^2 Q_s^2} \right). \quad (5)$$

Therefore, for sufficiently small z_{12} the perturbation series becomes unreliable and even more, since the large NLO correction is negative, the solution is unstable. This is indeed what has been observed in numerical solutions [10, 12, 13] and is demonstrated in Fig. 2.

4. Origin of the large logs and their resummation

Understanding the source of the large logarithms is necessary before trying to resum them to all orders. The DTL arises from diagrams in which the two successive




Figure 2: Unstable solution to the NLO BK equation [10]. Only the DTL of the NLO kernel is kept and the coupling is fixed at $\bar{\alpha} = 0.25$.

gluon emissions are strongly ordered in light-cone time x^+ , thus its origin is in the kinematics. For example, let us examine the diagram in Fig. 3 where we take $\mathbf{k} \ll \mathbf{p}$, $\tilde{\mathbf{k}} \ll \tilde{\mathbf{p}}$ and $k^+ \ll p^+$. Inspection of energy denominators of the type $(k^- + p^-)^{-1} = [(2k^2/k^+) + (2p^2/p^+)]^{-1} \equiv (1/\tau_k + 1/\tau_p)^{-1}$ which appear in light-cone perturbation theory, shows that the largest logarithmic terms occur when the lifetimes (or equivalently the light-cone energies [14]) of the two gluons are also strongly ordered, i.e. when $\tau_k \ll \tau_p$. Different hookings of the two gluons lead to 32 diagrams like the one in Fig. 3. Adding all these contributions and integrating over transverse momenta we find in the collinear regime

$$\Delta T_{12} = \bar{\alpha}^2 \int \frac{dp^+}{p^+} \frac{dk^+}{k^+} \Theta \left(p^+ \frac{z_{13}^2}{z_{14}^2} - k^+ \right) dz_{13}^2 dz_{14}^2 \frac{z_{12}^2}{z_{13}^2 z_{14}^2} T_{14}. \quad (6)$$

The step-function is the key element and arises by encoding the lifetime constraint. By further doing the longitudinal integrations and the one over the intermediate dipole size z_{13} , we obtain the part with the DTL in Eq. (3)². Thus, in order to resum the DTLs to all orders, we need to take into account all the diagrams with an arbitrary number of gluons emissions, in which the gluons are not only strongly ordered in their transverse and longitudinal momenta, but also in their lifetimes.

The STL also arises from successive emissions in which the second gluon is much softer, both in transverse and longitudinal momenta, than the first one, but now it is the region $\tau_k \sim \tau_p$ which gives the relevant contribution. Not surprisingly, the STL is of dynamical origin and is related to DGLAP evolution. This is supported by the fact that its coefficient $A_1 = 11/12$ is the first non-singular term in the small- ω expansion of the

²Other diagrams which are not time-ordered may contain double logarithms individually, but they cancel in the final answer [10].




Figure 3: Typical diagram leading to a large DTL. The softer gluon k is emitted after and absorbed before the harder one p .

relevant anomalous dimension, that is

$$P_T(\omega) = \int_0^1 dz z^\omega \left[P_{\text{gg}}(z) + \frac{C_F}{N_c} P_{\text{qg}}(z) \right] = \frac{1}{\omega} - A_1 + \dots \quad (7)$$

This suggests that in order to resum such STLs, it suffices to include A_1 as an anomalous dimension, i.e. as a power-law suppression in the evolution kernel.

Putting everything together, we arrive at the non-linear evolution equation [10, 11]

$$\frac{d\tilde{T}_{12}}{dY} = \frac{\bar{\alpha}}{2\pi} \int d^2 z_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left(\frac{z_{12}^2}{z_c^2} \right)^{\pm A_1 \bar{\alpha}} \frac{J_1(2\sqrt{\bar{\alpha}} L_{13} L_{23})}{\sqrt{\bar{\alpha}} L_{13} L_{23}} (\tilde{T}_{13} + \tilde{T}_{23} - \tilde{T}_{12} - \tilde{T}_{13} \tilde{T}_{23}), \quad (8)$$

which is of the LO BK type, but with two extra factors in the kernel that resum the STLs and DTLs³. Clearly, when such factors are expanded in the second non-trivial order in $\bar{\alpha}$ they reproduce the large logarithmic terms of the NLO BK equation as presented in Eq. (3). In Eq. (8) we have defined $L_{13} \equiv \ln(z_{13}^2/z_{12}^2)$ and similarly for L_{23} , while $z_c = \min\{z_{13}, z_{23}\}$. The positive sign in the power-law suppression is to be used in the regime in Eq. (2), while the minus sign is taken when $z_c \ll z_{12}$ and will resum STLs arising from successive gluon emissions which are still softer in longitudinal momentum but harder in the transverse space. Finally, one should note that \tilde{T}_{12} in Eq. (8) coincides with the physical amplitude T_{12} only for $Y \geq \ln(1/z_{12}^2 Q_0^2)$, with Q_0 the typical target scale, which is anyway the region in which high energy evolution can be trusted⁴.

5. Running coupling

There is one last source of potentially large NLO terms in the NLO BK equation. These are the logarithmic terms proportional to \bar{b} in Eq. (1) and they can

³So long as the DTLs are considered, a presumably equivalent to the order of accuracy, but non-local in Y , equation has been derived in [14]. In momentum space and at the linear level, the Bessel kernel has first appeared in [15].

⁴Moreover, the initial condition in Eq. (8) has to be modified in order to include large transverse logarithms in the impact factor [10].

get large when the scales in their arguments become very disparate. The scale μ , which is the same one at which the coupling is to be evaluated in Eqs. (1) and (8), must be chosen in a way to cancel such large logarithms. Even though the choice is not unique, it is the hardest scale which determines the running. Indeed, the smallest dipole prescription

$$\bar{\alpha}_{\min} = \bar{\alpha}(z_{\min}) \quad \text{with} \quad z_{\min} = \min\{z_{12}, z_{13}, z_{23}\}, \quad (9)$$

cancels the large logarithms in all dangerous regions. Another option is to choose μ so that all $\bar{\alpha}^2 \bar{b}$ terms vanish. Since in the present work we also neglect all pure $\mathcal{O}(\bar{\alpha}^2)$ terms, such a choice looks like the *fastest apparent convergence* scheme. One easily finds

$$\bar{\alpha}_{\text{fac}} = \left(\frac{1}{\bar{\alpha}_{12}} + \frac{z_{13}^2 - z_{23}^2}{z_{12}^2} \frac{\bar{\alpha}_{13} - \bar{\alpha}_{23}}{\bar{\alpha}_{13}\bar{\alpha}_{23}} \right)^{-1}, \quad (10)$$

which is equivalent to $\bar{\alpha}_{\min}$ in the extreme kinematical cases where one of the three dipoles is much smaller than the other two. We shall adopt one of the above two schemes, but we shall not consider the one introduced in [7] and which has been extensively used so far in phenomenology, as it is plagued by certain peculiarities.

6. Solutions, applications and outlook

Contrary to what happens to the NLO BK equation in (1), the resummed one in Eq. (8) admits well-defined solutions, whether the coupling is fixed or it runs. Such a stable solution is shown in Fig. 4. The DTLs, the STLs and the running of the coupling individually suppress the evolution and hence the final cumulative “speed” is substantially suppressed when compared to the one of the LO evolution.

Given appropriate forms for the initial condition, in [11] we have used the numerical solutions to the resummed BK equation in order to perform fits to the HERA data [16] for the ep reduced cross section (for similar fits, without inclusion of the STLs, see [17]). Given the range of validity for small- x dynamics, we have restricted ourselves in the region $x \leq 10^{-2}$ and $Q^2 \leq Q_{\max}^2$, where the upper limit Q_{\max}^2 in the photon virtuality has been varied even up to 400 GeV². With 4 free parameters we obtain good fits with a χ^2 per point around 1.1–1.2. They are also rather discriminatory: they favor (i) the running coupling version of the McLerran-Venugopalan (MV) model as an initial condition and (ii) the two running coupling prescriptions $\bar{\alpha}_{\min}$ and $\bar{\alpha}_{\text{fac}}$, with a slight preference to the former. Whether or not we include the power-law suppression factor due




Figure 4: Stable solution to the resummed BK equation (8).

the resummation of STLs in Eq. (8), does not affect the quality of the fits, however it does lead to more physical values of the fit parameters. For completeness, let us point out that the evolution speed extracted from such fits is $\lambda_s \equiv d \ln Q_s^2 / dY = 0.20 \div 0.24$.

It would be desirable to generalize Eq. (8) to include the pure $\mathcal{O}(\bar{\alpha}^2)$ terms, but this may not change much in the phenomenology, since many of such contributions are taken into account by one of the fit parameters.

References

- [1] F. Gelis, E. Iancu, J. Jalilian-Marian, R. Venugopalan, *Ann.Rev.Nucl.Part.Sci.* 60 (2010) 463, arXiv:1002.0333.
- [2] A. Dumitru, J. Jalilian-Marian, T. Lappi, B. Schenke, R. Venugopalan, *Phys.Lett.* B706 (2011) 219, arXiv:1108.4764.
- [3] E. Iancu, D.N. Triantafyllopoulos, *JHEP* 1111 (2011) 105, arXiv:1109.0302.
- [4] E. Iancu, D.N. Triantafyllopoulos, *JHEP* 1204 (2012) 025, arXiv:1112.1104.
- [5] I. Balitsky, *Nucl.Phys.* B463 (1996) 99, arXiv:hep-ph/9509348.
- [6] Y.V. Kovchegov, *Phys.Rev.* D60 (1999) 034008, arXiv:hep-ph/9901281.
- [7] I. Balitsky, *Phys.Rev.* D75 (2007) 014001, arXiv:hep-ph/0609105.
- [8] Y. V. Kovchegov, H. Weigert, *Nucl.Phys.* A784 (2007) 188, arXiv:hep-ph/0609090.
- [9] I. Balitsky and G.A. Chirilli, *Phys.Rev.* D77 (2008) 014019, arXiv:0710.4330.
- [10] E. Iancu, J.D. Madrigal, A.H. Mueller, G. Soyez, D.N. Triantafyllopoulos, *Phys.Lett.* B744 (2015) 293, arXiv:1502.05642.
- [11] E. Iancu, J.D. Madrigal, A.H. Mueller, G. Soyez, D.N. Triantafyllopoulos, arXiv:1507.03651.
- [12] E. Avsar, A. Stasto, D.N. Triantafyllopoulos, D. Zaslavsky, *JHEP* 1110 (2011) 138, arXiv:1107.1252.
- [13] T. Lappi, H. Mäntysaari, *Phys.Rev.* D91 (2015) 074016, arXiv:1502.02400.
- [14] G. Beuf, *Phys.Rev.* D89 (2014) 074039, arXiv:1401.0313.
- [15] A. Sabio Vera, *Nucl.Phys.* B722 (2005) 65, arXiv:hep-ph/0505128.
- [16] H1, ZEUS Collaboration, F. Aaron et al., *JHEP* 1001 (2010) 109, arXiv:0911.0884.
- [17] J. L. Albacete, arXiv:1507.07120.