

**Statistical optimization for passive scalar transport:
maximum entropy production versus maximum
Kolmogorov–Sinai entropy**

M. Mihelich, D. Faranda, Bérengère Dubrulle, D. Paillard

► **To cite this version:**

M. Mihelich, D. Faranda, Bérengère Dubrulle, D. Paillard. Statistical optimization for passive scalar transport: maximum entropy production versus maximum Kolmogorov–Sinai entropy. *Nonlinear Processes in Geophysics*, European Geosciences Union (EGU), 2015, 22, pp.187 - 196. <10.5194/npg-22-187-2015>. <cea-01490845>

HAL Id: cea-01490845

<https://hal-cea.archives-ouvertes.fr/cea-01490845>

Submitted on 16 Mar 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Statistical optimization for passive scalar transport: maximum entropy production versus maximum Kolmogorov–Sinai entropy

M. Mihelich, D. Faranda, B. Dubrulle, and D. Paillard

Laboratoire SPHYNX, CEA/IRAMIS/SPEC, CNRS URA 2464, 91191 Gif-sur-Yvette, France

Correspondence to: B. Dubrulle (berengere.dubrulle@cea.fr)

Received: 18 September 2014 – Published in Nonlin. Processes Geophys. Discuss.: 18 November 2014

Revised: 20 February 2015 – Accepted: 28 February 2015 – Published: 25 March 2015

Abstract. We derive rigorous results on the link between the principle of maximum entropy production and the principle of maximum Kolmogorov–Sinai entropy for a Markov model of the passive scalar diffusion called the Zero Range Process. We show analytically that both the entropy production and the Kolmogorov–Sinai entropy, seen as functions of a parameter f connected to the jump probability, admit a unique maximum denoted $f_{\max_{\text{EP}}}$ and $f_{\max_{\text{KS}}}$. The behaviour of these two maxima is explored as a function of the system disequilibrium and the system resolution N . The main result of this paper is that $f_{\max_{\text{EP}}}$ and $f_{\max_{\text{KS}}}$ have the same Taylor expansion at first order in the deviation from equilibrium. We find that $f_{\max_{\text{EP}}}$ hardly depends on N whereas $f_{\max_{\text{KS}}}$ depends strongly on N . In particular, for a fixed difference of potential between the reservoirs, $f_{\max_{\text{EP}}}(N)$ tends towards a non-zero value, while $f_{\max_{\text{KS}}}(N)$ tends to 0 when N goes to infinity. For values of N typical of those adopted by Paltridge and climatologists working on maximum entropy production ($N \approx 10$ – 100), we show that $f_{\max_{\text{EP}}}$ and $f_{\max_{\text{KS}}}$ coincide even far from equilibrium. Finally, we show that one can find an optimal resolution N_* such that $f_{\max_{\text{EP}}}$ and $f_{\max_{\text{KS}}}$ coincide, at least up to a second-order parameter proportional to the non-equilibrium fluxes imposed to the boundaries. We find that the optimal resolution N^* depends on the non-equilibrium fluxes, so that deeper convection should be represented on finer grids. This result points to the inadequacy of using a single grid for representing convection in climate and weather models. Moreover, the application of this principle to passive scalar transport parametrization is therefore expected to provide both the value of the optimal flux, and of the optimal number of degrees of freedom (resolution) to describe the system.

1 Introduction

A major difficulty in the modelling of nonlinear geophysical or astrophysical processes is the taking into account of all the relevant degrees of freedom. For example, fluid motions obeying Navier–Stokes equations usually require of the order of $N = Re^{9/4}$ modes to faithfully describe all scales between the injection scale and the dissipative scale (Frisch, 1995). In atmosphere, or ocean, where the Reynolds number exceeds 10^9 , this amounts to $N = 10^{20}$, a number too large to be handled by any existing computers (Wallace and Hobbs, 2006). The problem is even more vivid in complex systems such as planetary climate, where the coupling of litho- bio- and cryosphere with ocean and atmosphere increases the number of degrees of freedom beyond any practical figure. This justifies the long historical tradition of parametrization and statistical model reduction, to map the exact equations describing the system onto a set of simpler equations involving few degrees of freedom. The price to pay is the introduction of *free parameters*, describing the action of discarded degrees of freedom, that need to be prescribed.

When the number of free parameters is small, their prescription can be successfully done empirically through calibrating experiments or by a posteriori tuning (Rotstayn, 2000). When the number of parameters is large, such as in climate models where it reaches several hundreds (Murphy et al., 2004), such empirical procedure is inapplicable, because it is impossible to explore the whole parameter space. In that respect, it is of great interest to explore an alternative road to parametrization via application of a statistical optimization principle, such as minimizing or maximizing of a suitable cost functional. As discussed by Turkington (2013) and Pascale et al. (2012), this strategy usually leads to closed reduced equations with adjustable parameters in the closure

appearing as weights in the cost functional that can be computed explicitly. A renowned example in climate is given by a principle of maximum entropy production (MEP) that allowed (Paltridge, 1975) to derive the distribution of heat and clouds at the Earth surface with reasonable accuracy, without any parameters and with a model of a dozen of degrees of freedom (boxes). Since then, refinements of the Paltridge model have been suggested to increase its generality and range of prediction (Herbert et al., 2011). MEP states that a stationary non-equilibrium system chooses its final state in order to maximize the entropy production as is explained in Martyushev and Seleznev (2006). Rigorous justifications of its application have been searched using e.g. information theory (Dewar and Maritan, 2014) without convincing success. More recently, we have used the analogy of the climate box model of Paltridge with the asymmetric exclusion Markov process (ASEP) to establish numerically a link between the MEP and the principle of maximum Kolmogorov–Sinai entropy (MKS) (Mihelich et al., 2014). The MKS principle is a relatively new concept which extends the classical results of equilibrium physics (Monthus, 2011). This principle applied to Markov chains provides an approximation of the optimal diffusion coefficient in transport phenomena (Gómez-Gardeñes and Latora, 2008) or simulates random walk on irregular lattices (Burda et al., 2009). It is therefore a good candidate for a physically relevant cost functional in passive scalar modelling.

The goal of the present paper is to derive rigorous results on the link between MEP and MKS using a Markov model of the passive scalar diffusion called the Zero Range Process (Andjel, 1982). We find that there exists an optimal resolution N_* such that both maxima coincide to second order in the distance from equilibrium. The application of this principle to passive scalar transport parametrization is therefore expected to provide both the value of the optimal flux, and of the optimal number of degrees of freedom (resolution) to describe the system. This suggests that the MEP and MKS principle may be unified when the Kolmogorov–Sinai entropy is defined on opportunely coarse-grained partitions.

2 From passive scalar equation to ZRP model

The equation describing the transport of a passive scalar like temperature in a given velocity field $u(x, t)$ reads as

$$\partial_t T + u \partial_x T = \kappa \partial_x^2 T, \quad (1)$$

with appropriate boundary conditions, or equivalently, in non-dimensional form:

$$\partial_t T + u \partial_x T = \frac{1}{RePr} \partial_x^2 T, \quad (2)$$

where κ , Re and Pr are respectively the molecular diffusivity, the Reynolds and the Prandtl number. To solve this equation, one must know both the velocity field and the boundary

conditions, and use as many number of modes as necessary to describe the range of scales up to the scales at which molecular diffusivity takes place, i.e. roughly $(RePr)^{3/2}$ modes, where Re is the Reynolds number of the convective flow, and Pr is its Prandtl number. In geophysical flows, this number is too large to be handled even numerically (Troen and Mahrt, 1986). Moreover, in typical climate studies, the velocity flow is basically unknown as it must obey a complicated equation involving the influence of all the relevant climate components. In order to solve the equation, one must necessarily prescribe the heat flux $f = -uT + \kappa \nabla T$. The idea of Paltridge was then to discretize the passive scalar equation in boxes and prescribe the heat flux $f_{i(i+1)}$ between boxes i and $i + 1$ by maximizing the associated thermodynamic entropy production $\dot{S} = \sum_i f_{i(i+1)} (\frac{1}{T_{i+1}} - \frac{1}{T_i})$. Here, we slightly modify the Paltridge discretization approximation to make it amenable to rigorous mathematical results on Markov chains. For simplicity, we stick to a one-dimensional case (corresponding to boxes varying only in latitude) and impose the boundary conditions through two reservoirs located at each end of the chain (mimicking the solar heat flux at pole and equator). We consider a set of N boxes that can contain an arbitrary number $n \in \mathbb{N}$ of particles. We then allow transfer of particles in between two adjacent boxes via decorrelated jumps (to the right or to the left) following a 1-D Markov dynamics governed by a coupling with the two reservoirs imposing a difference of chemical potential at the ends. The resulting process is called the Zero Range Process (ZRP) (Andjel, 1982). The different jumps are described as follows. At each time step a particle can jump right with probability $p w_n$ or jump left with probability $q w_n$ where w_n is a parameter depending of the number of particles inside the box. Physically it represents the interactions between particles. At the edges of the lattice the probability rules are different: at the left edge a particle can enter with probability α and exit with probability γw_n whereas at the right edge a particle can exit with probability βw_n and enter with probability δ . Choices of different w_n give radically different behaviours. For example $w_n = 1 + b/n$ where $b \geq 0$ described condensation phenomena (Großkinsky et al., 2003) whereas $w_1 = w$ and $w_n = 1$ if $n \geq 2$ has been used to modelled road traffic. We consider in this paper the particular case where $w = 1$ by convenience of calculation. Moreover, without loss of generality we will take $p \geq q$ which corresponds to a particle flow from the left to the right and denote $f = p - q$. After a sufficiently long time the system reaches a non-equilibrium steady state. The interest of this toy model is that it is simple enough so that exact computations are analytically tractable.

Taking the continuous limit of this process, it may be checked that the fugacity z , which is a quantity related to the average particle density (see Eq. 9 below) of stationary solutions of a system consisting of boxes of size $\frac{1}{N}$, follows

the continuous equation (Levine et al., 2005) :

$$f \frac{\partial z}{\partial x} - \frac{1}{2N} \frac{\partial^2 z}{\partial x^2} = 0, \quad (3)$$

corresponding to a stationary solution of a non-dimensional passive scalar equation with non-dimensional velocity f and a non-dimensional diffusivity $\frac{1}{RePr} = \frac{1}{2N}$. Therefore, the fugacity of the Zero Range Process is a passive scalar obeying a convection–diffusion equation, with advection velocity controlled by the probability to jump to the right or to the left, and diffusivity controlled by the number of boxes: the larger the number of boxes (the finer the resolution) the smaller the diffusivity. This observation illuminates the well-known observation that the numerical diffusion of a discrete model of diffusion is inversely proportional to the resolution. The parameter f controls the regime: $f = 0$ corresponds to a purely conductive regime whereas the larger f the more convective the regime. In the sequel, we calculate the entropy production and the Kolmogorov–Sinai entropy function of f . These two quantities reach a maximum denoted respectively $f_{\max EP}$ and $f_{\max KS}$. The MEP principle (resp. the MKS principle) states that the system will choose $f = f_{\max EP}$ (resp $f = f_{\max KS}$).

We will first show that numerically $f_{\max EP} \approx f_{\max KS}$ even far from equilibrium for a number of boxes N roughly corresponding to the resolution taken by Paltridge (1975) in his climate model. This result is similar to what we found for the ASEP model (Mihelich et al., 2014) and thus gives another example of a system in which the two principles are equivalent. Moreover we will see analytically that $f_{\max EP}$ and $f_{\max KS}$ have the same behaviour in first order in the difference of the chemical potentials between the two reservoirs for N large enough. These results provide a better understanding of the relationship between the MEP and the MKS principles.

3 Notation and useful preliminary results

This Markovian process is a stochastic process with an infinite number of states in bijection with \mathbb{N}^N . In fact, each state can be written $n = (n_1, n_2, \dots, n_N)$ where n_i is the number of particle lying in site i . We call P_n the stationary probability to be in state n . In order to calculate this probability it is easier to use a quantum formalism than the Markovian formalism (Domb, 2000; Levine et al., 2005).

The probability to find m particles in the site k is equal to $p_k(n_k = m) = \frac{z_k^m}{Z_k}$ where Z_k is the analogue of the grand canonical repartition function and z_k is the fugacity between 0 and 1. Moreover $Z_k = \sum_{i=0}^{\infty} z_k^i = \frac{1}{1-z_k}$. So, finally

$$p_k(n_k = m) = (1 - z_k) z_k^m. \quad (4)$$

We can show that the probability P over the states is the tensorial product of the probability p_k over the boxes:

$$P = p_1 \otimes p_2 \otimes \dots \otimes p_N.$$

Thus events $(n_k = m)$ and $(n_{k'} = m')$ for $k \neq k'$ are independent and so

$$P(m_1, m_2, \dots, m_N) = p_1(n_1 = m_1) * \dots * p_N(n_N = m_N). \quad (5)$$

So finally

$$P(m_1, m_2, \dots, m_N) = \prod_{k=1}^N (1 - z_k) z_k^{m_k}. \quad (6)$$

Moreover, with the Hamiltonian equation found from the quantum formalism we can find the exact values of z_k function of the system parameters:

$$z_k = \frac{(\frac{p}{q})^{k-1} [(\alpha + \delta)(p - q) - \alpha\beta + \gamma\delta] - \gamma\delta + \alpha\beta(\frac{p}{q})^{N-1}}{\gamma(p - q - \beta) + \beta(p - q + \gamma)(\frac{p}{q})^{N-1}}, \quad (7)$$

and the flux of particles c :

$$c = (p - q) \frac{-\gamma\delta + \alpha\beta(\frac{p}{q})^{N-1}}{\gamma(p - q - \beta) + \beta(p - q + \gamma)(\frac{p}{q})^{N-1}}. \quad (8)$$

Finally, the stationary density is related to the fugacity by the relation

$$\rho_k = z_k \frac{\partial \log Z_k}{\partial z_k} = \frac{z_k}{1 - z_k}. \quad (9)$$

3.1 Entropy production

For a system subject to internal forces X_i and associated fluxes J_i the macroscopic entropy production is well known (Onsager, 1931) and takes the form

$$\sigma = \sum_i J_i \cdot X_i.$$

The physical meaning of this quantity is a measure of irreversibility: the larger σ the more irreversible the system.

In the case of the ZRP, irreversibility is created by the fact that $p \neq q$. We will parametrize this irreversibility by the parameter $f = p - q$ and we will take $p + q = 1$. In the remainder of the paper, we take, without loss of generality, $p \leq q$ which corresponds to a flow from left to right. Moreover, the only flux to be considered here is the flux of particles c and the associated force is due to the gradient of the density of particles $\rho : X = \nabla \log \rho$ (Balian, 1992).

Thus, when the stationary state is reached, i.e. when c is constant,

$$\sigma = \sum_{i=1}^{N-1} c \cdot (\log(\rho_i) - \log(\rho_{i+1})) = c \cdot (\log(\rho_1) - \log(\rho_N)). \quad (10)$$

Thus, according to Eqs. (7), (8), (9) and (10) when N tends to $+\infty$ we obtain:

$$\sigma(f) = \frac{\alpha f}{f + \gamma} \left(\log \left(\frac{\alpha}{f + \gamma - \alpha} \right) - \log \left(\frac{(\alpha + \delta)f + \gamma\delta}{f(\beta - \alpha - \delta) + \beta\gamma - \gamma\delta} \right) \right). \quad (11)$$

Because $f \geq 0$ the entropy production is positive if and only if $\rho_1 \geq \rho_N$ and iff $z_1 \geq z_N$. This is physically coherent be-

cause fluxes are in the opposite direction of the gradient. We remark that if $f = 0$ then $\sigma(f) = 0$. Moreover, when f increases $\rho_1(f)$ decreases and $\rho_2(f)$ increases until they take the same value. Thus there exists f , large enough, for which $\sigma(f) = 0$. Between these two values of f the entropy production has at least one maximum.

3.2 Kolmogorov–Sinai entropy

There are several ways to introduce the Kolmogorov–Sinai entropy which is a mathematical quantity introduced by Kolmogorov and developed by the renowned mathematicians Sinai and Billingsley (Billingsley, 1965). Nevertheless, for a Markov process we can give it a simple physical interpretation: the Kolmogorov–Sinai entropy is the time derivative of the Jaynes entropy (entropy over the path):

$$S_{\text{Jaynes}}(t) = - \sum_{\Gamma_{[0,t]}} p_{\Gamma_{[0,t]}} \log(p_{\Gamma_{[0,t]}}), \quad (12)$$

For a Markov Chain we thus have

$$S_{\text{Jaynes}}(t) - S_{\text{Jaynes}}(t-1) = - \sum_{(i,j)} \mu_{i,\text{stat}} p_{ij} \log(p_{ij}), \quad (13)$$

where $\mu_{\text{stat}} = \mu_{i,\text{stat}}$ $i = 1, \dots, N$ is the stationary measure and where the p_{ij} are the transition probabilities.

Thus the Kolmogorov–Sinai entropy takes the following form:

$$h_{\text{KS}} = - \sum_{(i,j)} \mu_{i,\text{stat}} p_{ij} \log(p_{ij}). \quad (14)$$

For the ZRP, we show in the Appendix that it can be written as

$$\begin{aligned} h_{\text{KS}} = & -(\alpha \log \alpha + \delta \log \delta + \gamma \log \gamma + \beta \log \beta \\ & + (N-1)(p \log(p) + q \log(q)) + (p \log(p) + q \log(q)) \\ & \sum_{i=1}^N (1 - z_i) + (\gamma \log(\gamma) + p \log(p))(1 - z_1) \\ & + (\beta \log(\beta) + q \log(q))(1 - z_N). \end{aligned} \quad (15)$$

4 Results

We start by pointing to some interesting properties of f_{maxEP} and f_{maxKS} , then present numerical experiments on the ZRP model and conclude with some analytical computations.

Note that for N, α, β, γ and δ fixed the entropy production as well as the Kolmogorov–Sinai entropy seen as functions of f both admit a unique maximum. When N tends to infinity and $f = 0$, using Eq. (7) (i.e. the symmetric case), we find that $z_1 = \frac{\alpha}{\gamma}$ and $z_N = \frac{\delta}{\beta}$. Thus, the system is coupled with two reservoirs with respective chemical potential $\frac{\alpha}{\gamma}$ (left) and $\frac{\delta}{\beta}$ (right). For $\frac{\alpha}{\gamma} \neq \frac{\delta}{\beta}$ the system is out of equilibrium. We assume, without loss of generality, $z_1 \geq z_N$, which corresponds

to a flow from left to right. As a measure of deviation from equilibrium we take $s = z_1 - z_N$: the larger s , the more density fluxes we expect into the system.

First we remark that f_{maxEP} hardly depends on N whereas f_{maxKS} depends strongly on N . This is easily understood because σ depends only on z_1 and z_N whereas h_{KS} depends on all the z_i . Moreover, the profile of the z_i depends strongly on N . In particular, for a fixed difference of potential between the reservoirs, $f_{\text{maxEP}}(N)$ tends towards a non-zero value, while $f_{\text{maxKS}}(N)$ tends to 0 when N goes to infinity.

Moreover, f_{maxEP} and f_{maxKS} coincide even far from equilibrium for N corresponding to the choice of Paltridge (1975) $N \approx 10 \sim 100$. For N fixed, as large as one wants, and for all ϵ , as small as one wants, there exists ν such that for all $s \in [0; \nu]$ $|f_{\text{maxEP}} - f_{\text{maxKS}}| \leq \epsilon$.

These observations are confirmed by the results presented in Figs. 1 and 3 where EP and KS are calculated using Eqs. (7) and (15) for $s = 0.13$ and three different partitions: $N = 20$, $N = 100$ and $N = 1000$. The figure shows that f_{maxEP} and f_{maxKS} coincide with good approximation for $N = 20$ and $N = 100$. But then when N increases $f_{\text{maxKS}}(N)$ tends to 0 whereas $f_{\text{maxEP}}(N)$ tends to a non-zero value.

In Fig. 2 we represent the entropy production (top) and KS entropy (bottom) as functions of f for $N = 1000$ and for three value of s : $s = 0.13$; $s = 0.2$; $s = 0.04$. This supports the claim that for N fixed, we can try different values of s such that $s \in [0; \nu]$ $|f_{\text{maxEP}} - f_{\text{maxKS}}| \leq \epsilon$. Figure 3 shows that Δf_{max} is minimum when the system is close to equilibrium whereas the further the system is from equilibrium (when s increases) the more Δf_{max} increases. Moreover, the optimal resolution where f_{maxEP} and f_{maxKS} coincide is approximately 10–100. Then Δf_{max} is maximum at $N = 500$ and $s = 0.05$. Δf_{max} is obviously linear in s , for small values of s , but the behaviour with N is more complicated.

Such numerical investigations help to understand why $f_{\text{maxKS}}(N)$ and $f_{\text{maxEP}}(N)$ have different behaviour functions of N , and why for N large enough f_{maxKS} and f_{maxEP} have the same behaviour of first order in the deviation from equilibrium measured by the parameter s . We will see that we can get a precise answer to such questions by doing calculations and introducing a sort of hydrodynamics approximation.

4.1 Taylor expansion

From Eq. (15) it is apparent that f_{maxKS} depends on N whereas from Eq. (10) we get that f_{maxEP} hardly depends on N . Indeed there is a difference between f_{maxEP} and f_{maxKS} , i.e. a difference between the two principles for the ZRP. Nevertheless, we have seen numerically that there is a range of N , namely $N \approx 10\text{--}100$ for which the maxima fairly coincide.

Using Eqs. (15), (7), and (11) we compute analytically the Taylor expansion of f_{maxEP} and f_{maxKS} in s . We will show the main result: f_{maxEP} and f_{maxKS} have the same Taylor expansion in first order in s for N large enough. Their Taylor




Figure 1. Entropy production calculated using Eq. (11) (left) and KS entropy calculated using Eqs. (7) and (15) (right) as functions of f for $s = 0.13$ and respectively $N = 20$, $N = 100$ and $N = 1000$.

expansions are different up to the second order in s but there exists an N , i.e. a resolution, such that $f_{\max_{\text{EP}}}$ and $f_{\max_{\text{KS}}}$ coincident up to the second order.

Let us start by computing $f_{\max_{\text{KS}}}$. It does not depend of the constant terms of h_{KS} in Eq. (15) and therefore we need only concern ourselves with

$$\begin{aligned}
 & -(p \log(p) + q \log(q)) \left(\sum_{i=1}^N (z_i) - 1 \right) + (\gamma \log(\gamma)) \\
 & + p \log(p)(1 - z_1) + (\beta \log(\beta) + q \log(q))(1 - z_N) \\
 & = N \cdot H(f, N, \alpha, \gamma, \beta, \delta).
 \end{aligned} \tag{16}$$

Using Eq. (7), the expression of $H(f, N, \alpha, \gamma, \beta, \delta)$ takes an easy form. To simplify the calculations, we restrict the space of parameter by assuming $\alpha + \gamma = 1$ and $\beta + \delta = 1$ and we parametrize the deviation from equilibrium by the parameter $\bar{s} = \alpha - \delta$. Moreover let us denote $a = \frac{1}{N}$. Thus, we have $H(f, N, \alpha, \gamma, \beta, \delta) = H(f, a, \alpha, \bar{s})$. In order to know the Taylor expansion to the first order in \bar{s} of $f_{\max_{\text{KS}}}$ we develop $H(f, a, \alpha, \bar{s})$ up to the second order in f ; i.e. we have $H(f, a, \alpha, \bar{s}) = C + Bf + Af^2 + o(f^2)$. Then we find

$f_{\max_{\text{KS}}} = -B/2A$ which we will develop in the power of \bar{s} . This is consistent if we assume $f \ll a$.

After some tedious but straightforward calculations, we get at the first order in \bar{s} :

$$f_{\max_{\text{KS}}}(\bar{s}) = \frac{1}{4} \frac{(1 - \alpha) - a(\alpha + 2)}{\alpha(1 - \alpha) + 2a\alpha(\alpha - 1)} \bar{s} + o(\bar{s}) \tag{17}$$

and so

$$f_{\max_{\text{KS}}}(\bar{s}) = \frac{1}{4\alpha} \bar{s} + \frac{3a}{4(\alpha - 1)} \bar{s} + o(\bar{s}) + o(a\bar{s}). \tag{18}$$

We repeat the same procedure starting from Eq. (11) and we obtain

$$f_{\max_{\text{EP}}}(\bar{s}) = \frac{\bar{s}}{4\alpha} + o(\bar{s}) + o(a). \tag{19}$$

Thus, since $a = \frac{1}{N} \ll 1$ the behaviour of $f_{\max_{\text{KS}}}(\bar{s})$ and $f_{\max_{\text{EP}}}(\bar{s})$ is the same for \bar{s} small enough.

We remark that we can strictly find the same result by solving the hydrodynamics continuous approximation given by Eq. (3). This equation is a classical convection–diffusion




Figure 2. Entropy production (left) and KS entropy (right) as functions of f for $N = 1000$ and respectively $s = 0.13$, $s = 0.2$ and $s = 0.04$.




Figure 3. 2-D plot representing $\Delta f_{\max} = f_{\max_{EP}} - f_{\max_{KS}}$ in the (N, s) space.

equation. We remark that, by varying f , we change the convective behaviour: $f = 0$ corresponds to a purely diffusive regime whereas by increasing f we enhance the role of convection. If the system is near equilibrium then $f_{\max_{EP}} \approx f_{\max_{KS}} \approx 0$ and the system is purely diffusive. When the system is out of equilibrium $f_{\max_{EP}}$ and $f_{\max_{KS}}$ are different from 0 and the system corresponds to an (optimal) trade-off between purely diffusive and convective behaviour.

One can verify this numerically: we first calculate the exact values of the entropy production function of f using Eq. (7) and the Kolmogorov–Sinai entropy function of f us-

ing Eqs. (7) and (15). Then we approximate these two curves with a cubic spline approximation in order to find $f_{\max_{EP}}$ and $f_{\max_{KS}}$.

In order to find the optimal resolution N_* we can go one step further by expanding $f_{\max_{EP}}$ and $f_{\max_{KS}}$ up to the second order in \bar{s} :

$$f_{\max_{EP}}(\bar{s}) = \frac{\bar{s}}{4\alpha} + \frac{\bar{s}^2(\alpha + 1)}{8\alpha^2(\alpha - 1)} + o(\bar{s}^2) + o(a), \quad (20)$$

$$f_{\max_{KS}}(\bar{s}) = \frac{1}{4} \frac{(1 - \alpha) - a(\alpha + 2)}{\alpha(1 - \alpha) + 2a\alpha(\alpha - 1)} \bar{s} + \frac{(1 - \alpha)^2 + a(\alpha^2 - 2\alpha + 1)}{8\alpha^2(\alpha - 1)^2(1 - 2a)} \bar{s}^2 + o(\bar{s}^2). \quad (21)$$

Thus, $f_{\max_{EP}}$ and $f_{\max_{KS}}$ coincide in second order in \bar{s} iff a satisfies the quadratic equation

$$(4\alpha - 6\alpha^2 + 6\alpha^3 - 4\bar{s} + 3\alpha^2\bar{s})a^2 - \frac{1}{2}(8\alpha - 8\bar{s} + 3\alpha^2\bar{s} - 6\alpha^2 + 6\alpha^3)a - (1 - \alpha) = 0. \quad (22)$$




Figure 4. $f_{\max_{EP}}$ (left) and $f_{\max_{KS}}$ (right) as functions of \bar{s} for $\alpha = 0.5$ and $N = 100$. Both $f_{\max_{KS}}$ and $f_{\max_{EP}}$ have a linear behaviour with slope respectively 0.48 and 0.49, close to $\frac{1}{4\alpha} = 0.5$.




Figure 5. $f_{\max_{KS}(\bar{s})}$ (left) and $f_{\max_{EP}(\bar{s})}$ (right) as functions of α and in black the curve $f(\bar{s}) = \frac{1}{4\alpha}\bar{s}$. Note that the approximation $f_{\max_{KS}(\bar{s})} \approx f_{\max_{EP}(\bar{s})} \approx \frac{1}{4\alpha}\bar{s}$ is good.

This equation has a unique positive solution because the leading coefficient is positive for s small enough ($4\alpha - 6\alpha^2 + 6\alpha^3 - 4\bar{s} + 3\alpha^2\bar{s}$) ≥ 0 and the constant term is negative $-(1-\alpha) \leq 0$. We remark that the optimal resolution $N_* = \frac{1}{a^*}$ depends on the parameters of the system namely on the degree of non-equilibrium. This fact can be the explanation for two well-known issues in climate/weather modelling. First, it explains that, when downgrading or upgrading the resolution of convection models, the relevant parameters must be changed as they depend on the grid size. Second, it suggests that if the resolution is well tuned to represent a particular range of convective phenomena, it might fail in capturing the dynamics out of this range: since finer grids are needed to better represent deep convection phenomena, the deviations between model and observations observed in the distribution of extreme convective precipitation may be due to an inadequacy of the grid used.

5 Conclusions

We have shown how a simple 1-D Markov process, the Zero Range Process, can be used to obtain rigorous results on the problem of parametrization of the passive scalar transport problem, relevant to many geophysical applications including temperature distribution in climate modelling. Using this model, we have derived rigorous results on the link between a principle of maximum entropy production and the principle of maximum Kolmogorov–Sinai entropy using a Markov model of the passive scalar diffusion called the Zero Range Process. The Kolmogorov–Sinai entropy seen as a function of the convective velocity admits a unique maximum. We

show analytically that both have the same Taylor expansion at the first order in the deviation from equilibrium. The behaviour of these two maxima is explored as a function of the resolution N (equivalent to the number of boxes, in the box approximation). We found that, for a fixed difference of potential between the reservoirs, the maximal convective velocity predicted by the maximum entropy production principle tends towards a non-zero value, while the maximum predicted using Kolmogorov–Sinai entropy tends to 0 when N goes to infinity. For values of N typical of those adopted by climatologists ($N \approx 10-100$), we show that the two maxima nevertheless coincide even far from equilibrium. Finally, we show that there is an optimal resolution N_* such that the two maxima coincide to second order in \bar{s} , a parameter proportional to the non-equilibrium fluxes imposed to the boundaries. The fact that the optimal resolution depends on the intensity of the convective phenomena to be represented, points to new interesting research avenues, e.g. the introduction of convective models with adaptive grids optimized with maximum entropy principles on the basis of the convective phenomena to be represented.

The application of this principle to passive scalar transport parametrization is therefore expected to provide both the value of the optimal flux, and of the optimal number of degrees of freedom (resolution) to describe the system. It would be interesting to apply it to a more realistic passive scalar transport problem, to see if it would yield a model that could be numerically handled (i.e. corresponding to a number of boxes that is small enough to be handled by present computers). In view of applications to atmospheric convection, it

would be interesting to apply this procedure to the case of an active scalar, coupled with a Navier–Stokes equation for the velocity. In such a case, the role of f will be played by the turbulent subgrid Reynolds stresses. The heat fluxes and N^* will be fixed by the coarse-graining length, and the optimization procedure will in principle provide the optimum subgrid Reynolds stresses at a given resolution N . Moreover, by imposing coincidence of MKS and MEP, one could get both the Reynolds stresses, heat fluxes and the optimum resolution. Further, on the theoretical side, it will be interesting to study whether for general dynamical systems, there exists a smart way to coarse grain the Kolmogorov–Sinai entropy such that its properties coincide with the thermodynamic entropy production. This would eventually justify the use of the MEP principle and explain the deviations as well as the different representations of it due to the dependence of the dynamic (Kolmogorov–Smirnov–Tsallis–Jaynes) entropies on the kind of partition adopted.

Appendix A: Computation of the K-S entropy

In this appendix, we compute the Kolmogorov–Sinai entropy for the Zero Range Process, starting from its definition in Eq. (14). In the frame of our Zero Range Process, we use Eqs. (14) and (6) to write it as

$$\begin{aligned}
h_{\text{KS}} &= -\sum_i \mu_{i\text{stat}} \sum_j p_{ij} \log(p_{ij}) = -\sum_{m_1=0}^{+\infty} \dots \sum_{m_N=0}^{+\infty} \\
&P(m_1, m_2, \dots, m_N) \sum_j p_{(m_1, \dots, m_N) \rightarrow j} \log(p_{(m_1, \dots, m_N) \rightarrow j}) \\
&= -\sum_{m_1=0}^{+\infty} P(m_1) \dots \sum_{m_N=0}^{+\infty} \\
&P(m_N) \sum_j p_{(m_1, \dots, m_N) \rightarrow j} \log(p_{(m_1, \dots, m_N) \rightarrow j}). \quad (\text{A1})
\end{aligned}$$

We thus have to calculate $\sum_j p_{(m_1, \dots, m_N) \rightarrow j} \log(p_{(m_1, \dots, m_N) \rightarrow j})$ that we will refer to as (*). We will take $p + q = \alpha + \delta = \beta + \gamma = 1$ and $dt = \frac{1}{N}$ in order to neglect the probabilities to stay in the same state compare to the probabilities of changing state. There are five different cases to consider:

1. if $\forall i m_i \geq 1$ so the possible transitions are:
 $(m_1, m_2, \dots, m_N) \rightarrow (m_1 \pm 1, m_2, \dots, m_N)$ with respective probabilities α and δ ;
 $(m_1, m_2, \dots, m_N) \rightarrow (m_1, m_2, \dots, m_N \pm 1)$ with respective probabilities γ and β ;
and $(m_1, \dots, m_k, \dots, m_N) \rightarrow (m_1, \dots, m_k \pm 1, \dots, m_N)$ with respective probabilities p and q ;
thus,

$$\begin{aligned}
(*) &= \alpha \log \alpha + \delta \log \delta + \gamma \log \gamma + \beta \log \beta \\
&+ (N-1)(p \log(p) + q \log(q)); \quad (\text{A2})
\end{aligned}$$

2. if $m_1 \geq 1$ and $m_N \geq 1$ and let i be the number of m_i between 2 and $N-1$ equal to 0. With the same argument as previously we have

$$\begin{aligned}
(*) &= \alpha \log \alpha + \delta \log \delta + \gamma \log \gamma + \beta \log \beta \\
&+ (N-1-i)(p \log(p) + q \log(q)); \quad (\text{A3})
\end{aligned}$$

3. if $m_1 = 0$ and $m_N \geq 1$ and let i the number of m_i between 2 and $N-1$ equal to 0 we have

$$\begin{aligned}
(*) &= \alpha \log \alpha + \delta \log \delta + \beta \log \beta \\
&+ (N-2-i)p \log(p) + (N-1-i)q \log(q); \quad (\text{A4})
\end{aligned}$$

4. the same applies if $m_1 \geq 1$ and $m_N = 0$ and let i the number of m_i between 2 and $N-1$ equal to 0 we have

$$\begin{aligned}
(*) &= \alpha \log \alpha + \delta \log \delta + \gamma \log \gamma \\
&+ (N-1-i)p \log(p) + (N-2-i)q \log(q); \quad (\text{A5})
\end{aligned}$$

5. finally, if $m_1 = 0$ and $m_N = 0$ and let i the number of m_i between 2 and $N-1$ equal to 0 we have

$$\begin{aligned}
(*) &= \alpha \log \alpha + \delta \log \delta \\
&+ (N-2-i)(p \log(p) + q \log(q)). \quad (\text{A6})
\end{aligned}$$

Using Eq. (4) we find that $P(m_k = 0) = 1 - z_k$ and $\sum_{i=1}^{+\infty} P(m_k = i) = z_k$.
We thus obtain that

$$\begin{aligned}
h_{\text{KS}} &= -(\alpha \log \alpha + \delta \log \delta + \gamma \log \gamma + \beta \log \beta) \\
&+ (N-1)(p \log(p) + q \log(q)) \\
&+ (p \log(p) + q \log(q)) \\
&\left(\sum_{r=0}^N r \sum_{i_1 \dots i_N i=i_1, \dots, i_r} \prod_{i \neq i_1 \dots i_r} (1 - z_i) \prod_{i \neq i_1 \dots i_r} z_i \right) \\
&+ (\gamma \log(\gamma) + p \log(p)) z_N (1 - z_1) \\
&\left(\sum_{i_2 \dots i_{N-1} i=i_2, \dots, i_r} \prod_{i \neq i_2 \dots i_r} (1 - z_i) \prod_{i \neq i_2 \dots i_r} z_i \right) \\
&+ (\beta \log(\beta) + q \log(q)) z_1 (1 - z_N) \\
&\left(\sum_{i_2 \dots i_{N-1} i=i_2, \dots, i_r} \prod_{i \neq i_2 \dots i_r} (1 - z_i) \prod_{i \neq i_2 \dots i_r} z_i \right) \\
&+ (\beta \log(\beta) + \gamma \log \gamma + p \log p + q \log q) \\
&\left(\sum_{i_2 \dots i_{N-1} i=i_2, \dots, i_r} \prod_{i \neq i_2 \dots i_r} (1 - z_i) \prod_{i \neq i_2 \dots i_r} z_i \right). \quad (\text{A7})
\end{aligned}$$

This expression, though complicated at first sight, can be simplified. Indeed, interested in the function $F(a) = \prod_{k=1}^N (z_k + a(1 - z_k))$ and by deriving subject to a we show that

$$\sum_{r=0}^N r \sum_{i_1 \dots i_N i=i_1, \dots, i_r} \prod_{i \neq i_1 \dots i_r} (1 - z_i) \prod_{i \neq i_1 \dots i_r} z_i = \sum_{i=1}^N (1 - z_i). \quad (\text{A8})$$

Thus we can simplify the earlier equation to obtain

$$\begin{aligned}
h_{\text{KS}} &= -(\alpha \log \alpha + \delta \log \delta + \gamma \log \gamma + \beta \log \beta) \\
&+ (N-1)(p \log(p) + q \log(q)) \\
&+ (p \log(p) + q \log(q)) \sum_{i=1}^N (1 - z_i) \\
&+ (\gamma \log(\gamma) + p \log(p))(1 - z_1) \\
&+ (\beta \log(\beta) + q \log(q))(1 - z_N). \quad (\text{A9})
\end{aligned}$$

Acknowledgements. M. Mihelich thanks IDEEX Paris Saclay for financial support.

Edited by: J. M. Redondo

Reviewed by: two anonymous referees

References

- Andjel, E. D.: Invariant measures for the zero range process, *Ann. Probabil.*, 525–547, 1982.
- Balian, R.: *Physique statistique et thermodynamique hors équilibre*, Ecole Polytechnique, 1992.
- Billingsley, P.: *Ergodic theory and information*, Wiley, 1965.
- Burda, Z., Duda, J., Luck, J. M., and Waclaw, B.: Localization of the maximal entropy random walk, *Phys. Rev. Lett.*, 102, 160602, 2009.
- Dewar, R. C. and Maritan, A.: A theoretical basis for maximum entropy production, in: *Beyond the Second Law*, 49–71, Springer, 2014.
- Domb, C.: *Phase transitions and critical phenomena*, Vol. 19, Academic Press, 2000.
- Frisch, U.: *Turbulence: the legacy of AN Kolmogorov*, Cambridge University Press, 1995.
- Gómez-Gardeñes, J. and Latora, V.: Entropy rate of diffusion processes on complex networks, *Phys. Rev. E*, 78, 065102, 2008.
- Großkinsky, S., Schütz, G. M., and Spohn, H.: Condensation in the zero range process: stationary and dynamical properties, *J. Stat. Phys.*, 113, 389–410, 2003.
- Herbert, C., Paillard, D., Kageyama, M., and Dubrulle, B.: Present and Last Glacial Maximum climates as states of maximum entropy production, *Q. J. Roy. Meteorol. Soc.*, 137, 1059–1069, 2011.
- Levine, E., Mukamel, D., and Schütz, G.: Zero-range process with open boundaries, *J. Stat. Phys.*, 120, 759–778, 2005.
- Martyushev, L. M. and Seleznev, V. D.: Maximum entropy production principle in physics, chemistry and biology, *Phys. Rep.*, 426, 1–45, 2006.
- Mihelich, M., Dubrulle, B., Paillard, D., and Herbert, C.: Maximum Entropy Production vs. Kolmogorov-Sinai Entropy in a Constrained ASEP Model, *Entropy*, 16, 1037–1046, 2014.
- Monthus, C.: Non-equilibrium steady states: maximization of the Shannon entropy associated with the distribution of dynamical trajectories in the presence of constraints, *J. Stat. Mech.*, p. P03008, 2011.
- Murphy, J. M., Sexton, D. M., Barnett, D. N., Jones, G. S., Webb, M. J., Collins, M., and Stainforth, D. A.: Quantification of modelling uncertainties in a large ensemble of climate change simulations, *Nature*, 430, 768–772, 2004.
- Onsager, L.: Reciprocal relations in irreversible processes. I., *Phys. Rev.*, 37, 2265–2279, 1931.
- Paltridge, G. W.: Global dynamics and climate—a system of minimum entropy exchange, *Q. J. Roy. Meteorol. Soc.*, 101, 475–484, 1975.
- Pascale, S., Gregory, J. M., Ambaum, M. H., and Tailleux, R.: A parametric sensitivity study of entropy production and kinetic energy dissipation using the FAMOUS AOGCM, *Clim. Dynam.*, 38, 1211–1227, 2012.
- Rotstayn, L. D.: On the “tuning” of autoconversion parameterizations in climate models, *J. Geophys. Res.-Atmos.* (1984–2012), 105, 15495–15507, 2000.
- Troen, I. and Mahrt, L.: A simple model of the atmospheric boundary layer; sensitivity to surface evaporation, *Boundary-Lay. Meteorol.*, 37, 129–148, 1986.
- Turkington, B.: An optimization principle for deriving nonequilibrium statistical models of hamiltonian dynamics, *J. Stat. Phys.*, 152, 569–597, 2013.
- Wallace, J. M. and Hobbs, P. V.: *Atmospheric science: an introductory survey*, Vol. 92, Academic press, 2006.