

## Analysis of intrachromosomal homologous recombination in mammalian cell, using tandem repeat sequences

S. Lambert, Y. Saintigny, F Delacote, F Amiot, B. Chaput, M. Lecomte, S.  
Huck, P. Bertrand, B. Lopez

► **To cite this version:**

S. Lambert, Y. Saintigny, F Delacote, F Amiot, B. Chaput, et al.. Analysis of intrachromosomal homologous recombination in mammalian cell, using tandem repeat sequences. Mutation Research/Mutation Research Genomics, Elsevier, 1999, 433, pp.159 - 168. cea-01938132

**HAL Id: cea-01938132**

**<https://hal-cea.archives-ouvertes.fr/cea-01938132>**

Submitted on 28 Nov 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Review

# Analysis of intrachromosomal homologous recombination in mammalian cell, using tandem repeat sequences

S. Lambert<sup>1</sup>, Y. Saintigny<sup>1</sup>, F. Delacote, F. Amiot, B. Chaput, M. Lecomte, S. Huck, P. Bertrand, B.S. Lopez<sup>\*</sup>

*UMR 217 CNRS, CEA, DSV, DRR, 60-68 Av. du Général Leclerc BP6 92 265 Fontenay aux Roses, Cedex, France*

Received 6 January 1998; received in revised form 25 January 1999; accepted 10 February 1999

## Abstract

In all the organisms, homologous recombination (HR) is involved in fundamental processes such as genome diversification and DNA repair. Several strategies can be devised to measure homologous recombination in mammalian cells. We present here the interest of using intrachromosomal tandem repeat sequences to measure HR in mammalian cells and we discuss the differences with the ectopic plasmids recombination. The present review focuses on the molecular mechanisms of HR between tandem repeats in mammalian cells. The possibility to use two different orientations of tandem repeats (direct or inverted repeats) in parallel constitutes also an advantage. While inverted repeats measure only events arising by strand exchange (gene conversion and crossing over), direct repeats monitor strand exchange events and also non-conservative processes such as single strand annealing or replication slippage. In yeast, these processes depend on different pathways, most of them also existing in mammalian cells. These data permit to devise substrates adapted to specific questions about HR in mammalian cells. The effect of substrate structures (heterologies, insertions/deletions, GT repeats, transcription) and consequences of DNA double strand breaks induced by ionizing radiation or endonuclease (especially the rare-cutting endonuclease ISce-I) on HR are discussed. Finally, transgenic mouse models using tandem repeats are briefly presented. © 1999 Elsevier Science B.V. All rights reserved.

*Keywords:* Homologous recombination; Mammalian cells; Tandem repeat sequences

## 1. Introduction

Homologous recombination (HR) plays a central role in various fundamental processes determining genome organization and rearrangement such as molecular evolution [1,2] mating type switching in

yeast (for review, see Ref. [3]), antigenic variation of the trypanosome (for review, see Ref. [4]), or for diversification of immunoglobulin genes in chicken [5] or in rabbit [6], faithful chromosomal segregation during meiosis in yeast (for review, Refs. [7,8]), and DNA repair (for review, see Ref. [9]).

On one hand, HR is involved in the maintenance of genome integrity by repairing damaged DNA; on the other hand, it may also contribute to genome instability since recombination between homologous

<sup>\*</sup> Corresponding author. Tel.: +33-1-46-54-88-35; Fax: +33-1-46-54-91-80; E-mail: lopez@dsvidf.cea.fr

<sup>1</sup> Equal contribution.

repeated sequences dispersed through the genome may lead to duplications, inversions, deletions, translocations [10]. Moreover, gene conversion with pseudogenes can inactivate a functional allele [11]. Finally, several lines of evidences connect HR with cancer predisposition or prevention. First, carcinogens stimulate intrachromosomal HR [12–15]. Second, intrachromosomal HR is elevated in DNA repair deficient cell lines, in Ataxia telangiectasia cell lines or p53 defective cells lines, all of these phenotypes being associated with cancer predisposition [16–20]. Third, Rad51, a putative recombination protein in mammalian cells, has been shown to interact with the products of the tumor suppressor genes *p53*, *BRCA1*, *BRCA2* [21–25].

The use of intrachromosomal tandem repeat represents a good strategy to measure HR in mammalian cells and has been often used because (i) it is the easiest way to introduce the two partners of recombination in a broad variety of cells and in one round of cell transfection, (ii) these systems permit efficient measure of HR and (iii) it allows to select for gene conversion or crossover and deletion events; (iv) depending on the orientation of the two markers, it can measure recombination arising by different mechanisms such as strand exchange (SE), single strand annealing (SSA) or replication slippage [26]. These mechanisms involving different pathways in yeast, it is thus essential to be able to distinguish between them in mammalian cells also. Here we present mechanisms of HR deduced from the knowledge in yeast, and also some molecular characteristics of HR between tandem repeat in mammalian cells. These data aid in the design of substrates adapted to more specific questions about the regulation of HR in mammalian cells.

### 1.1. Mechanisms and pathways of homologous recombination between tandem repeat: the knowledge from *Saccharomyces cerevisiae*

Two different orientations of the tandem repeat recombination substrate can be used: direct repeat (DR) or inverted repeat (IR). In bacteria and yeast, RecA/Rad51 protein promotes strand exchange (SE). SE recombination can act on DR as well as on IR

and lead either to gene conversion or to crossing over (Fig. 1). Gene conversion between both DR and IR leaves intact the general structure of the locus. Depending on the orientation of the substrates, crossing over results in different structures: crossing over between IR leads to the inversion of the intervening sequence (Fig. 1B, Fig. 2A); crossing over between DR leads to the deletion of the intervening sequence either by an unequal sister chromatid exchange or by an intrachromatid exchange (Fig. 2B,C). In addition to SE recombination, direct repeats (DR) permit also to monitor deletion events arising by *RAD51* independent mechanisms such as single strand annealing (SSA) (Fig. 3), replication slippage and other mechanisms [26]. Because of their orientation, inverted repeats measure neither SSA nor replication slippage.

Extensive studies in yeast have determined genes involved in the control of HR. SE and SSA represent the main pathways using homologous sequences to achieve double strand break (DSB) repair. Besides DSB repair, replication slippage involved pathways different from those required for SE and SSA [26]. In *Saccharomyces cerevisiae*, the SE mechanism is dependent on the *RAD52* epistasis group including *RAD51*, *RAD52*, *RAD54*, *RAD55*, *RAD57* genes, whereas SSA does not involve these genes. SSA is dependent upon *SRS2*, on the nucleotide excision repair proteins Rad1 and Rad10 and on the mismatch repair proteins Msh2 and Msh3 [27,28]. Homologues to most of these genes have been described in mammalian cells (Table 1). Clearly, the study of HR in mammalian cells would benefit from the complementary use of the two kinds of substrates (DR and IR).

### 1.2. Molecular events of intrachromosomal recombination between duplicated sequences in mammalian cells

Two kinds of substrates have been used. One used two *LAC Z* sequences, recombinant cells being detected by the blue coloration. The second type of substrates used selectable genes (herpes simplex virus TK gene in *tk*<sup>-</sup> cells, neomycin resistance gene, hygromycin resistance gene...), recombinant cells




Fig. 1. Principle of the strategy to measure intrachromosomal homologous recombination between tandem repeat sequences. (A) Direct repeats, (B) inverted repeats (the letters a and z show the orientation of the intergenic sequence), (C) truncated repeats. The parental construction contains two inactive copies of a reporter gene (hatched boxes). This reporter gene can confer resistance to selective media. An alternative strategy uses the  $\beta$ -galactosidase gene whose expression can be monitored by histochemistry. In the parental form the two copies are inactivated ( $R^-$ ) by a mutation, a deletion or an insertion (black boxes) located at different positions on each copy. An alternative strategy uses two truncated but overlapping regions (C) black boxes. In most of the systems, another selection gene ( $S^+$  or  $S^-$ ) is used (i) to select the transfected cells with the recombination substrate, (ii) to analyze the distribution of recombination events (deletion or gene conversion). Recombination between the duplicated sequences creates a functional gene ( $R^+$ ) either by gene conversion (a conservative event) or by deletion of the intergenic sequence (non-conservative event). Gene conversion can occur with direct as well as with inverted repeat sequences. Deletions can only occur with direct repeat sequences by crossing over after strand exchange (SE), by single strand annealing (SSA) or other events such as replication slippage. Crossovers between the inverted repeat sequences do not lead to deletion but to inversion of the intervening sequence. With two truncated copies, a deletion creates a functional gene ( $R^+$ ).

forming clones when the selection was applied. Additionally, DSB repair has been studied following *in vivo* endonucleases treatment.

### 1.2.1. Spontaneous intrachromosomal recombination is a conservative mechanism

Analysis of the structure of recombinants has revealed that spontaneous intrachromosomal recombination is mainly a conservative mechanism in

mouse L-cells with gene conversion representing approximately 80% of the recombination events [29]. This was observed using direct repeat or inverted repeat sequences [30]. The majority of the events giving rise to crossover products involved unequally paired of sister chromatids after DNA replication (see Fig. 2) [31]. These results contrast with those of extrachromosomal recombination which follow a non-conservative mechanism in mouse L-cells [32–




Fig. 2. Models for crossing-over events after SE. (A) Crossing over between two inverted repeats leads to the inversion of the intervening sequence. The arrows indicate the orientation of the two copies, (X) symbolizes the exchange and the numbers are here to orient the intergenic sequence. (B) and (C) recombination between two direct repeats (hatched boxes). The arrows indicate the orientation of the two copies. The lines represent the duplex DNA. (B) Recombination between two mispaired chromatids. (C) Intrachromatid recombination. The two copies (hatched boxes) are paired. After the exchange (X), the resolution of the Holliday junction can lead to gene conversion (see Fig. 1) or to a crossing over event. In (B), the crossing over leads to the formation of a unique active gene and to the deletion of the intergenic sequence on one chromatid and to a triplication on the other chromatid. In (C), the crossing over leads to the maintenance of only one copy on the chromosome and one loop that can be eliminated. In some cases, the loop can be integrated elsewhere in the genome. (D) Crossover between two truncated copies. The black box represents the overlapping sequence. (A, C, D) correspond to reciprocal intrachromatid exchanges.

34]. It cannot be excluded that these ratios may differ in other cell types.

### 1.2.2. Heteroduplex formation

The different models for HR involved intermediate structures in which DNA-strand exchanges create hybrids and heteroduplex DNA between the two recombining molecules [35–37]. Biochemical studies of mitotically dividing mammalian cells indicate that

exchange between sister chromatids involved inter-strand transfer of DNA [38,39]. In addition, heteroduplex formation during HR promoted by human nuclear extracts was observed in a cell free system [40]. The analysis of the intrachromosomal recombination products between tandem repeats has shown that recombinant colonies result from unrepaired heteroduplex DNA [31,41]. Since particular structures may affect the formation, the stability or the resolu-




Fig. 3. The single strand annealing model [32]. The two complementary strands of the duplex DNA are drawn here. The heavy lines symbolized the homologous repeat sequences and the arrows the orientation of the two repeats. (1) A double strand break can occur even in the non-homologous intervening sequence. (2) After degradation by a single strand exonuclease, single stranded tails (ss-tails) are created. If the repeats are in a direct orientation, complementary ss-tails are created. (3) Annealing of the two complementary single stranded regions results in a structure leading to the deletion of the intervening sequence after the resolution of this structure. (4) SSA is exclusively a non conservative mechanism. If the repeats are in an inverted orientation, the ss-tails (step 2) are identical but not complementary and SSA cannot act.

tion of the heteroduplex recombination intermediates, the effect on recombination of the DNA struc-

ture (insertions or deletions leading to loops formation into the heteroduplex molecule the homology requirements...) have been studied.

### 1.2.3. Effect of the sequence structure: insertions, deletions, sequence homology requirements, microsatellites, transcription

**1.2.3.1. Insertions and deletions.** The molecular nature of insertion or deletion mutations (corresponding to the black boxes in Figs. 1 and 2) in the copies of the duplication can influence the efficiency of HR. DNA strand exchange is able to propagate through heterologous sequences forming heteroduplex DNA bearing loops. The resolution of such intermediates leads more frequently to the excision of the loop with an efficiency correlated to the size of the loop [42].

**1.2.3.2. Homology requirements.** Two alternative strategies can be drawn to address the question of how much homology is required for HR in mammalian cells. The first uses two truncated molecules with different lengths of overlapping of uninterrupted homology (Fig. 2D). When the two interacting molecules share length of homologies between 295 bp and 1.8 kb, the rate of gene conversion is

Table 1  
Comparison between strand exchange (SE) and single strand annealing (SSA)

Process	Products	Orientation of the substrates	Pathway in <i>S. cerevisiae</i>	Homologues in human cells	
Strand exchange	<u>Gene conversion</u>	Direct repeat	<i>RAD51</i> <sup>a</sup>	<i>HsRAD51</i> <sup>a</sup>	
		Inverted repeat	<i>RAD52</i> <sup>b</sup>	<i>HsRAD52</i>	
	<u>Crossing over</u>		<i>RAD54</i>	<i>HsRAD54</i>	
		Deletions	Direct repeat	<i>RAD55</i> <sup>a</sup>	see: <sup>a</sup>
		Inversions	Inverted repeat	<i>RAD59</i>	see: <sup>a</sup>
Single strand annealing	Deletions	Direct repeat	<i>RAD1</i>	<i>XP-F</i> <sup>c</sup>	
			<i>RAD10</i>	<i>ERCC1</i>	
			<i>MSH2</i>	<i>HsMSH2</i>	
			<i>MSH3</i>	<i>HsMSH3</i>	
			<i>SRS2</i>		

<sup>a</sup>In yeast *RAD55* and *RAD57* share homologies with *RAD51*. In mammalian cells, beside *HsRAD51*, at least 6 other *RAD51* homologues have been described: *XRCC2*, *XRCC3*, *RAD51B/HREC2*, *RAD51H3*, *RAD51C*, *RAD51D*. Some of these homologues presumably correspond to *RAD55* and to *RAD57*.

<sup>b</sup>Defines the epistasis group for homologous recombination in yeast.

<sup>c</sup>*Xeroderma pigmentosum* group F.

directly proportional to the length of uninterrupted homology. This rate is reduced 7 fold with 200 bp homology and 100 fold with 95 bp length of homology [43].

The second strategy uses two molecules of approximately the same size but containing sequence polymorphism. Waldman and Liskay used a HSV-*tk* duplication in which one *TK* copy came from HSV type 1 and the other copy from HSV type 2 (homeologous recombination). These *TK* genes share 81% of homology. The authors observed that with 19% divergence, the rate of intrachromosomal HR was reduced 1000 fold relative to the rate of HR between two identical HSV1-*tk* sequences. In contrast, the rate of intramolecular or intermolecular extrachromosomal recombination was only reduced by a factor 3 to 15 [44]. These results also argue in favor of distinct mechanisms between extrachromosomal and intrachromosomal recombination. Moreover, if efficient intrachromosomal recombination required a minimum of 134 to 232 bp of uninterrupted homology, a single-nucleotide heterology in this minimal region of homology was sufficient to inhibit efficient recombination [45]. In addition, when recombination initiates in a perfectly homologous sequence, it is able in a second step to propagate through an adjacent sequences exhibiting 19% heterologies [45]. Finally, Yang and Waldman [46] show that gene conversion involved transfer of uninterrupted blocks of information from 35 to more than 330 bp. Taken together, these results are consistent with the notion that more than 200 bp of homology are required to initiate efficient gene conversion in mammalian cells. This has also been found in *in vitro* reactions in cell free systems [47].

**1.2.3.3. Effect of GT repeats.** A variety of DNA sequences may play direct or indirect roles in recombination by their effects on the DNA structures. It was proposed that GT and GC repeats, which can form Z-DNA, may influence recombination [48]. It has been shown that GT, GC repeats and minisatellite repeats, stimulated extrachromosomal recombination of transfected DNA [49,50]. However, using an intrachromosomal assay, a (GT)<sub>29</sub> repeat has been shown to be unable to stimulate recombination and to influence the distribution of the recombination events [51].

**1.2.3.4. Transcription.** Transcription stimulates homologous recombination in *S. cerevisiae* [52,53]. Stimulation by transcription of extrachromosomal recombination [54] and intrachromosomal recombination [55] has been reported in CHO cells. Alleles transcribed at high levels recombined about 2 to 7-fold more frequently than identical alleles transcribed at low level. In line with this, preferential repair of UV damage has been shown to abolish the transcription–stimulation of HR [56]. Finally, transcription has no effect on recombination induced by a DNA double strand break [57].

### 1.3. Stimulation of homologous recombination by DNA double strand breaks

In different species, DNA-damaging agents were shown to stimulate homologous recombination. DNA double strand breaks (DSB) are the main lesion involved in the stimulation of homologous recombination in different organisms. It is used to initiate recombination during meiosis in yeast *S. cerevisiae* (for review, see Refs. [7,8]). The DSB is also one of the most genotoxic lesion induced by ionizing radiation and can be repaired by two general mechanisms: (i) non homologous end joining (NHEJ), which does not necessarily involve sequence homologies and promote the end-joining of the broken extremities; (ii) homology-directed repair which involves homologous sequences and corresponds to SE and SSA mechanisms. Ionizing radiation stimulates interallelic recombination in the endogenous *TK* locus in a human lymphoblast cell line [58]. However, the effect of ionizing radiation on intrachromosomal HR between tandem repeat sequences varies according to the cells and/or to the substrates used. Although, HR between two LacZ sequences in CV-1 cells is stimulated by ionizing radiation [59], it does not stimulate HR between HSV-TK sequences in mouse L-cells [12]. The fact that the cell lines used are different could explain this result. Another explanation could involved the differences in the substrates, LacZ vs. TK sequences. In the former case, recombinant are scored by coloration ( $\beta$ -galactosidase activity) some hours following the treatment. In contrast TK<sup>+</sup> recombination events can be scored on surviving colonies several weeks after the treatment. Thus,

in the former system, recombination events can be scored before the death of the cells generally arising several days after exposure to radiation. In contrast, the second system scores only surviving colonies.

Another way to produce DSB in target DNA uses nucleases. Restriction enzymes corresponding to single sites present in the target were electroporated into CHO cells or human cells [42,60]. In these cases, recombination was increased by more than 10-fold. Electroporation of the rare cutting yeast endonuclease *PI-SceI* also stimulated recombination provided there was a corresponding cleavage site in the duplication [60]. The yeast endonuclease *I-SceI* has provided a useful tool to study targeted DSB in mammalian cells, as already reviewed [61]. *I-SceI* recognizes a cleavage site of 18 bp long. Due to this large restriction sequence, there is probably no *I-SceI* site in the mammalian genome and expression of the *I-SceI* enzyme in mammalian cells is not toxic. However, when the *I-SceI* restriction site is present in the duplication, at least 80% of the molecules are cleaved after transfection of a plasmid expressing *I-SceI* enzyme [62]. Induction of a site-directed DSB into the recombination substrates strongly stimulates both homologous and non-homologous recombination [63,64,57]. HR is stimulated 100 times but non-homologous recombination is stimulated 1000 times [64]. However, using a physical analysis, homology-directed repair of *I-SceI*-induced DSB's is found to account for 30–50% of the observed events [65]. *I-SceI* induced recombination produced mainly deletion events (80%), that are interpreted as a result from SSA. Finally, in contrast with spontaneous HR, transcription does not stimulate DSB-induced recombination [57], although we cannot exclude that the efficiency of DSB-stimulation is so great that it would mask the effect of transcription.

#### 1.4. Transgenic mice models

The assay using duplicated repeats to measure intrachromosomal recombination can be envisioned *in vivo* in transgenic mice. In that case, recombination can only be recorded using genes giving a coloration such as the  $\beta$ -galactosidase gene, but not with a gene conferring resistance to a drug. Two models were developed to measure recombination in

specific tissues. One model has been developed to analyze the lineage of cells in the myotome; thus, the expression of the recombination substrate was driven by a promoter which confers expression specifically to cells of this compartment: the promoter of the  $\alpha$ -subunit of acetylcholine receptor. The descendants of the recombinant cells are histochemically identified, permitting the analysis of the lineage of cells in the intact embryos [66]. It appears that the frequency of recombination in this tissue is similar to the frequency measured in cultured cells: between 1 and  $2 \times 10^{-6}$  [66].

Another transgenic model also uses a duplication of  $\beta$ -galactosidase genes driven by a meiosis specific promoter. Germ line gene conversion was analyzed in transgenic male gametes. Spermatids which undergo intrachromosomal gene conversion produce functional  $\beta$ -galactosidase ( $\text{lacZ}^+$ ), visualized by histochemical staining. Approximately 2% of the spermatids, produced by a combination of meiotic and mitotic conversion events, were  $\text{LacZ}^+$  [67].

## 2. Conclusions

Numerous differences exist between intrachromosomal and extrachromosomal HR. For example, intrachromosomal recombination is conservative whilst plasmid recombination is non-conservative in mouse L-cells; in addition heterologies and GT repeats differently affected plasmid and intrachromosomal recombination. Moreover, the number of plasmid copies is not controlled and it is well established that plasmids are submitted to extensive nuclease attack after transfection. Chromatin structure and the nuclear organization may also account for the differences observed between plasmid and intrachromosomal recombination. Thus, we believe it is most suitable to measure HR between two intrachromosomal tandem repeat sequences.

The other possibilities to measure HR between two intrachromosomal sequences are interallelic and ectopic recombination, which are however technically complicate to set up. Furthermore, interallelic HR as well as ectopic HR are rather inefficient in mammalian cells [68,42,69]. Finally, the possibility of using two different orientations of tandem repeats



(direct or inverted) and the use of the rare-cutting endonuclease ISce-I provide a set of complementary approaches to study HR in mammalian cells.

Using these of strategies, connections between HR and other fundamental cellular processes have been described. Firstly, transcription stimulates HR but not DSB-induced HR [57]. Connections with cell cycle control and DNA repair have also been studied: alteration of *ATM* or p53 functions lead to an increase of HR [18–20]; carcinogens and DNA damaging agents treatments stimulate both gene conversion and deletion [12]. More generally, the efficiency of repair of DNA damages diminishes the stimulation of recombination [17,56]; finally, inhibition of Poly(ADP-ribose)polymerase (PARP) by 3-methoxybenzamide increases recombination between 3 and 4-fold in mouse L-cells [70].

Very little is known on the genetic control of homologous recombination in mammalian cells. Overexpression of human *RAD52* stimulates recombination between two *lacZ* direct repeats about 3-fold, in monkey cells and confers resistance to ionizing radiation [59]. The Rad52 stimulation acts via the single stranded DNA binding protein RPA which is also involved in replication and DNA excision repair [71]. One question would be to understand how overexpression of only one component of a multiprotein complex can stimulate the whole process, i.e., homologous recombination. One explanation could be that Rad52 is the limiting factor of the complex. However, it has recently been shown that overexpression of the Hamster Rad51 protein into CHO, also confers resistance to ionizing radiation and stimulates HR between direct repeat sequences [72]; this result is not in accordance with this hypothesis. The combined analysis of HR between direct repeats, between inverted repeats and of the DSB-induced HR should afford essential information to understand these process in mammalian cells.

## Acknowledgements

This work was supported by CEA, CNRS, ARC (1366) and Electricité de France. We thank C. White and C. Ducray for helpful comments.

## References

- [1] S.A. Liebhaber, M. Goosens, Y.-K. Kan, Homology and concerted evolution at the  $\alpha 1$  and  $\alpha 2$  loci of human  $\alpha$ -globin, *Nature* 290 (1981) 26–29.
- [2] D. Baltimore, Gene conversion: some implications for immunoglobulin Genes, *Cell* 26 (1981) 295–296.
- [3] J.E. Haber, Mating type gene switching in *Saccharomyces cerevisiae*, *TIG* 8 (1992) 446–452.
- [4] L.H.T. Van der Ploeg, Antigenic variation in African trypanosome, *TIG* 8 (1992) 452–457.
- [5] C.A. Reynaud, V. Anquez, H. Grimal, J.-C. Weill, A hyperconversion mechanism generates the chicken light chain preimmune repertoire, *Cell* 48 (1987) 379–388.
- [6] R.S. Becker, K.L. Knight, Somatic diversification of immunoglobulin heavy chain VDJ genes: evidence for somatic gene conversion in rabbits, *Cell* 63 (1990) 987–997.
- [7] S.G. Roeder, Chromosome synapsis and genetic recombination, *TIG* 6 (1990) 385–389.
- [8] N. Kleckner, Meiosis: how could it work?, *Proc. Natl. Acad. Sci. U.S.A.* 93 (1996) 8167–8174.
- [9] E.C. Friedberg, G.C. Walker, W. Siede, *DNA Repair and Mutagenesis*, ASM Press, Washington, DC, 1995.
- [10] R.J. Bollag, A.S. Waldman, R.M. Liskay, Homologous recombination in mammalian cells, *Annu. Rev. Genet.* 23 (1989) 199–225.
- [11] M. Amor, L.K. Parker, H. Globerman, M.L. New, P.C. White, Mutation in the CYP21B gene (Ile-172-Asn) causes steroid 21-hydroxylase deficiency, *Proc. Natl. Acad. Sci. U.S.A.* 85 (1988) 1600–1604.
- [12] Y. Wang, V.M. Maher, R.M. Liskay, J.J. McCormick, Carcinogens can induce homologous recombination between duplicated chromosomal sequences in mouse L cells, *Mol. Cell. Biol.* 8 (1988) 196–202.
- [13] D. Hellgren, Mutagen-induced recombination in mammalian cells in vitro, *Mut. Res.* 284 (1992) 37–51.
- [14] D.L. Ludwig, J.R. Stringer, Spontaneous and induced homologous recombination between *lacZ* chromosomal direct repeats in CV-1 cells, *Somat. Cell. Mol. Genet.* 20 (1994) 11–25.
- [15] H. Zhang, T. Tsujimura, N.P. Bhattacharyya, V.M. Maher, J.J. McCormick, O6-methylguanine induces intrachromosomal homologous recombination in human cells, *Carcinogenesis* 17 (1996) 2229–2235.
- [16] T. Tsujimura, V.M. Maher, A.R. Godwin, R.M. Liskay, J.J. McCormick, Frequency of intrachromosomal homologous recombination induced by UV radiation in normally repairing and excision repair-deficient human cells, *Proc. Natl. Acad. Sci. U.S.A.* 87 (1990) 1566–1570.
- [17] N.P. Bhattacharyya, V.M. Maher, J.J. McCormick, Effect of nucleotide excision repair in human cells on intrachromosomal homologous recombination induced by UV and 1-nitrosopyrene, *Mol. Cell. Biol.* 10 (1990) 3945–3951.
- [18] S.M. Meyn, High spontaneous intrachromosomal recombination rates in ataxia–telangiectasia, *Science* 260 (1993) 1327–1330.

- [19] P. Bertrand, D. Rouillard, A. Boulet, C. Levalois, T. Soussi, B.S. Lopez, Increase of spontaneous intrachromosomal homologous recombination in mammalian cells expressing a mutant p53 protein, *Oncogene* 14 (1997) 117–1122.
- [20] K.L. Mekeel, W. Tang, L.A. Kachnic, C.-M. Luo, J.S. DeFrank, S.N. Powell, Inactivation of p53 results in high rates of homologous recombination, *Oncogene* 14 (1997) 1847–1857.
- [21] H.-W. Sturzbecher, B. Donzelmann, W. Henning, U. Knippschild, S. Buchhop, p53 is linked directly to homologous recombination processes via RAD51/RecA protein interaction, *EMBO J.* 15 (1996) 1992–2002.
- [22] S. Buchhop, M.K. Gibson, X.W. Wang, P. Wagner, H.-W. Sturzbecher, C.C. Harris, Interaction of p53 with the human Rad51 protein, *Nucleic Acids Res.* 25 (1997) 3868–3874.
- [23] R. Scully, J. Chen, A. Plug, Y. Xiao, D. Weaver, J. Feunteun, T. Ashley, D.M. Livingston, Association of BRCA1 with Rad51 in mitotic and meiotic cells, *Cell* 88 (1997) 265–275.
- [24] R. Mizuta, J.M. LaSalle, H.-L. Cheng, A. Shinohara, H. Ogawa, N. Copeland, N.A. Jenkins, M. Lalonde, F.W. Alt, RAB22 and RAB163/mouse BRCA2: proteins that specifically interact with the RAD51 protein, *Proc. Natl. Acad. Sci. U.S.A.* 94 (1997) 6927–6932.
- [25] L.Y. Marmorstein, T. Ouchi, S.A. Aaronson, The BRCA2 gene product functionally interacts with p53 and RAD51, *Proc. Natl. Acad. Sci. U.S.A.* 95 (1998) 13869–13874.
- [26] H.L. Klein, Genetic control of intrachromosomal recombination, *BioEssays* 17 (1995) 147–159.
- [27] E.L. Ivanov, N. Sugawara, J. Fishman-Lobell, J.E. Haber, Genetic requirements for the single-strand annealing pathway of double-strand break repair in *Saccharomyces cerevisiae*, *Genetics* 142 (1996) 693–704.
- [28] N. Sugawara, F. Pâques, M. Colaiàcovo, J.E. Haber, Role of Msh2 and Msh3 repair proteins in double-strand break induced recombination, *Proc. Natl. Acad. Sci. U.S.A.* 94 (1997) 9214–9219.
- [29] R.M. Liskay, J.L. Stachelek, A. Letsou, Homologous recombination between repeated chromosomal sequences in mouse cells, *Cold Spring Harbor Symp. Quant. Biol.* 49 (1984) 183–189.
- [30] R.J. Bollag, R.M. Liskay, Conservative intrachromosomal recombination between inverted repeats in mouse cells: association between reciprocal exchange and gene conversion, *Genetics* 119 (1988) 161–169.
- [31] A. Letsou, R.M. Liskay, Effect of the molecular nature of mutation on the efficiency of intrachromosomal gene conversion in mouse cells, *Genetics* 117 (1987) 759–769.
- [32] F.L. Lin, K. Sperle, N. Sternberg, Model for homologous recombination during transfer of DNA into mouse L cells: role for DNA ends in the recombination process, *Mol. Cell. Biol.* 4 (1984) 1020–1034.
- [33] S. Chakrabarti, M.M. Seidman, Intramolecular recombination between transfected sequences in mammalian cells is non-conservative, *Mol. Cell. Biol.* 6 (1986) 2520–2526.
- [34] F. Lin, K. Sperle, N. Sternberg, Extrachromosomal recombination in mammalian cells as studied with single- and double stranded DNA substrates, *Mol. Cell. Biol.* 7 (1987) 129–140.
- [35] R. Holliday, A mechanism for gene conversion in fungi, *Genet. Res.* 5 (1964) 282–306.
- [36] M.S. Meselson, C.M. Radding, A general model for genetic recombination, *Proc. Natl. Acad. Sci. U.S.A.* 72 (1975) 358–361.
- [37] J.W. Szostak, T.L. Orr-Weaver, R.J. Rothstein, F. Stahl, The double-strand-break repair model for recombination, *Cell* 33 (1983) 25–35.
- [38] J. Rommelaere, A. Miller-Faures, Detection by density equilibrium centrifugation of recombinant-like DNA molecules in somatic mammalian cells, *J. Mol. Biol.* 98 (1975) 195–218.
- [39] P.D. Moore, R. Holliday, Evidence for the formation of hybrid DNA during mitotic recombination in Chinese hamster cells, *Cell* 8 (1976) 573–579.
- [40] B. Lopez, S. Rousset, J. Coppey, Homologous recombination intermediates between two duplex DNA catalysed by human cell extracts, *Nucleic Acids Res.* 15 (1987) 5643–5655.
- [41] R.J. Bollag, D.R. Elwood, E.D. Tobin, A.R. Godwin, R.M. Liskay, Formation of heteroduplex DNA during mammalian intrachromosomal gene conversion, *Mol. Cell. Biol.* 12 (1992) 1546–1552.
- [42] A.R. Godwin, R.J. Bollag, D.M. Christie, R.M. Liskay, Spontaneous and restriction enzyme-induced chromosomal recombination in mammalian cells, *Proc. Natl. Acad. Sci. U.S.A.* 91 (1994) 12554–12558.
- [43] R.M. Liskay, A. Letsou, J.L. Stachelek, Homology requirements for efficient gene conversion between duplicated chromosomal sequences in mammalian cells, *Genetics* 115 (1987) 161–167.
- [44] A.S. Waldman, R.M. Liskay, Differential effect of base-pair mismatch on intrachromosomal versus extrachromosomal recombination in mouse cells, *Proc. Natl. Acad. Sci. U.S.A.* 84 (1987) 5340–5344.
- [45] A.S. Waldman, R.M. Liskay, Dependence of intrachromosomal recombination in mammalian cells on uninterrupted homology, *Mol. Cell. Biol.* 8 (1988) 5350–5357.
- [46] D. Yang, A.S. Waldman, Fine-resolution analysis of products of intrachromosomal homeologous recombination in mammalian cells, *Mol. Cell. Biol.* 17 (1997) 3614–3628.
- [47] B.S. Lopez, P. Bertrand-Mercat, E. Corteggiani, J. Coppey, Directional recombination is initiated at a double strand break in human nuclear extracts, *Nucleic Acids Res.* 20 (1992) 501–506.
- [48] J.A. Blaho, R.D. Wells, Left-handed Z-DNA and genetic recombination, *Prog. Nucleic Acid Res. Mol. Biol.* 37 (1989) 107–126.
- [49] W.P. Wahls, P.D. Moore, Homologous recombination enhancement conferred by the Z-DNA motif d(TG)<sub>30</sub> is abrogated by Simian Virus 40 T antigen binding to adjacent DNA sequences, *Mol. Cell. Biol.* 10 (1990) 794–800.
- [50] W.P. Wahls, L.J. Wallas, P.D. Moore, Hypervariable minisatellite DNA is a hotspot for homologous recombination in human cells, *Cell* 60 (1990) 95–103.
- [51] R.G. Sargent, R.V. Merrihew, R. Nairn, G. Adair, M. Meuth, J.H. Wilson, The influence of a (GT)<sub>29</sub> microsatellite se-

- quence on homologous recombination in the hamster adenine phosphoribosyl transferase gene, *Nucleic Acids Res.* 24 (1996) 746–753.
- [52] B.J. Thomas, R. Rothstein, Elevated recombination rates in transcriptionally active DNA, *Cell* 56 (1989) 619–630.
- [53] K. Voelkel-Meiman, R.L. Keil, G.S. Roeder, Recombination-stimulating sequences in Yeast Ribosomal DNA correspond to sequences regulating transcription by RNA polymerase I, *Cell* 48 (1987) 1071–1079.
- [54] J.A. Nickoloff, R.J. Reynolds, Transcription stimulates homologous recombination in mammalian cells, *Mol. Cell. Biol.* 10 (1990) 4837–4845.
- [55] J.A. Nickoloff, Transcription enhances intrachromosomal homologous recombination in mammalian cells, *Mol. Cell. Biol.* 12 (1992) 5311–5318.
- [56] W.P. Deng, J.A. Nickoloff, Preferential repair of UV damage in highly transcribed DNA diminishes UV-induced intrachromosomal recombination in mammalian cells, *Mol. Cell. Biol.* 14 (1994) 391–399.
- [57] D.G. Taghian, J.A. Nickoloff, Chromosomal double-strand breaks induce gene conversion at high frequency in mammalian cells, *Mol. Cell. Biol.* 17 (1997) 6386–6393.
- [58] M.B. Benjamin, J.B. Little, X-rays induce interallelic homologous recombination at the human thymidine kinase gene, *Mol. Cell. Biol.* 12 (1992) 2730–2738.
- [59] M.S. Park, Expression of human RAD52 confers resistance to ionizing radiation in mammalian cells, *J. Biol. Chem.* 270 (1995) 15467–15470.
- [60] M. Brenneman, F.S. Gimble, J.H. Wilson, Stimulation of intrachromosomal homologous recombination in human cells by electroporation with site-specific endonuclease, *Proc. Natl. Acad. Sci. U.S.A.* 93 (1996) 3608–3612.
- [61] M. Jasin, Genetic manipulation of genome with rare-cutting endonucleases, *TIG* 12 (1996) 224–228.
- [62] P. Rouet, M. Jasin, Introduction of double-strand breaks into the genome of mouse cells by expression of a rare-cutting endonuclease, *Mol. Cell. Biol.* 14 (1994) 8096–8106.
- [63] P. Rouet, F. Sih, M. Jasin, Expression of a site-specific endonuclease stimulates homologous recombination in mammalian cells, *Proc. Natl. Acad. Sci. U.S.A.* 91 (1994) 6064–6068.
- [64] G.R. Sargent, M.A. Brenneman, J.H. Wilson, Repair of site-specific double-strand breaks in a mammalian chromosome by homologous and illegitimate recombination, *Mol. Cell. Biol.* 17 (1997) 267–277.
- [65] F. Liang, M. Han, P.J. Romanienko, M. Jasin, Homology-directed repair is a major double-strand break repair pathway in mammalian cells, *Proc. Natl. Acad. Sci. U.S.A.* 95 (1998) 5172–5177.
- [66] C. Bonnerot, J.-F. Nicolas, Clonal analysis in the intact mouse embryo by intragenic homologous recombination, *C. R. Acad. Sci. Paris* 316 (1993) 1207–1217.
- [67] J.R. Murti, M. Bumbulis, J.C. Schimenti, High-frequency germ line gene conversion in transgenic mice, *Mol. Cell. Biol.* 12 (1992) 2545–2552.
- [68] F.-L. Lin, N. Sternberg, Homologous recombination between overlapping thymidine kinase gene fragments stably integrated into mouse cell genome, *Mol. Cell. Biol.* 4 (1984) 852–861.
- [69] M.J. Shulman, C. Collins, A. Connor, L.R. Read, M.D. Baker, Interchromosomal recombination is suppressed in mammalian somatic cells, 14 (1995) 4102–4107.
- [70] A.S. Waldman, B.C. Waldman, Stimulation of intrachromosomal homologous recombination in mammalian cells by inhibitors of poly(ADP-ribosylation), *Nucleic Acids Res.* 19 (1991) 5943–5947.
- [71] M.S. Park, D.L. Ludwig, E. Stigger, S.-H. Lee, Physical interaction between human RAD52 and RPA is required for homologous recombination in mammalian cells, *J. Biol. Chem.* 271 (1996) 18996–19000.
- [72] S. Vispé, C. Cazaux, C. Lesca, M. Defais, Overexpression of Rad51 protein stimulates homologous recombination and increases resistance of mammalian cells to ionizing radiation, *Nucleic Acids Res.* 26 (1998) 2859–2864.