Proteogenomic insights into uranium tolerance of a Chernobyl's *Microbacterium* bacterial isolate

Nicolas Gallois, Béatrice Alpha-Bazin, Philippe Ortet, mohamed Barakat, Laurie Piette, Justine Long, C. Berthomieu, J. Armengaud, Virginie Chapon

► To cite this version:

Nicolas Gallois, Béatrice Alpha-Bazin, Philippe Ortet, mohamed Barakat, Laurie Piette, et al.. Proteogenomic insights into uranium tolerance of a Chernobyl's *Microbacterium* bacterial isolate. Journal of Proteomics, 2018, 177, pp.148-157. 10.1016/j.jprot.2017.11.021 . cea-01944196

HAL Id: cea-01944196

<https://hal-cea.archives-ouvertes.fr/cea-01944196>

Submitted on 5 Dec 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Proteogenomic insights into uranium tolerance of a Chernobyl's

Microbacterium bacterial isolate

Gallois Nicolas¹, Alpha-Bazin Béatrice², Ortet Philippe³, Barakat Mohamed³, Piette Laurie¹, Long Justine³, Berthomieu Catherine¹, Armengaud Jean^{2#}, Chapon Virginie¹

¹CEA, CNRS, Aix-Marseille Université, UMR 7265 Biologie Végétale et Microbiologie

Environnementales, Laboratoire des Interactions Protéine Métal, 13108 Saint-Paul-lez-Durance, France.

²Laboratoire Innovations technologiques pour la Détection et le Diagnostic (Li2D), Service de

Pharmacologie et Immunoanalyse (SPI), CEA, INRA, F-30207 Bagnols sur Cèze, France.

³CEA, CNRS, Aix-Marseille Université, UMR 7265 Biologie Végétale et Microbiologie

Environnementales, Laboratoire d'écologie microbienne de la rhizosphère et d'environnements extrêmes, 13108 Saint-Paul-lez-Durance, France

[#]Corresponding author at:

Laboratory «Innovative technologies for Detection and Diagnostics», CEA-Marcoule, DRF-Li2D, PRAE

Marcel Boiteux – Départementale 765, BP 17171, F-30200 Bagnols-sur-Cèze cedex, France

Tel: +33 4 66 79 62 77; Fax: +33 4 66 79 64 60.

E-mail address: jean.armengaud@cea.fr

URL: <http://ibitecs.cea.fr/dsv/ibitecs/Pages/services/spi/li2d.aspx>

Keywords

Radionuclides, toxicity, proteogenomics, bioinformatics, toxicoproteomics, *Microbacterium*

Abstract

Microbacterium oleivorans A9 is a uranium-tolerant actinobacteria isolated from the trench T22 located near the Chernobyl nuclear power plant. This site is contaminated with different radionuclides including uranium. To observe the molecular changes at the proteome level occurring in this strain upon uranyl exposure and understand molecular mechanisms explaining its uranium tolerance, we established its draft genome and used this raw information to perform an in-depth proteogenomics study. High-throughput proteomics were performed on cells exposed or not to 10 μ M uranyl nitrate sampled at three previously identified phases of uranyl tolerance. We experimentally detected and annotated 1,532 proteins and highlighted a total of 591 proteins for which abundances were significantly differing between conditions. Notably, proteins involved in phosphate and iron metabolisms show high dynamics. A large ratio of proteins more abundant upon uranyl stress, are distant from functionally-annotated known proteins, highlighting the lack of fundamental knowledge regarding numerous key molecular players from soil bacteria.

Biological significance

Microbacterium oleivorans A9 is an interesting environmental model to understand biological processes engaged in tolerance to radionuclides. Using an innovative proteogenomics approach, we explored its molecular mechanisms involved in uranium tolerance. We sequenced its genome, interpreted high-throughput proteomic data against a six-reading frame ORF database deduced from the draft genome, annotated the identified proteins and compared protein abundances from cells exposed or not to uranyl stress after a cascade search. These data show that a complex cellular response to uranium occurs in *Microbacterium oleivorans* A9, where one third of the experimental proteome is modified. In particular, the uranyl stress perturbed the phosphate and iron metabolic pathways. Furthermore, several transporters have been identified to be specifically associated to

uranyl stress, paving the way to the development of biotechnological tools for uranium decontamination.

1. Introduction

Radionuclides (RNs) are naturally present in the environment but are redistributed by anthropogenic activities *e.g.* chemical nitrogen fertilizer use, weapons manufacturing, nuclear research and nuclear fuel production. Several major nuclear accidents such as Chernobyl and Fukushima led to the dispersion of several RNs through the atmosphere in microparticles [1] and in the soil [2]. In Chernobyl, radionuclides-contaminated wastes were buried on site in about 800 trenches. Besides ^{137}Cs and ^{90}Sr , other RNs are present in the Chernobyl trench T22, including ^{60}Co , ^{154}Eu , $^{238, 239, 240}\text{Pu}$, ^{241}Am and $^{235, 238}\text{U}$.

Uranium (U) belongs to the actinide series. It exhibits both radiological and chemical toxicities, and a long half-life, resulting in environmental and human health major concern [3]. As a hard Lewis acid and according to the HSAB theory which qualitatively explains the stability of metal complexes [4], uranyl tends to interact with hard Lewis base such as phosphate, sulfate, carbonate or amine by ionic interactions. It can displace other hard Lewis acids like Ca^{2+} and Fe^{3+} in highly oxygenated site. Moreover, different studies have confirmed the affinity between uranium and phosphate or carbonate [5-9], or the replacement of iron by uranyl [10].

Bacteria can interact directly or indirectly with RNs and change their speciation, playing a major role in their mobility and transfer in the environment. The use of both molecular- and culture-based approaches to study bacterial diversity in the trench T22 demonstrated that RNs-contaminated soils host an unexpected diversity of bacteria [11].

During their sampling campaign at Chernobyl, Chapon and colleagues have constructed a large collection of cultured bacteria from the trench T22 soils offering the opportunity to study the RN

tolerance of these bacteria [11]. The strain *Microbacterium oleivorans* A9 (referred as *Microbacterium* sp. A9_sp3_-1_2 in [12]) is one of the most uranium-tolerant isolate of this collection. In a previous paper, the interactions between uranium and this bacterium have been investigated and specific exposure conditions in which the bacteria were kept alive while being exposed to soluble forms of uranium have been set-up. With this tightly controlled exposure conditions, it has been shown that *Microbacterium oleivorans* A9 exhibit three sequential mechanisms involved in uranium detoxification: a rapid metal removal within the first 30 minutes, then an active U(VI) release in the exposure medium accompanied by a phosphate efflux and a final biomineralization step of uranium in autunite-like mineral phases [12]. For this reason, it constitutes a relevant model to study the mechanisms involved in uranium tolerance and in uranium-bacteria interactions. In bacteria, survival in metal-contaminated environment is mainly achieved through active efflux pumps [13]. If such systems are well-known for a wide variety of non-essential metals, involvement of an efflux system in detoxification of uranium has not yet been described, although up-regulation of genes encoding metal efflux pumps has already been reported for *Desulfotomaculum reducens* exposed to U(VI) in anaerobic conditions [14].

The cellular response of *Escherichia coli*, *Acidithiobacillus ferrooxidans* and two *Anabaena* strains upon uranium exposure has already been studied with 2D-PAGE based proteomic methodology [15-17]. These studies highlighted an impact of uranium on the bacterial proteome, but the low throughput of this approach allowed identifying only 11, 18, 45 and 27 proteins respectively. These may represent only a partial view of the proteins modulated upon uranium stress. Two studies using high-throughput tandem mass spectrometry have deepened the proteome changes occurring in two other bacteria, *Geobacter sulfurreducens* and *Caulobacter crescentus* [18, 19]. These papers show that resistance to uranium is a complex cellular response with induction of multiple stress response systems. These four studies, based on gram-negative organisms, have used different uranium concentrations (from 50 to 500 μ M) as well as different exposure conditions making any comparison difficult.

Proteogenomics, the alliance of genomics and proteomics, has proved its efficiency in supporting the existence of genes encoding hypothetical proteins or with unknown function, and even unannotated in the genome for several organisms [20]. This methodology is even more appropriate when a quick focus on mechanisms for which the main players are unknown is required [21-23]. Because, proteogenomics is based on the assignment of MS/MS spectra with a six-frame translation of the genome, time-consuming, resource-dependent and hard-working genome sequence completion steps are bypassed. For example, the annotation of *Deinococcus deserti* by means of proteogenomics has shown its added value to correct annotation errors in other *Deinococcus* sp. genomes [21, 24]. The exploration of the halotolerance molecular mechanisms of *Tistlia consotensis* by proteogenomics based on a draft genome sequence straightforwardly highlighted major changes in response to hypo-osmotic and hyper-osmotic conditions [22, 25].

Here, the molecular changes occurring in *Microbacterium oleivorans* A9 upon uranium exposure were deciphered by means of a proteogenomics approach consisting in draft genome sequencing and high-throughput proteome coverage by next-generation proteomics. Proteomic changes of cells exposed to 10 μ M uranyl for 30 min, 4 h and 24 h were investigated. This experimental strategy allowed identification without *a priori* of soluble and membrane proteins with significantly differing abundances compared to controls, which could be players of the defense mechanism of *Microbacterium oleivorans* A9 against uranium. Furthermore, these genome and proteome datasets represent invaluable resources to gain insights into the physiology and role of *Microbacterium* in soils.

2. Experimental procedures

2.1 Uranium exposure

The experimental procedure used to expose the bacteria to uranium is described in [12]. *Microbacterium oleivorans* A9 strain was routinely cultivated in 0.1 X Tryptic Soy Broth (TSB, Difco Laboratories) at 30°C with shaking. Cells at exponential growth phase were harvested by centrifugation for 10 min at 5000 *g*. From this stage on, samples were maintained at 25°C throughout the experiment. The resulting cell pellets were washed twice in 0.1 M NaCl pH 5.0 and were re-suspended at about 6×10^9 cells mL⁻¹ in 0.1 M NaCl pH 5.0 with 0 (control) or 10 µM U(VI). U(VI) was added as uranyl nitrate UO₂(NO₃)₂·6H₂O (Sigma–Aldrich) from a 7.51 mM stock solution in 16 mM HNO₃. No precipitate was observed along the exposure conditions. The nitrate concentration was adjusted to 0.416 mM by adding NaNO₃ when needed. Bacteria exposed to U(VI) and controls were incubated at 25°C with shaking. Four independent biological replicates were made for each condition and for each time point. Fractions of 1 ml of cell suspension were taken after 0.5, 4, and 24 h of uranium exposure. Samples were centrifuged for 5 min at 8000 *g*. The resulting supernatants were removed and the pellets were conserved at -80 °C until proteomic analysis.

2.2 Genome of *Microbacterium oleivorans* A9

Genomic DNA was sequenced on a HiSeq 2000 sequencing platform (Illumina) by the GenoScreen Company and genome *de novo* assembling was performed on the reads using ABYSS [26]. The Whole Genome Shotgun project data have been deposited at DDBJ/EMBL/GenBank under the accession number MTIO000000000. For each contig, the ORF translated sequences were extracted from stop codon to stop codon in each of the six possible reading frames. The ORF-derived polypeptide list was restricted to those exhibiting a length of at least 50 amino acids. Proteomic data and proteogenomic procedures were then used to discriminate between false, correct but undetected, and correct and detected protein sequences as described earlier [21, 27]. The ORF sequences for which at least one spectrum has been assigned were kept to create a second database. This step allowed us to identify

more proteins and to improve qualitatively and quantitatively the MS/MS spectra attribution. The protein sequences were given a cluster of orthologous groups (COG) assignment based on the reference database [28]. Furthermore, enzymes annotation using PRIAM profiles [29] associated to KEGG database [30] allowed the identification of metabolic pathways. An in-house batch program was developed to emphasize metabolic pathways highlighted by proteomic analysis.

2.3 Proteome sample preparation and trypsin in-gel proteolysis

Cell pellets (between 2.0 and 4.6 mg, 3.1 mg average, wet weight) were dissolved in a given volume (100 μ L for 1.7 mg of bacterial pellet) of 1X NuPAGE™ LDS Sample Buffer (Invitrogen) supplemented with 2% β -mercaptoethanol. Samples were sonicated for 5 min in a transonic 780H sonicator and subjected to a 5 min incubation at 99°C prior to SDS-PAGE. Samples were subjected to a 3 min short migration in denaturing conditions as previously described [31]. For this, samples were loaded onto a 4-12% gradient 10-well NuPAGE Bis-Tris (Invitrogen) gel operated with NuPAGE MES (Invitrogen) as running buffer at 200 V. After gel staining with Simply Blue Safe Stain, a ready-to-use Coomassie G-250 stain from Invitrogen, the whole proteome content (comprising both soluble and membrane proteins) from each sample was excised as a single piece of gel polyacrylamide band. The 24 resulting polyacrylamide bands were washed with MilliQ water and dehydrated with $\text{CH}_3\text{CN}:\text{NH}_4\text{HCO}_3$ 50mM (1:1 v:v) before addition of 100% CH_3CN . Gel pieces were then dried for 2 to 5 min under vacuum. For in-gel digestion, dry gel pieces were rehydrated with 100 mM NH_4HCO_3 containing 25 mM DTT and incubated for 10 min at 56°C. After removal of the solution, the gel pieces were further treated with 100 mM NH_4HCO_3 containing 55 mM iodoacetamide at room temperature and dried under vacuum as before. The gel bands were then processed for in-gel proteolysis with trypsin in presence of 0.01% ProteaseMax (Promega) as previously described [32].

2.4 NanoLC-MS/MS analysis

NanoLC–MS/MS experiments for the 24 resulting peptide mixtures were performed using a Q-Exactive HF mass spectrometer (ThermoFisher) coupled to an UltiMate 3000 LC system (Dionex-LC

Packings) in similar conditions as those previously described [33]. Peptide mixtures (10 µl) were loaded and desalted on-line on a reverse phase precolumn (Acclaim PepMap 100 C18, 5 µm bead size, 100 Å pore size, 5 mm × 300 µm id) from LC Packings. Peptides were then resolved onto a reverse phase Acclaim PepMap 100 C18 column (3 µm, 100 Å, 500 mm x 75 µm id) at a flow rate of 0.3 µl/min with a 180 min linear gradient of CH₃CN/0.1% formic acid and injected into the Q-Exactive HF mass spectrometer. The Q-Exactive HF instrument was operated according to a Top20 data-dependent acquisition method as previously described [33]. Briefly, a scan cycle was initiated with a full scan of peptide ions in the ultra-high-field Orbitrap analyzer, followed by selection of a single precursor and its dissociation in high energy collisional mode, and MS/MS scans on the 20 most abundant precursor ions. Full scan mass spectra were acquired with an Automatic Gain Control Target set at 3x10⁶ ions and a resolution of 60,000 from *m/z* 350 to 1,800. MS/MS scan was initiated at a resolution of 15,000 when the ACG target reached 1x10⁵ ions with a threshold intensity of 83,000 and potential charge states of 2+ and 3+. Precursor ions were selected with a dynamic exclusion of 10 sec for increasing the reliability of spectral count measurements.

2.5 Interpretation of mass spectrometry data

The recorded MS/MS spectra for the 24 samples were merged after being searched against the home-made ORF database. First, peak lists were generated with the MASCOT DAEMON software (version 2.3.2) from Matrix Science using the extract_msn.exe data import filter from the Xcalibur FT package (version 2.0.7) proposed by ThermoFisher. Data import filter options were set as previously described [34] at 400 (minimum mass), 5000 (maximum mass), 0 (grouping tolerance), 0 (intermediate scans), and 1000 (threshold). MS/MS spectra were searched with MASCOT against the ORF database with the following parameters: tryptic peptides with a maximum of 2 miss cleavages during proteolytic digestion, a mass tolerance of 5 ppm on the parent ion and 0.02 Da on the MS/MS, fixed modification for carbamidomethylated Cys (+57.0215) and variable modification for oxidized Met (+15.9949). All peptide matches with a peptide score above its query threshold set at $p \leq 0.05$ with the ORF database and rank 1 were parsed using the IRMa 1.31.1c software [35]. MS/MS spectra

assigned to several loci were systematically removed. The average error for the determination of peptide mass is 0.59 ppm and the mean MASCOT score is 50.7. A protein was considered validated when at least two different peptides were detected. False-positive identification of proteins was estimated using a reverse decoy database as below 0.1% with these parameters.

The mass spectrometry proteomics data have been deposited in a public repository at the ProteomeXchange Consortium (<http://proteomecentral.proteomexchange.org>) via the PRIDE partner repository [36] with the dataset identifier PXD005794 and project DOI 10.6019/PXD005794.

The number of MS/MS spectra per protein (spectral counts) was determined for the four replicates of each of the three time points for both conditions. The protein abundances were compared for each time point between the uranyl exposure and the control conditions. For each of these comparisons, the list of non-redundant proteins detected among the six corresponding datasets was established. The total spectral count of each polypeptide was used to rank the proteins from the highest to the lowest detection intensities. The statistical protein variation among the four replicates samples of the two specific exposure conditions compared was calculated using the T-Fold option of the PatternLab 2.0 software [37]. This module allows normalizing the spectral count datasets, calculating the average fold changes with statistics (t-test), and estimating the resulting theoretical false discovery rate. MS/MS data were compiled in Excel (Microsoft) and converted for PatternLab with an Excel home-designed macro. Normalization was done taking into account the total number of spectral count for each sample, taking at least two readings per protein. A minimum value of 1 was added systematically to all spectral count values in order to consider missing values as a standard PatternLab normalization. Parameters for the comparisons were as follows: minimum fold change of 1.5, minimum p-value of 0.05 and BH-FDR Alfa of 0.15. Normalized spectral abundance factor (NSAF) for each protein was calculated using the formula: (spectral counts / theoretical molecular weight) x 1000, see [34].

3. Results and discussion

3.1 *Microbacterium oleivorans* A9 genome annotation

The main characteristics of *Microbacterium oleivorans* A9 genome is described in [38]. Briefly, the genome size is around 2.95 Mbp. The G+C content of the genome is 68.33%. COG assignment for the resulting polypeptide dataset shows that two classes are predominant: proteins involved in metabolism of amino acids (74 proteins) on the first hand, and those involved in metabolism of carbohydrates (66 proteins) on the other. A search against the BacMet database [39], which comprises antibacterial biocides and metal resistance genes, highlights the presence of 104 unique metal resistance genes such as *actP* gene which encodes a copper-transporting P-type ATPase, *arsB* gene which encodes an arsenic pump membrane protein or *yieF* gene which encodes a chromate reductase. We found a 16S rRNA gene copy in the genome, exhibiting 99% sequence identity with *Microbacterium oleivorans* 16S rRNA gene. Moreover, the assembling results are strikingly similar to those obtained for *Microbacterium oleivorans* strain RIT293 (RefSeq NZ_JFY000000000.1) [40] with a median total length of 2.9 Mbp, a G+C content of 69% and a total of 2,732 protein-coding genes. This genome version comprises 11 contigs with a N50 parameter at 467,109 bp.

3.2 Shotgun proteogenomics of *Microbacterium oleivorans* A9

Proteogenomics quick panorama

Figure 1 shows the strategy applied here for quickly deciphering the proteome dynamics from resting cells exposed or not to 10 μ M uranium. A total of 1,221,282 MS/MS spectra were recorded. Most of these spectra were of high quality as revealed by the high percentage of assignment obtained (61.1%) at p-value 0.05. For interpreting MS/MS spectra, a six reading frame translated ORF database was created. This proteogenomics database comprised a total of 30,853 putative polypeptide sequences, totaling 4,903,573 amino acids with an average of 159 amino acids per putative polypeptide and 2,064 as longest length. A total of 746,092 MS/MS spectra could be assigned in first intention, allowing certifying the presence of 1,504 proteins detected with at least two different

peptides (Supplementary data Table S1). Because of the unusual large size of the six reading frame ORF database (30,853 putative polypeptide sequences instead of 2813 annotated coding DNA sequences), a second round of search was initiated after reducing the database to the proteins detected in the first round as proposed previously for improving the sensitivity of proteogenomics studies [41]. The second search led to the assignment of 747,621 peptide-to-spectrum matches, certifying the detection of 17,027 unique peptide sequences and 1,532 non-redundant proteins identified with at least two different peptides (Supplementary data Table S2). This two-round search allows increasing the number of proteins validated with at least two different peptides (1532 instead of 1504) and enhances the number of assigned MS/MS spectra (747 621 instead of 746 092). This dataset compares favorably with previous studies of bacteria exposed to uranium, as a dataset of less than 950 proteins were reported from *Caulobacter crescentus* [19] and 1,363 proteins were listed from *Geobacter sulfurreducens* [18]. The proteomic dataset obtained here represents 54% (including 15% of membrane proteins) of the predicted protein-coding genes found in the draft genome of *Microbacterium oleivorans* A9 [38]. The distribution of the peptides detected during our experiment along the chimeric genome of *Microbacterium oleivorans* A9 is uniform and confirm the validity of this proteogenomics approach, as previously discussed for the proteogenomics mapping of the alphaproteobacterium *Tistlia consotensis* [22].

Functional categorization of the detected proteins and relative abundances

The dataset of 1,532 identified proteins were classified into 22 COG categories. **Table 1** shows the functional classification of the mass-spectrometry certified global proteome. Most of the detected proteins (50%) are distributed in five classes: hypothetical proteins (15%), amino acid metabolism (11%), carbohydrate metabolism (9%), translation, ribosomal structure, and biogenesis (9%) and transcription (6%). The remaining 17 classes represent less than 5% each (**Table 1**). The prevalence of the same classes has been shown in previous studies [18, 19], but here, the large number of uncharacterized proteins is indicative of an important lack of knowledge regarding soil gram-positive bacteria. 36 proteins account for 25% of the total proteome detected, and thus are the key molecular

draft horses within the cells. These 36 most abundant proteins detected throughout the study as evaluated with their normalized spectral abundance factor (NSAF) are given in Supplementary data Table S2. The most abundant protein is the HU bacterial nucleoid DNA-binding protein involved in DNA structuration. Two of the top five most abundant proteins are the chaperonin GroEL (1.28%) and the co-chaperonin GroES (1.06%). These two proteins form the binary GroEL/GroES complex, which mediates proper folding of many proteins. The sub-unit GroEL is specifically known for its important role both in normal and stressful conditions as well as in the resistance towards toxic metals [42]. As expected the ribosomal proteins are abundant since the cells were collected during the exponential growth phase. The 100 most abundant proteins represent 46.1% of total NSAF. Amongst these, 16 proteins (8.3% of total NSAF) are components of the small ribosomal subunit and 27 proteins (10.6% total NSAF) belong to the large ribosomal subunit. This abundance of ribosomal proteins has already been observed [23, 34]. Within the most abundant proteins, several are involved in energy metabolism such as the F0F1-type ATP synthase (0.78% and 0.57% respectively).

3.3 Proteome dynamics upon uranium stress

Common differentially produced proteins over the time course of the experiment

Proteins with significant abundance changes (fold change ≥ 1.5 and p-value ≤ 0.05) in uranium-exposed condition versus control were listed at each time point. A total of 592 proteins met the criteria: 391 at 0.5h, 294 at 4h and 122 at 24h. The decrease of differentially produced proteins over time is due to the non-growing exposure conditions. **Figure 2** shows a Venn diagram representation where the numbers of these proteins are reported for each condition, highlighting the proteins specific of a given time point and those common in two or three time points. A relatively small set of 24 proteins were shared along the whole kinetic. Remarkably seven of these 24 proteins are predicted to be ABC-type transporters. Their spectrum activities appeared to be broad as they share significant sequence similarities with dipeptide/oligopeptide/nickel, Fe³⁺-hydroxamate, multidrug or enterocholin transporters. A serine protease and a DNA-binding transcriptional regulator belonging

294 to the PadR family are also found more abundant in presence of uranium for the three time points:
 295 1.7, 1.6 and 4.6-fold; 2.1-fold, 2.6-fold and 2.0-fold, respectively. Members of the PadR family
 296 regulate different pathways such as multidrug resistance and detoxification [43].
 297 A total of 128 proteins were found significantly more abundant in 0.5h and 4h samples of exposure
 298 to uranium compared to controls. These 128 proteins covered a large variety of molecular functions
 299 making difficult to draw specific uranium responsive pathways. Six ABC-type transporters were
 300 observed among this large panel of up-regulated proteins. They are predicted to be involved in
 301 amino acid, sugar, cobalamin, Fe^{3+} siderophore and glycerol-3-phosphate transport. These data
 302 reinforce the hypothesis that bacteria deploy a rather large panel of transport machineries to
 303 respond to uranium exposure. Four proteins involved in cell division are also significantly more
 304 abundant at these two initial phases of uranium exposure: the cell division proteins FtsI and FtsQ,
 305 and two chromosome partitioning ATPases. Such observation reinforces the idea that cells are viable
 306 and active.
 307 Once again, a link with iron metabolism is highlighted in the proteins common after 0.5h and 24h of
 308 uranium exposure with the presence of ABC-type Fe^{3+} /spermidine/putrescine transport system (2.0
 309 and 1.8-fold), the ferredoxin-NADP reductase (1.9 and 1.9-fold), the deferrochelataase/peroxidase
 310 EfeB (1.7 and 1.8-fold) and the NADPH-dependent ferric siderophore reductase (1.8 and 1.6-fold).
 311 Interestingly, a protein predicted to be a DNA-binding transcriptional regulator belonging to the ArsR
 312 family is found significantly more abundant at 0.5h but less abundant after 24h uranium exposure.
 313 This regulator is known to repress expression of stress-induced operons involved in toxic metal ion
 314 tolerance [44]. This suggests that *Microbacterium oleivorans* A9 first represses these genes when
 315 cells are at the initial stage of uranium contact, but need the expression of these genes when the
 316 uranium is intracellularly biomineralized and so less bioavailable.
 317 A comparison of the significant changes of protein abundances between 4h and 24h of exposure
 318 shows that 17 proteins are in common for these two time points. Three are related to
 319 osmoregulation (DNA-binding response regulator of the OmpR family and of the AcrR family,

maltooligosyltrehalose synthase) which is consistent with the exposure condition used *i.e.* NaCl 0.1M.

Proteins specifically modulated at a given time point

A large set of proteins (391) has been found significantly modulated in terms of abundance after 0.5h exposure to U(VI) as compared to the control. **Table 2** shows the proteins with the most important changes (fold-change ≥ 3.5). The chorismate mutase exhibited the highest fold-change (5.8-fold), while its homologue from *Geobacter sulfurreducens* has been previously shown to decrease in response to uranium [18]. This protein is involved in phenylalanine and tyrosine synthetic pathway through the conversion of chorismate in prephenate. As reported in **Figure 3**, five other proteins from the same pathway have an increased abundance upon uranium exposure, including the isochorismate synthase (2.6-fold) and the aminotransferase/4-amino-4-deoxychorismate lyase (2.0-fold). A protein involved in ABC-type glycerol-3-phosphate transport, two proteins responsible of acyl transfer to glycerol-3-phosphate and the CDP-glycerol glycerophosphotransferase were more abundant in uranium condition. Glycerol-3-phosphate appears as an important metabolite which may provide phosphate for complexing and immobilizing uranium [45]. Major changes affected the phosphate metabolism as evidenced by the important number (59) of proteins related to this metabolism. A large panel of phosphatases (16 polypeptides) is found in higher amounts after uranium stress. These enzymes are involved in phosphate removal mainly from sugar (fructose, trehalose) and from amino acids (serine, threonine).

Regarding the two other time points analyzed by high-throughput proteomics, the most modulated protein detected after 4h is also the protein with the higher abundance after 24h of uranium exposure compared to control (11-fold and 25-fold respectively). This protein shares some sequence similarities (63% identity) to the uncharacterized Ykol membrane protein of *Microbacterium oleivorans* LKL04 (**Table 2**). It presents a PepSY domain (pfam 03413) hypothetically involved in inhibition of a peptidase activity. Based on the TMHMM predictive tool [46], one transmembrane segment has been identified in the *Microbacterium oleivorans* A9 protein homolog from residues 39

to 61, suggesting that the longest part of this protein is extracellular (residues 62 to 247). Regarding the synteny in *Microbacterium oleivorans* A9 genome, a signal transduction histidine kinase is adjacent to this protein, indicating a possible involvement in signaling function. One of the four proteins for which the abundance decreased at 4h of uranium exposure is the glutaredoxin which can interact physically with a mercury/uranyl reductase (MerA) and plays a role in toxic metals resistance [47]. After 24h exposure, 12 transporters involved in the transport of diverse compounds (lipoprotein, Fe^{3+} -hydroxamate, dipeptide/oligopeptide/nickel, enterochelin, Fe^{3+} /spermidine/putrescine) are more abundant in the uranium condition compared to the control (**Table 2**). A DNA-binding response regulator of the NarL/FixJ family is also more detected (5.3-fold). The homologous regulator from *Escherichia coli* has been characterized and found to control nitrate- and nitrite-regulated gene expression. This is concordant with the lower detection (-2.1-fold) of a ferredoxin subunit of the nitrite reductase. In the present exposure condition, nitrate has been added in both control and uranium conditions and probably contributes to modify the production of these two proteins.

Phosphate metabolism

In our previous study, we have evidenced a phosphate efflux concomitant to the uranium efflux between 0.5h and 4h exposure and the intracellular formation of autunite, a mineral made of U, P and Ca, suggesting a strong connection between phosphate and uranium [12]. Among these proteins, the phosphohistidine phosphatase SixA is the most up-produced protein (3.5 fold-change at 0.5h and 4h). Its precise role is still unclear but its involvement in a signal transduction circuitry through down-regulation of the ArcB-to-ArcA phosphorelay has been shown under anaerobic conditions in *Escherichia coli* [56]. On the other hand, the increase abundance of the broad specificity phosphatase PhoE only after 4h (2.3 fold-change) is congruent with the phosphate efflux observed in our previous experiment [12]. Several components of an ABC-type glycerol-3-phosphate transport system have also a significant positive fold-change. No such component is found more abundant after 24h exposure probably because the phosphate is complexed with uranium leading to autunite

accumulation. The glycerol-3-phosphate has been demonstrated as a source of phosphate for the precipitation of uranium [45]. A polyphosphate kinase shows an up-production after 0.5 h (fold-change of 1.8). It can act as a defense mechanism by constituting a phosphate reserve that will be used latter to form uranyl-phosphate complex. As reported in **Figure 4**, the genomic context of this kinase reveals a locus with seven adjacent genes, possibly organized as an operon. These genes are all related to phosphate metabolism. The operon comprises a transcriptional regulatory protein, the phosphohistidine phosphatase SixA and four components of an ABC-type phosphate transport system. Five of this six genes encode proteins which have been found more abundant in uranium condition but with low statistical confidence (p-value above 0.05) or with significant but low fold-change. Noteworthy, this cluster of genes is conserved amongst several *Microbacterium* species from our collection (data not shown). In this work, several proteins involved in phosphate metabolism were modulated upon uranium exposure. The relationships between uranium and phosphate metabolism have also been evidenced by others [57-59].

Reminding that uranium can bind to highly oxygenated sites such as phosphate and carbonate, it is interesting to point out that a component of an ABC-type nitrate/sulfonate/bicarbonate transport system shows a significant increase in terms of abundance (1.9 fold-change) after an exposure of 4h, in the same time as uranium and phosphate efflux occur. Several cations transporters (K^+ , Mn^{2+} , Zn^{2+} , Mg^{2+} , Co^{2+}) are also induced upon uranium exposure. One can speculate that one of these transporters may be involved in uranyl efflux.

Iron metabolism

In our previous study and unlike to phosphate metabolism, we obtained no element suggesting an effect of uranium on iron metabolism [12]. Here, however, a strong impact on iron metabolism is highlighted through the large number of proteins related to iron metabolism more abundant under uranyl stress (19 out of 43 proteins classified in the “Inorganic ion metabolism and transport” category). Most of them are components of the siderophore iron uptake system, either ABC-transport type subunits or siderophore modification enzymes. Among these 19 proteins, two ABC-

398 type Fe^{3+} -hydroxamate transport systems and one ABC-type enterocholin transport system are found
399 systematically more abundant at the three time points (fold changes of 1.7, 2.1 and 2.8 respectively).
400 Hydroxamate and enterocholin are siderophores, low molecular compounds produced by bacteria to
401 scavenge iron (Fe(III)) in iron starvation conditions. The involvement of siderophores in tolerance to
402 toxic metals has already been demonstrated [48]. The ATPase component of an ABC-type
403 cobalamin/ Fe^{3+} -siderophore transport system is also found in higher proportion at 0.5h and 4h with
404 fold changes of 1.6 and 1.7 respectively. Moreover, the abundance of a NADPH-dependent ferric
405 siderophore reductase increased upon uranium exposure: fold change of a 1.8 and 1.6 at 0.5h and
406 24h, respectively. This protein is involved in the release of iron from siderophore. Involvement of
407 siderophores in the resistance to uranium has been poorly studied, but hydroxamate has been
408 shown to bind uranium and to be able to chelate more efficiently uranium with the carbonate form
409 [49]. In addition, induction of siderophore production in response to uranium stress has been shown
410 for the marine cyanobacterium *Synechococcus elongatus* BDU 130911 [50]. Taken together, these
411 data suggest that uranium could interact with siderophore in *Microbacterium oleivorans* A9. One
412 attractive hypothesis is that uranium could then enter the cells via this pathway.

413 Two components (EfeO and EfeB) of the EfeUOB transporter were also detected in higher amounts in
414 our dataset. This transporter is involved in iron uptake in *Bacillus subtilis* [51]. EfeB is a peroxidase
415 involved in oxidation of ferrous iron to ferric iron which allows its binding by the EfeO component.
416 This protein shows a regular increase at 0.5h, 4h, and 24h with fold changes of 1.7, 1.8 and 1.8,
417 respectively. Moreover, EfeB presents the ability to reduce the reactive oxygen species formed in
418 presence of ferrous iron and thus, confers protection to the cell. The prediction of both a cupredoxin-
419 like domain and an imelysin peptidase with a highly conserved HXXE motif confirm that we are
420 dealing with the EfeO component and not the EfeM for which the cupredoxin-like domain is absent
421 [52]. The EfeO component is a ferric iron binding protein and is able to transfer its substrate to the
422 permease EfeU. The variation in the abundance of EfeO during the experiment is not significant. The
423 EfeU component remained undetected despite its prediction in the genome. The EfeUOB transporter

of *Escherichia coli* has been shown to be induced under low pH condition [53]. This three-component transporter is also found in *Bacillus subtilis* and was shown to be involved in high-affinity uptake of both ferrous (Fe(II)) and ferric iron (Fe(III)) [51]. Taken together, these results suggest that uranium exposure is perceived by *Microbacterium oleivorans* A9 as an iron starvation, thus enhancing the synthesis of iron uptake systems. This result is in line with what has been shown for *Geobacter sulfurreducens* in which uranium induces the Fur operon [18]. Other biological molecules involved in iron metabolism, e.g. transferrin, have been shown to be able to link uranium and can constitute a way for uranium to be transported in the cells [55].

4. Concluding remarks

The high-throughput proteogenomic methodology applied to *Microbacterium oleivorans* A9 allows to quickly identify a large number of proteins and highlights those that could be involved in uranium tolerance. The proteome coverage obtained here is higher than previous proteomic studies devoted at exploring the proteome under uranium stress [18, 19]. *Microbacterium oleivorans* A9 modifies drastically its proteome upon uranium exposure. As previously hypothesized, proteins involved in phosphate metabolism are found in larger abundances under uranium stress due to higher synthesis or poorer degradation or both of these mechanisms. More surprisingly, proteins related to iron metabolism were detected with significant changes in abundance. The detailed relationships between uranium response and iron metabolism are unknown at the moment. However, based on our current results it can be hypothesized that uranium may enter the cells via siderophore transportation. This will be further investigated. The detection of 15% of functionally unassigned proteins highlights the lack of knowledge regarding proteins with potentially key roles in response to uranium stress. Focusing on the most interesting protein candidates could be done by a comparative analysis using high-throughput proteogenomics between *Microbacterium oleivorans* A9 and other *Microbacterium* isolates differing in uranium sensitivity such as *Microbacterium lemovicum* ViU22 [60] which is uranium-sensitive.

449

450 Acknowledgments

451 Nicolas Gallois was financially supported by the Commissariat à l'Energie Atomique et aux Energies
452 Alternatives (CEA) by means of the program "Thèse Amont-Aval" directed by Yves Bréchet, Haut-
453 Commissaire à l'Energie Atomique. The project was supported by CEA through the Toxicology
454 program (BEnUr). We thank Microb&co for all the writing advices given during the 7th ICME .

455

456 References

457

458

- 459 [1] Y. Abe, Y. Iizawa, Y. Terada, K. Adachi, Y. Igarashi, I. Nakai, Detection of uranium and chemical
460 state analysis of individual radioactive microparticles emitted from the Fukushima nuclear accident
461 using multiple synchrotron radiation X-ray analyses, *Anal Chem* 86(17) (2014) 8521-8525.
462 [2] D. Bugai, V. Kashparov, L. Dewiére, Y. Khomutinin, S. Levchuk, V. Yoschenko, Characterization of
463 subsurface geometry and radioactivity distribution in the trench containing Chernobyl clean-up
464 wastes, *Environ Geol* 47(6) (2005) 869-881.
465 [3] C.R. Cothorn, W.L. Lappenbusch, J.A. Cotruvo, Health effects guidance for uranium in drinking
466 water, *Health physics* 44 Suppl 1 (1983) 377-84.
467 [4] R.G. Pearson, Hard and Soft Acids and Bases, *J Am Chem Soc* 85(22) (1963) 3533-&.
468 [5] R.J. Reeder, M. Nugent, C.D. Tait, D.E. Morris, S.M. Heald, K.M. Beck, W.P. Hess, A. Lanzirotti,
469 Coprecipitation of uranium(VI) with calcite: XAFS, micro-XAS, and luminescence characterization,
470 *Geochim Cosmochim Acta* 65(20) (2001) 3491-3503.
471 [6] N. Renninger, R. Knopp, H. Nitsche, D.S. Clark, J.D. Keasling, Uranyl precipitation by *Pseudomonas*
472 *aeruginosa* via controlled polyphosphate metabolism, *Appl Environ Microbiol* 70(12) (2004) 7404-12.
473 [7] P. Yong, L.E. Macaskie, Role of citrate as a complexing ligand which permits enzymically-mediated
474 uranyl ion bioaccumulation, *Bull Environ Contam Toxicol* 54(6) (1995) 892-9.
475 [8] M.L. Merroun, S. Selenska-Pobell, Bacterial interactions with uranium: an environmental
476 perspective, *J Contam Hydrol* 102(3-4) (2008) 285-295.
477 [9] A. Nakajima, T. Horikoshi, T. Sakaguchi, Studies on the Accumulation of Heavy-Metal Elements in
478 Biological-Systems .5. Ion Effects on the Uptake of Uranium by *Chlorella-Regularis*, *Agr Biol Chem*
479 *Tokyo* 43(3) (1979) 625-629.
480 [10] M. Bouby, I. Billard, M. J., I. Rossini, Complexation of uranium VI with the siderophore
481 pyoverdine, *Radiochim Acta* 80 (1998) 95-100.
482 [11] V. Chapon, L. Piette, M.-H. Vesvres, F. Coppin, C.L. Marrec, R. Christen, N. Theodorakopoulos, L.
483 Février, S. Levchuk, A. Martin-Garin, C. Berthomieu, C. Sergeant, Microbial diversity in contaminated
484 soils along the T22 trench of the Chernobyl experimental platform, *Appl Geochem* 27(7) (2012) 1375-
485 1383.
486 [12] N. Theodorakopoulos, V. Chapon, F. Coppin, M. Floriani, T. Vercouter, C. Sergeant, V. Camilleri,
487 C. Berthomieu, L. Fevrier, Use of combined microscopic and spectroscopic techniques to reveal

interactions between uranium and *Microbacterium* sp. A9, a strain isolated from the Chernobyl exclusion zone, J Hazard Mater 285 (2015) 285-293.

[13] D.H. Nies, Microbial heavy-metal resistance, Appl Microbiol Biotechnol 51(6) (1999) 730-50.

[14] P. Junier, E.D. Vecchia, R. Bernier-Latmani, The Response of *Desulfotomaculum reducens* MI-1 to U(VI) Exposure: A Transcriptomic Study, Geomicrobiol J 28(5-6) (2011) 483-496.

[15] A. Khemiri, M. Carriere, N. Bremond, M.A. Ben Mlouka, L. Coquet, I. Llorens, V. Chapon, T. Jouenne, P. Cosette, C. Berthomieu, *Escherichia coli* response to uranyl exposure at low pH and associated protein regulations, PloS one 9(2) (2014) e89863.

[16] L. Dekker, F. Arsene-Ploetze, J.M. Santini, Comparative proteomics of *Acidithiobacillus ferrooxidans* grown in the presence and absence of uranium, Res Microbiol 167(3) (2016) 234-239.

[17] B. Panda, B. Basu, C. Acharya, H. Rajaram, S.K. Apte, Proteomic analysis reveals contrasting stress response to uranium in two nitrogen-fixing *Anabaena* strains, differentially tolerant to uranium, Aquatic toxicology 182 (2017) 205-213.

[18] R. Orellana, K.K. Hixson, S. Murphy, T. Mester, M.L. Sharma, M.S. Lipton, D.R. Lovley, Proteome of *Geobacter sulfurreducens* in the presence of U(VI), Microbiology 160(Pt 12) (2014) 2607-2617.

[19] M.C. Yung, J. Ma, M.R. Salemi, B.S. Phinney, G.R. Bowman, Y. Jiao, Shotgun proteomic analysis unveils survival and detoxification strategies by *Caulobacter crescentus* during exposure to uranium, chromium, and cadmium, J Proteome Res 13(4) (2014) 1833-1847.

[20] J. Armengaud, Next-generation proteomics faces new challenges in environmental biotechnology, Curr Opin Biotechnol 38 (2016) 174-182.

[21] A. de Groot, R. Dulerio, P. Ortet, L. Blanchard, P. Guerin, B. Fernandez, B. Vacherie, C. Dossat, E. Jolivet, P. Siguier, M. Chandler, M. Barakat, A. Dedieu, V. Barbe, T. Heulin, S. Sommer, W. Achouak, J. Armengaud, Alliance of proteomics and genomics to unravel the specificities of Sahara bacterium *Deinococcus deserti*, PLoS genetics 5(3) (2009) e1000434.

[22] C. Rubiano-Labrador, C. Bland, G. Miotello, P. Guerin, O. Pible, S. Baena, J. Armengaud, Proteogenomic insights into salt tolerance by a halotolerant alpha-proteobacterium isolated from an Andean saline spring, J Proteomics 97 (2014) 36-47.

[23] X. Zhu, S. Xie, J. Armengaud, W. Xie, Z. Guo, S. Kang, Q. Wu, S. Wang, J. Xia, R. He, Y. Zhang, Tissue-specific proteogenomic analysis of *Plutella xylostella* larval midgut using a multialgorithm pipeline, Molecular & cellular proteomics 15(6) (2016) 1791-807.

[24] M. Baudet, P. Ortet, J.C. Gaillard, B. Fernandez, P. Guerin, C. Enjalbal, G. Subra, A. de Groot, M. Barakat, A. Dedieu, J. Armengaud, Proteomics-based Refinement of *Deinococcus deserti* Genome Annotation Reveals an Unwanted Use of Non-canonical Translation Initiation Codons, Molecular & Cellular Proteomics 9(2) (2010) 415-426.

[25] C. Rubiano-Labrador, C. Bland, G. Miotello, J. Armengaud, S. Baena, Salt Stress Induced Changes in the Exoproteome of the Halotolerant Bacterium *Tistlia consotensis* Deciphered by Proteogenomics, PloS one 10(8) (2015) e0135065.

[26] J.T. Simpson, K. Wong, S.D. Jackman, J.E. Schein, S.J. Jones, I. Birol, ABySS: a parallel assembler for short read sequence data, Genome Res 19(6) (2009) 1117-23.

[27] J. Armengaud, C. Bland, J. Christie-Oleza, G. Miotello, Microbial proteogenomics, gaining ground with the avalanche of genome sequences, J Bacteriol Parasitol S3-001 (2011).

[28] R.L. Tatusov, M.Y. Galperin, D.A. Natale, E.V. Koonin, The COG database: a tool for genome-scale analysis of protein functions and evolution, Nucleic Acids Res 28(1) (2000) 33-6.

[29] C. Claudel-Renard, C. Chevalet, T. Faraut, D. Kahn, Enzyme-specific profiles for genome annotation: PRIAM, Nucleic Acids Res 31(22) (2003) 6633-9.

[30] M. Kanehisa, Y. Sato, M. Kawashima, M. Furumichi, M. Tanabe, KEGG as a reference resource for gene and protein annotation, Nucleic Acids Res 44(D1) (2016) D457-62.

[31] E.M. Hartmann, F. Allain, J.C. Gaillard, O. Pible, J. Armengaud, Taking the shortcut for high-throughput shotgun proteomic analysis of bacteria, Methods Mol Biol 1197 (2014) 275-285.

[32] G. Clair, S. Roussi, J. Armengaud, C. Duport, Expanding the known repertoire of virulence factors produced by *Bacillus cereus* through early secretome profiling in three redox conditions, Molecular & cellular proteomics 9(7) (2010) 1486-98.

- [33] G. Klein, C. Mathe, M. Biola-Clier, S. Devineau, E. Drouineau, E. Hatem, L. Marichal, B. Alonso, J.C. Gaillard, G. Lagniel, J. Armengaud, M. Carriere, S. Chedin, Y. Boulard, S. Pin, J.P. Renault, J.C. Aude, J. Labarre, RNA-binding proteins are a major target of silica nanoparticles in cell extracts, *Nanotoxicology* 10(10) (2016) 1555-1564.
- [34] J.A. Christie-Oleza, B. Fernandez, B. Nogales, R. Bosch, J. Armengaud, Proteomic insights into the lifestyle of an environmentally relevant marine bacterium, *ISME J* 6(1) (2012) 124-35.
- [35] V. Dupierriis, C. Masselon, M. Court, S. Kieffer-Jaquinod, C. Bruley, A toolbox for validation of mass spectrometry peptides identification and generation of database: IRMa, *Bioinformatics* 25(15) (2009) 1980-1.
- [36] R.G. Cote, J. Griss, J.A. Dienes, R. Wang, J.C. Wright, H.W. van den Toorn, B. van Breukelen, A.J. Heck, N. Hulstaert, L. Martens, F. Reisinger, A. Csordas, D. Ovelleiro, Y. Perez-Rivevol, H. Barsnes, H. Hermjakob, J.A. Vizcaino, The PRoteomics IDentification (PRIDE) Converter 2 framework: an improved suite of tools to facilitate data submission to the PRIDE database and the ProteomeXchange consortium, *Molecular & cellular proteomics* 11(12) (2012) 1682-9.
- [37] P.C. Carvalho, J. Hewel, V.C. Barbosa, J.R. Yates Iii, Identifying differences in protein expression levels by spectral counting and feature selection, *Genet Mol Res* 7(2) (2008) 342-356.
- [38] P. Ortet, N. Gallois, J. Long, M. Barakat, V. Chapon, Draft genome sequence of *Microbacterium oleivorans* strain A9, a bacterium isolated from Chernobyl radionuclide-contaminated soil, *Genome announcement* 999(5) (2017) e00092-17.
- [39] C. Pal, J. Bengtsson-Palme, C. Rensing, E. Kristiansson, D.G. Larsson, BacMet: antibacterial biocide and metal resistance genes database, *Nucleic Acids Res* 42(Database issue) (2014) 737-743.
- [40] H.Y. Gan, H.M. Gan, M.A. Savka, A.J. Triassi, M.S. Wheatley, L.B. Smart, E.S. Fabio, A.O. Hudson, Whole-genome sequences of 13 endophytic bacteria isolated from shrub willow (*salix*) grown in geneva, new york, *Genome announcement* 2(3) (2014) e00288-14.
- [41] P. Jagtap, J. Goslinga, J.A. Kooren, T. McGowan, M.S. Wroblewski, S.L. Seymour, T.J. Griffin, A two-step database search method improves sensitivity in peptide sequence matches for metaproteomics and proteogenomics studies, *Proteomics* 13(8) (2013) 1352-7.
- [42] J.S. Weissman, H.S. Rye, W.A. Fenton, J.M. Beechem, A.L. Horwich, Characterization of the active intermediate of a GroEL-GroES-mediated protein folding reaction, *Cell* 84(3) (1996) 481-90.
- [43] G. Fibriansah, A.T. Kovacs, T.J. Pool, M. Boonstra, O.P. Kuipers, A.M. Thunnissen, Crystal structures of two transcriptional regulators from *Bacillus cereus* define the conserved structural features of a PadR subfamily, *PloS one* 7(11) (2012) e48015.
- [44] S. Silver, T. Phung le, A bacterial view of the periodic table: genes and proteins for toxic inorganic ions, *J Ind Microbiol Biotechnol* 32(11-12) (2005) 587-605.
- [45] R.J. Martinez, M.J. Beazley, M. Taillefert, A.K. Arakaki, J. Skolnick, P.A. Sobecky, Aerobic uranium (VI) bioprecipitation by metal-resistant bacteria isolated from radionuclide- and metal-contaminated subsurface soils, *Environ Microbiol* 9(12) (2007) 3122-33.
- [46] A. Krogh, B. Larsson, G. von Heijne, E.L. Sonnhammer, Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes, *Journal of molecular biology* 305(3) (2001) 567-80.
- [47] C. Cassier-Chauvat, F. Chauvat, Responses to oxidative and heavy metal stresses in cyanobacteria: recent advances, *Int J Mol Sci* 16(1) (2015) 871-886.
- [48] M. Rajkumar, N. Ae, M.N.V. Prasad, H. Freitas, Potential of siderophore-producing bacteria for improving heavy metal phytoextraction, *Trends Biotechnol* 28(3) (2010) 142-149.
- [49] K.F. Mo, Z. Dai, D.S. Wunschel, Production and characterization of desmaloichrome relative binding affinity for uranyl ions in relation to other siderophores, *J Nat Prod* (2016) 1492-9.
- [50] V. Rashmi, M. Shylajanaciyar, R. Rajalakshmi, S.F. D'Souza, D. Prabakaran, L. Uma, Siderophore mediated uranium sequestration by marine cyanobacterium *Synechococcus elongatus* BDU 130911, *Bioresour Technol* 130 (2013) 204-210.
- [51] M. Miethke, C.G. Monteferrante, M.A. Marahiel, J.M. van Dijl, The *Bacillus subtilis* EfeUOB transporter is essential for high-affinity acquisition of ferrous and ferric iron, *Biochim Biophys Acta* 1833(10) (2013) 2267-2278.

- [52] M.B. Rajasekaran, S.A. Mitchell, T.M. Gibson, R. Hussain, G. Siligardi, S.C. Andrews, K.A. Watson, Isolation and characterisation of EfeM, a periplasmic component of the putative EfeUOBM iron transporter of *Pseudomonas syringae* pv. *syringae*, *Biochem Bioph Res Co* 398(3) (2010) 366-371.
- [53] J. Cao, M.R. Woodhall, J. Alvarez, M.L. Cartron, S.C. Andrews, EfeUOB (YcdNOB) is a tripartite, acid-induced and CpxAR-regulated, low-pH Fe²⁺ transporter that is cryptic in *Escherichia coli* K-12 but functional in *E. coli* O157:H7, *Mol Microbiol* 65(4) (2007) 857-75.
- [54] B.D. Stewart, R.T. Amos, S. Fendorf, Effect of uranium(VI) speciation on simultaneous microbial reduction of uranium(VI) and iron(III), *J Environ Qual* 40(1) (2011) 90-97.
- [55] M. Hemadi, N.T. Ha-Duong, J.M. El Hage Chahine, Can uranium be transported by the iron-acquisition pathway? Ur uptake by transferrin, *J Phys Chem B* 115(14) (2011) 4206-4215.
- [56] M. Matsubara, T. Mizuno, The SixA phospho-histidine phosphatase modulates the ArcB phosphorelay signal transduction in *Escherichia coli*, *FEBS letters* 470(2) (2000) 118-24.
- [57] L.E. Macaskie, G. Basnakova, Microbially-enhanced chemisorption of heavy metals: a method for the bioremediation of solutions containing long-lived isotopes of neptunium and plutonium, *Environ Sci Technol* 32(1) (1998) 184-7.
- [58] M.J. Beazley, R.J. Martinez, P.A. Sobecky, S.M. Webb, M. Tallefert, Uranium biomineralization as a result of bacterial phosphatase activity: Insights from bacterial isolates from a contaminated subsurface, *Environ Sci Technol* 41(16) (2007) 5701-5707.
- [59] S. Kulkarni, C.S. Misra, A. Gupta, A. Ballal, S.K. Apte, Interaction of Uranium with Bacterial Cell Surfaces: Inferences from Phosphatase-Mediated Uranium Precipitation, *Appl Environ Microbiol* 82(16) (2016) 4965-74.
- [60] L. Mondani, L. Piette, R. Christen, D. Bachar, C. Berthomieu, V. Chapon, *Microbacterium lemovicicum* sp. nov., a bacterium isolated from a natural uranium-rich soil, *Int J Syst Evol Microbiol* 63(Pt 7) (2013) 2600-6.

618 **Table 1. Functional classification of the proteins detected by at least two peptides.**

COG category	Relative abundance (%)	Number of proteins
Hypothetical protein	14,6	223
Amino acid transport and metabolism	11,0	168
Translation, ribosomal structure and biogenesis	9,3	142
Carbohydrate transport and metabolism	8,8	135
Transcription	6,2	95
Energy production and conversion	5,5	84
Coenzyme transport and metabolism	5,3	82
General function	5,1	78
Lipid transport and metabolism	4,4	67
Nucleotide transport and metabolism	4,3	66
Posttranslational modification, protein turnover, chaperones	4,0	61
Cell wall/membrane/envelope biogenesis	3,9	60
Replication, recombination and repair	3,9	60
Signal transduction mechanisms	3,6	55
Function unknown	2,8	43
Inorganic ion transport and metabolism	2,8	43
Defense mechanism	1,3	20
Cell cycle control, cell division, chromosome partitioning	1,1	17
Secondary metabolites metabolism	1,0	16
Cell motility	0,5	8
Intracellular trafficking, secretion, and vesicular transport	0,5	8
RNA processing and modification	0,1	1

619

Table 2 - Key proteins with major changes after uranium exposure

ORFeome ID ^a	Functional annotation	Functional classification	Fold change ^b	p value	Best hit ^c
After 0.5h					
Contig10_25943_-2	Chorismate mutase	Amino acid metabolism	5.75	0.16E-03	CHMU_METJA
Contig1_343955_-2	Alpha-glucosidase, glycosyl hydrolase family GH31	Carbohydrate metabolism	4.50	0.02E-03	YCR2_ESCVU
Contig6_5891_-2	Acetyl-CoA acetyltransferase	Lipid metabolism	4.00	7.25E-06	BKTB_CUPNH
Contig11_20212_1	Beta-galactosidase/beta-glucuronidase	Carbohydrate metabolism	3.85	0.02E-03	BGAL_ARTSB
Contig3_31861_1	Sugar kinase of the NBD/HSP70 family	Carbohydrate metabolism	3.80	0.02E-03	XYLR_BACPZ
Contig3_278050_-1	DNA-binding transcriptional regulator, PadR family	Transcription	3.75	0.20E-03	YWZG_BACSU
Contig7_116064_-3	Dipeptidyl aminopeptidase/acylaminoacyl peptidase	Amino acid metabolism	3.75	0.15E-03	DAPB3_PSEMX
Contig8_12690_-3	tRNA A37 threonylcarbamoyladenosine modification protein TsaB	Translation, ribosome	3.67	0.09E-03	Y3455_MYCBO
Contig2_261531_-3	Phosphohistidine phosphatase SixA	Signal transduction	3.50	0.10E-03	Y1276_MYCTO
Contig1_23131_1	Serine protease, subtilisin family	PTM, protein turnover	3.50	0.05E-03	HLV_NATA1
Contig8_9448_1	Hypothetical protein	Hypothetical	3.50	0.03E-03	Y3037_MYCTO
Contig1_334362_3	ABC-type glycerol-3-phosphate transport system	Carbohydrate metabolism	3.50	2.97E-03	YURO_BACSU
After 4h					
Contig11_87308_2	Uncharacterized membrane protein Ykol	Function unknown	10.75	3.99E-06	No results
Contig1_430924_1	D-arabinose 1-dehydrogenase, Zn-dependent alcohol dehydrogenase family	Carbohydrate metabolism	4.60	9.21E-03	ADH_SULTO
Contig3_147088_1	DNA-binding transcriptional regulator/sugar metabolism, DeoR/GlpR family	Transcription	4.58	1.72E-04	AGAR_ECO57
Contig5_94994_2	Hypothetical protein	Hypothetical	3.83	1.64E-03	No results
Contig11_20081_2	6-phosphogluconolactonase/Glucosamine-6-P isomerase/deaminase	Carbohydrate metabolism	3.80	2.68E-02	NAGB_STRCO
Contig5_120376_1	Hypothetical protein	Hypothetical	3.75	1.57E-02	No results
Contig5_105896_-2	Adenosylmethionine-8-amino-7-oxononanoate aminotransferase	Coenzyme metabolism	3.55	2.92E-03	Y3329_MYCTU
Contig3_278050_-1	DNA-binding transcriptional regulator, PadR family	Transcription	3.50	5.59E-03	YWZG_BACSU
Contig2_261531_-3	Phosphohistidine phosphatase SixA	Signal transduction	3.50	2.28E-02	Y1276_MYCTO
After 24h					
Contig11_87308_2	Uncharacterized membrane protein Ykol	Function unknown	25	1.15E-04	No results
Contig5_20481_3	ABC-type lipoprotein export system	Cell wall/membrane	8.18	7.26E-07	Y1508_METJA
Contig5_94994_2	Hypothetical protein	Hypothetical	5.25	6.46E-03	No results
Contig5_161786_-2	DNA-binding response regulator, NarL/FixJ family	Signal transduction	5.25	4.78E-04	DEGU_BREBE
Contig1_136842_3	Periplasmic serine protease, S1-C subfamily	PTM, protein turnover	4.33	8.18E-04	DEGP_SALTY
Contig3_196395_-3	Hypothetical protein	Hypothetical	4.28	2.74E-03	TITIN_MOUSE
Contig15_39811_-1	DNA-binding transcriptional regulator, AcrR family	Transcription	3.75	4.04E-03	MIFTR_MYCTO

^a ORFeome ID refers to Supplementary Data Table S2.^b Fold changes represent the expression with uranium exposure normalized to expression without uranium.^c The best hit correspond to the first hit obtained by BLASTp against Swissprot database.

Figure 1 – Shotgun nanoLC-MS/MS strategy used for the proteome analysis of *Microbacterium*

***oleivorans* A9 under uranium exposure or not.** A total of 24 samples were processed: four biological replicates for each time point at 0,5h, 4h and 24h of exposure. After lysis, the proteins were subjected to a short SDS-PAGE migration before trypsin proteolysis in-gel. A Q-Exactive HF mass spectrometer was used to identify the resulting peptides.

Figure 2 – Venn diagram showing the number of proteins with significant abundance changes in the three sampling time points.

Figure 3 – Example of the phenylalanine, tyrosine and tryptophan biosynthesis pathways. The boxes represent the branched metabolic pathways. The enzymes are identified through their EC number. Proteins detected by proteogenomics are highlighted in red, proteins more abundant upon uranium exposure are indicated in blue and proteins predicted in the genome but not detected by proteomics are in grey. The genes' ID corresponding to the enzymes are as follow: 1.3.1.12:

MicroBact_A9DB#Contig4_204950_-2, 2.4.2.18: MicroBact_A9DB#Contig4_125788_-1, 2.5.1.19:

MicroBact_A9DB#Contig2_13726_-1, 2.5.1.54: MicroBact_A9DB#Contig4_169845_-3, 2.6.1.1:

MicroBact_A9DB#Contig12_70604_2, 2.6.1.9: MicroBact_A9DB#Contig7_122729_2, 2.7.1.71:

MicroBact_A9DB#Contig1_401394_-3, 4.1.1.48: MicroBact_A9DB#Contig1_371158_-1, 4.2.1.10:

MicroBact_A9DB#Contig1_399335_-2, 4.2.1.20: MicroBact_A9DB#Contig1_369105_-3, 4.2.1.51:

MicroBact_A9DB#Contig5_42426_3, 4.2.1.91: MicroBact_A9DB#Contig3_67122_3, 4.2.3.5:

MicroBact_A9DB#Contig1_401926_-1, 5.4.99.5: MicroBact_A9DB#Contig10_25943_-2, 6.1.1.1:

MicroBact_A9DB#Contig7_15747_3, 6.1.1.20: MicroBact_A9DB#Contig7_133127_2.

Figure 4 – Genomic context of the polyphosphate kinase. Fold-change (FC) are indicated for the first time point of sampling (0.5h). Proteins for which the gene name is in bold were detected by proteomics.

646

647 **List of supplementary material**

648 **Supplementary Table S1.** List of proteins and their spectral count in A9 samples in the first
649 proteogenomic search.

650 **Supplementary Table S2.** List of proteins and their spectral count in A9 samples in the
651 second round search. NSAF: Normalized Spectral Abundance Factor; SC: Spectral Count

652

Figure

[Click here to download Figure: Proteogenomics of Microbacterium sp. A9_Four figures revised version.pptx](#)

Microbacterium oleivorans A9 chromosome