

Development of operando solid-state Nuclear Magnetic Resonance Spectroscopy (ssNMR) for metal-oxygen batteries

Magali Gauthier, Alan Wong

► **To cite this version:**

Magali Gauthier, Alan Wong. Development of operando solid-state Nuclear Magnetic Resonance Spectroscopy (ssNMR) for metal-oxygen batteries. IUPAC 2019, Jul 2019, Paris, France. cea-02327967

HAL Id: cea-02327967

<https://hal-cea.archives-ouvertes.fr/cea-02327967>

Submitted on 23 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Development of operando solid-state Nuclear Magnetic Resonance Spectroscopy (ssNMR) for metal-oxygen batteries

Magali Gauthier, Alan Wong

NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif sur Yvette Cedex, France

Rechargeable metal-O₂ batteries have attracted much attention in recent years as a possible alternative to the widely used lithium-ion batteries. This is particularly the case for lithium and sodium-oxygen batteries, due to their potential high energy density.¹ However, great challenges remain in the development of metal-O₂ batteries, including the stabilization of the discharge products, poor cyclability and the need for new cathode design. On a more fundamental aspect, one must completely grasp the understanding of the underlying electrochemical mechanisms taking place inside the M-O₂ battery. Clear identifications of the discharge electrochemical pathways and their products (M-O₂, M₂O₂,...), as well as the reactivity of the electrolyte, are crucial. Discharge products have already been observed using many *ex situ* techniques such as XRD or Raman, however it may not provide a realistic view of the reactions and may lead to biased interpretation. Conversely, *In situ/operando* techniques² can capture the dynamic changes unequivocally in the cell environment during cycling and provide unprecedented information on intermediate phases that may be undetectable through *ex situ* analyses.

In the past few years, *ex situ* ssNMR emerged as a valuable tool for characterizing metal-O₂ cells on *ex situ* discharged and charged samples (Fig 4).^{3,4} *Operando* ssNMR appears thus of considerable interest to monitor the structural and electronic changes in M-O₂ upon cycling and to obtain valuable insights on the transition of chemical phases. To this aim, we recently designed an *operando* cell dedicated to metal-O₂ batteries that enables for O₂ gas to react at the cathode without compromising the NMR experiments. The electrochemical performance in the Li-O₂ configuration is quite similar to what is expected in the literature and validates the design of the cell. We will detail the working principle of the cell, its performance, and present the evolution of NMR signals during cycling.



Operando ⁷Li NMR on a Li-O₂ cell

References

1. Landa-Medrano et al.. *J. Phys. Chem. Lett.* **7**, 1161–1166 (2016).
2. Liang et al.. *Small Methods* **1**, 1700150 (2017).
3. Leskes et al. *J. Phys. Chem. C* **117**, 26929–26939 (2013).
4. Reeve et al. *J. Am. Chem. Soc.* **139**, 595–598 (2017).