

## One-step formation of stable multiple emulsions

Patrick Guenoun, Marine Protat, Noemie Bodin, Nadège Pantoustier, Jean Daillant, Frédéric Gobeaux, Florent Malloggi, Patrick Perrin

► **To cite this version:**

Patrick Guenoun, Marine Protat, Noemie Bodin, Nadège Pantoustier, Jean Daillant, et al.. One-step formation of stable multiple emulsions. 253rd ACS meeting. Interfacial Phenomena and the Oil-Water Interface, Apr 2017, San Francisco, United States. cea-02328442

**HAL Id: cea-02328442**

**<https://hal-cea.archives-ouvertes.fr/cea-02328442>**

Submitted on 23 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**ABSTRACT SYMPOSIUM NAME:** Interfacial Phenomena & the Oil-Water Interface

**ABSTRACT SYMPOSIUM PROGRAM AREA NAME:** COLL

**CONTROL ID:** 2606961

**PRESENTATION TYPE:** Oral Only : Consider for Sci-Mix

**TITLE:** One-step formation of stable multiple emulsions

**AUTHORS (FIRST NAME, LAST NAME):** Patrick M. Guenoun<sup>1</sup>, Marine Protat<sup>1</sup>, Noémi Bodin<sup>1</sup>, Nadège Pantoustier<sup>2</sup>, Jean Daillant<sup>3</sup>, Frédéric Gobeaux<sup>1</sup>, Florent Malloggi<sup>1</sup>, Patrick Perrin<sup>2</sup>

**INSTITUTIONS (ALL):**

1. CEA Saclay, Université Paris Saclay, Gif sur Yvette, France.

2. ESPCI, Paris, France.

3. SOLEIL, Saint Aubin, France.

**ABSTRACT BODY:**

**Abstract:** Stimuli-responsive multiple emulsions formed in a one-step mechanical emulsification process are shown to be stable for months. These emulsions are stabilized by amphiphilic copolymers synthesized by atom transfer radical polymerization (ATRP). Depending on pH, ionic strength and temperature, different emulsion types are obtained, including water-continuous (W/O/W) and oil-continuous (O/W/O) multiple emulsions.

In particular W/O/W emulsions can be formed with biocompatible molecules like poly(dimethylsiloxane)-b-poly(dimethylaminoethyl methacrylate) copolymer and Miglyol® 812 as an oil phase. These emulsions enable the encapsulation and controlled release of hydrophilic species with the three stimuli: pH, ionic strength and temperature. Furthermore, we studied the formation conditions of multiple water – toluene – polystyrene-b-poly(styrene-st-dimethylaminoethyl methacrylate) emulsions. The conformation of the polymer adsorbed at the water – oil interface was probed using an original neutron reflectivity set-up. Our results evidence a correlation between emulsion type and polymer conformation. More precisely, the formation of multiple emulsions is promoted by a decrease in the curvature of the microemulsions formed in water, in agreement with cryo transmission electron microscopy and small angle neutron scattering. Finally, we exhibit predictive criteria for the formation of multiple emulsions based on interfacial tension and polymer partitioning measurements.