



**HAL**  
open science

## Reaction mechanisms in swelling clays under ionizing radiation: impact on the geological storage of the nuclear waste

Maxime Lainé, Thierry Allard, Etienne Balan, Jean-Louis Robert, Sophie Le Caër

### ► To cite this version:

Maxime Lainé, Thierry Allard, Etienne Balan, Jean-Louis Robert, Sophie Le Caër. Reaction mechanisms in swelling clays under ionizing radiation: impact on the geological storage of the nuclear waste. 16th International Clay Conference, Jul 2017, Granada, Spain. cea-02338975

**HAL Id: cea-02338975**

**<https://cea.hal.science/cea-02338975>**

Submitted on 30 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# REACTION MECHANISMS IN SWELLING CLAYS UNDER IONIZING RADIATION: IMPACT ON THE GEOLOGICAL STORAGE OF THE NUCLEAR WASTE

LAINÉ MAXIME (1), ALLARD THIERRY (2), BALAN ETIENNE (2), ROBERT JEAN-LOUIS (2), LE CAËR SOPHIE (1)\*

(1) CEA/Saclay, DSM/IRAMIS/NIMBE/LIONS UMR-3685, F-91191 Gif-sur-Yvette cedex, France, (2) IMPMC, Sorbonne Universities, UPMC, UMR-7590, 4 Place Jussieu, F-75252 Paris cedex 05, France

The influence of ionizing radiation on clay minerals is poorly known, in spite of their use as a major component of the engineered barrier in High Level Nuclear Waste Repositories (HLNWR). In this context, the production of H<sub>2</sub> by clay minerals under ionizing radiation could be a real issue. It can, e.g., lead to the loss of radionuclide retention properties by creating cracks in the engineered barrier. It is thus important to determine H<sub>2</sub> formation reaction mechanisms and to understand the role of several parameters on this production, such as the water amount, the presence of impurities and the nature of the clay mineral.

We studied synthetic clay minerals (talc, which has no water molecules, montmorillonite and saponite which exhibit charge excess in different locations) [1-2] and also natural montmorillonites from Camp Berteau [3]. The reactivity under electron irradiation is deciphered thanks to H<sub>2</sub> production measurements at well-defined relative humidities together with electron spin resonance experiments that allow detecting the defects created upon irradiation. Comparing the reactivity induced in the synthetic samples enables understanding the impact of the presence of two-dimensionally confined water molecules and the influence of the charge location in the sheets of the clay mineral. Moreover, the comparison between synthetic and natural talc evidences the strong role played by impurities on the H<sub>2</sub> production yields [1]. We find that H<sub>2</sub> production is due to the presence of structural -OH groups and that the confinement of water molecules is a crucial factor. Moreover, the charge location in the sheets of the clay minerals plays only a marginal role [1-2]. All these results will be discussed and the corresponding reaction mechanisms will be presented. Lastly, the role of impurities in natural clay minerals such as Fe<sup>3+</sup> that strongly decrease the H<sub>2</sub> production will be discussed.

## References:

- [1] Lainé M., Allard T., Balan E., Martin F., Von Bardeleben H.J., Robert J.-L., Le Caër S. (2016). Reaction mechanisms in talc under ionizing radiation: evidence of a high stability of H<sup>•</sup> atoms, *J. Phys. Chem. C*, pp 2087–2095.
- [2] Lainé M., Balan E., Allard T., Paineau E., Jeunesse P., Mostafavi M., Robert J.-L., Le Caër S. (2017). Reaction mechanisms in swelling clays under ionizing radiation: influence of the water amount and of the nature of the clay mineral. *RCS Adv.*, pp 526-534.
- [3] Fourdrin C., Aarrachi H., Latrille C., Esnouf S., Bergaya F., Le Caër S. (2013). Water Radiolysis in Exchanged-Montmorillonites: The H<sub>2</sub> Production Mechanisms. *Environ. Sci. Technol.*, pp 9530-9537.