

# Accelerated ageing of electrolytes in Lithium-ion batteries: the point of view of a radiation chemist

Sophie Le Caër

► **To cite this version:**

Sophie Le Caër. Accelerated ageing of electrolytes in Lithium-ion batteries: the point of view of a radiation chemist. Conférence de Physique-Chimie (CPC16), Oct 2016, Nancy, France. cea-02339028

**HAL Id: cea-02339028**

**<https://hal-cea.archives-ouvertes.fr/cea-02339028>**

Submitted on 30 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Accelerated ageing of electrolytes in Lithium-ion batteries: the point of view of a radiation chemist

Sophie Le Caër

LIONS, NIMBE UMR 3685, CEA, CNRS, Université Paris Saclay, Bât. 546, F-91191 Gif-sur-Yvette Cedex, France. email: [sophie.le-caer@cea.fr](mailto:sophie.le-caer@cea.fr)

Since the early nineties, the rechargeable Li-ion battery (LIB) technology has dominated the electronic market. These batteries have become essential components in portable electronic applications. Since then, ageing processes are a growing global concern, essentially for their applications in electric and hybrid vehicles.

Ageing phenomena occurring in diethylcarbonate (DEC), DEC/LiPF<sub>6</sub> propylene carbonate (PC) and PC/LiPF<sub>6</sub> solutions, selected as model systems, have been studied using gamma- and pulse- radiolysis as a tool to generate similar species as the ones occurring in electrolysis of Li-ion batteries and to mimic the processes (Fig. 1) [1-3]. We prove that similar results were obtained in the ageing of an electrochemical cell filled with the same model solution [2]. This highlights the interest of the radiolysis approach in the field of ageing of electrolytes. Radiolysis has indeed the following major advantages: i) ageing processes are strongly accelerated (minutes/hours as compared to weeks and months in conventional battery studies); ii) time-resolved experiments are possible, enabling to study the system on multiple temporal scales (from picoseconds to minutes and days) [2]. This allows measuring rate constants, and writing very detailed reaction mechanisms; iii) the possibility to study the reactivity of each solvent without/with the salt (of course studies without the salt has no sense in the battery field!), leading to a very accurate understanding of the behavior of the system; iv) the possibility to perform a quick screening of many electrolytes, to identify rapidly the most promising ones. All these points will be discussed.




Fig. 1. Scheme comparing primary electron transfers at the electrodes in the electrolytic charge/ageing processes of a Li-ion battery (with molar DEC/ LiPF<sub>6</sub> solution as a model electrolyte) (top), and after ionization in the bulk in the radiolytic process with the same medium (bottom).

### References

- [1] D. Ortiz, V. Steinmetz, D. Durand, S. Legand, V. Dauvois, P. Maître, S. Le Caër, *Nature Comm.*, **6**, doi:10.1038/ncomms7950 (2015).
- [2] D. Ortiz, I. Jimenez Gordon, J.P. Baltaze, O. Hernandez-Alba, S. Legand, V. Dauvois, G. Si Larbi, U. Schmidhammer, J.-L. Marignier, J.-F. Martin, J. Belloni, M. Mostafavi, S. Le Caër, *ChemSusChem*, **8**, 3605 (2015).
- [3] S. Le Caër, D. Ortiz, J.-L. Marignier, U. Schmidhammer, J. Belloni, M. Mostafavi, *J. Phys. Chem. Lett.*, **7**, 186 (2016)