



HAL
open science

Following the growth of *Saccharomyces Cerevisiae* in standard culture conditions using hyperpolarized ^{129}Xe NMR

Céline Boutin, Estelle Léonce, Patrick Berthault, Guillaume Carret

► To cite this version:

Céline Boutin, Estelle Léonce, Patrick Berthault, Guillaume Carret. Following the growth of *Saccharomyces Cerevisiae* in standard culture conditions using hyperpolarized ^{129}Xe NMR. EUROMAR 2018, Jul 2018, Nantes, France. cea-02339659

HAL Id: cea-02339659

<https://cea.hal.science/cea-02339659>

Submitted on 30 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Hyperpolarization

EUROMAR2018/222

Following the growth of *Saccharomyces Cerevisiae* in standard culture conditions using hyperpolarized ^{129}Xe NMR

Celine Boutin¹, Estelle Leonce¹, Patrick Berthault¹, Guillaume Carret¹

¹IRAMIS, CEA Saclay, Gif-sur-Yvette, France

If selected, I agree to present my work during an oral communication: Yes

Abstract: Xenon has several interesting properties for the NMR study of biological cells: 1) its nuclear polarization can be boosted via optical pumping, which increases the detection sensitivity by several orders of magnitude. 2) It is nontoxic and soluble in biologic medium and crosses the plasma membrane while keeping its polarization. 3) It has a wide range of chemical shift which makes its NMR signature sensitive to fine cell changes. In particular with cell suspensions two distinct signals are observed on the ^{129}Xe NMR spectrum, corresponding to xenon in the bulk and xenon inside the cells.[1] This has led to promising applications, such as the discrimination of cells sensitive and resistant to chemotherapy.[2]


In this work, we studied *Saccharomyces Cerevisiae* cells at different times of a culture via hyperpolarized ^{129}Xe NMR (xenon was hyperpolarized using our home made optical pumping setup) and controlled the number of cells by the measure of the optical density at 600 nm. For each time stage of the yeast culture, a sample was withdrawn and after hyperpolarized xenon addition, the corresponding ^{129}Xe spectra were analyzed in a 11.7 Tesla spectrometer: the chemical shift splitting, the in-out exchange rate and the proportion of xenon inside the cells were extracted and compared with classical analyses using optical density. The evolution of the spectral signature of ^{129}Xe in *S. Cerevisiae* is dependent of the culture time as presented in the figure. Hypotheses of explanations will be presented.

In parallel, new methodologies, compatible with narrow-bore NMR spectrometers, were developed to allow the in situ follow-up of cell cultures by NMR, using microfluidic system and micro-NMR detection. Surface treatment such as parylene coating was shown to increase biocompatibility.

[1] C. Boutin et al., Hyperpolarized ^{129}Xe NMR signature of living biological cells, *NMR Biomed.* 24 (2011) 1264.

[2] France Patent 12 52922 (2012) C. Boutin, P. Berthault, Procédé de détermination de la résistance cellulaire aux médicaments.

You can insert an image with the button on the right:



Disclosure of Interest: None Declared