



Mapping the electrocatalytic activity of modified Carbon Nanotubes by Scanning ElectroChemical Microscopy for Oxygen Reduction

O Henrotte, B. Jousselme, S. Campidelli, V. Derycke, R. Cornut

► To cite this version:

O Henrotte, B. Jousselme, S. Campidelli, V. Derycke, R. Cornut. Mapping the electrocatalytic activity of modified Carbon Nanotubes by Scanning ElectroChemical Microscopy for Oxygen Reduction. Journées d'Electrochimie 2017, Jun 2017, Bordeaux, France. cea-02341312

HAL Id: cea-02341312

<https://cea.hal.science/cea-02341312>

Submitted on 31 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mapping the electrocatalytic activity of modified Carbon Nanotubes by Scanning ElectroChemical Microscopy for Oxygen Reduction.

O. Henrotte^a, B. Jousselme^a, S. Campidelli^a, V. Derycke^a, and R. Cornut^a

^a LICSEN, NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay 91191 Gif-sur-Yvette Cedex, France

The massive adoption of Proton Exchange Membrane Fuel Cell (PEMFC) technology is critically depending on the finding of new materials to substitute noble metals for Oxygen Reduction Reaction (ORR).

In this field, performances are usually evaluated with Rotating disk electrodes (RDE). Unfortunately in this configuration the signature of the electrocatalytic activity is convoluted with different contributions such as the porosity, the electronic conductivity, the roughness, etc. In this project we propose to use Scanning ElectroChemical Microscopy (SECM)^[1] to investigate the electrocatalytic activity of modified carbon nanotubes (NTC)^[2].

Our approach consists in the study of individual and small aggregates of these nano-objects in order to acquire an image of their electrocatalytic activity. For this we acquire concordant SECM and SEM images of NTC agglomerates as illustrated in figure 1. This methodology permits to identify the origin of the achieved performances without distortion due to the averaging of the response when working at the material level. This permits to better understand the ORR mechanisms and improve the state of the art methodologies for the synthesis of noble metal free electrocatalysts.




Fig. 1: a) SECM image (done with 0.3 μm diameter gold electrode in H₂SO₄ 0.5M) of the ORR activity of NTCs b) SEM image of the same area.

[1] D. Polcari, P. Dauphin-Ducharme and J. Mauzeroll. *Chem. Rev.*, **2016**, 116, 13234–13278.

[2] P. D. Tran, A. Morozan, S. Archambault, J. Heidkamp, P. Chenevier, H. Dau, M. Fontecave, A. Martinent, B. Jousselme and V. Artero. *Chem. Sci.*, **2015**, 6, 2050–2053.