

## Charged clusters in liquids

I. Chikina, A Varlamov, V. Shikin

► **To cite this version:**

I. Chikina, A Varlamov, V. Shikin. Charged clusters in liquids. International Colloquium "Nanomatériaux: Microstructures et propriétés", Mar 2017, Tunis, Tunisia. cea-02341534

**HAL Id: cea-02341534**

**<https://hal-cea.archives-ouvertes.fr/cea-02341534>**

Submitted on 31 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

I Chikina , A Varlamov , V B Shikin

L.I.O.N.S., UMR 3685 CEA-CNRS, NIMBE, CEA Saclay, CEA-F-91191, Gif sur Yvette, France, EU

### Charged clusters in liquids

We discuss the instability of the water surface in applied vertical electric field induced by the thermal dissociation of molecules. It is shown that the breakdown phenomena occurring as the result of such instability are characterized by two relaxation times. The first one,  $\tau_{jet}$ , is the discharge time of the surface. The second,  $T^*$ , characterizes the process of charge accumulation in the layer near the liquid-vapor boundary. The latter depends on the properties of the diluted charged solution and intensity of the vertical electric field. Since  $T^* \gg \tau_{jet}$  it determines the occurring cyclic alternation of the breakdowns. The nature of both times as well as the methods for their measurement are discussed.