

A doubly responsive probe for the detection of Cys4-tagged proteins

Emilie Mari, C. Boutin, E. Léonce, B Rousseau, M. Erard, P. Berthault

► **To cite this version:**

Emilie Mari, C. Boutin, E. Léonce, B Rousseau, M. Erard, et al.. A doubly responsive probe for the detection of Cys4-tagged proteins. European Molecular Imaging Meeting, Mar 2016, Utrecht, Netherlands. cea-02346358

HAL Id: cea-02346358

<https://hal-cea.archives-ouvertes.fr/cea-02346358>

Submitted on 5 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A doubly responsive probe for the detection of Cys4-tagged proteins

E. Mari^{abc}, C. Boutin^c, E. Léonce^c, B. Rousseau^a, M. Erard^b and P. Berthault^c

^a iBiTec-S/SCBM, LabEx LERMIT, CEA Saclay, 91191 Gif-sur-Yvette, France.

^b Laboratoire de Chimie Physique, Université Paris Sud, 91405 Orsay Cedex, France

^c NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay 91191 Gif-sur-Yvette, France

Introduction

Full understanding of intracellular phenomena involves sensitive and non-invasive detection. A less disruptive method than labeling of fluorescent proteins uses binding between a tag of only six natural amino acids that can be genetically incorporated into the protein of interest and a small molecule called FIAsh[1]. This molecule has the ability to fluoresce only when it binds to its 4Cys-tag target. Another technique based on ¹²⁹Xe NMR has emerged. Xenon is hyperpolarized to enhance the NMR signal by orders of magnitude and its reversible encapsulation in functionalized host systems gives it a specific spectral signature[2]. Capability of the noble gas to cross cell membranes without losing its polarization[3] enables *in cellulo* investigations.

Here we report the first design and study of a dual fluorescence- and ¹²⁹Xe NMR-based sensor of Cys4-tagged proteins[4].

Methods

The biosensor consists of a cryptophane functionalized by a FIAsh derivative through an ethylene diamine spacer. Laser-polarized xenon is reversibly encapsulated in the cryptophane. Interaction between the biosensor and a model peptide is studied in this work.

Results

The biosensor reveals some promising properties. When bound to the target peptide, fluorescence increases by a factor 24 and the caged Xe NMR frequency is shifted by a large value (8 and 23 ppm resp. for two isomers of the biosensor). Its internalization in Cos7 cells is evidenced by confocal fluorescence.

Conclusions

This doubly smart probe is highly promising for monitoring, studying, detecting proteins *in cellulo*. Structural, chemical and lateral resolutions are combined by the bimodality of this new concept.

Acknowledgement

PhD thesis funded by Idex Paris Saclay. Supports from the French Ministry of Research (project ANR-12-BSV5-0003), from the *Fondation pour la Recherche Médicale* (project DCM20111223065) and from INSERM (convention #PC201326) are acknowledged. Dr. Brotin (ENS Lyon) synthesized the cryptophane core.

References

- [1] B. A. Griffin et al., *Science*, 281 (1998) 269-272
- [2] P. Berthault et al., *Prog. NMR Spectrosc.* 55 (2009) 35-60
- [3] C. Boutin et al., *NMR Biomed.* 2011, 24, 1264-1269
- [4] N. Kotera et al., *Chem. Comm.* 51 (2015) 11482-11484