

Electrochemical characterisation of the Ce(IV) limiting carbonate complex

C. Riglet-Martial, P. Vitorge, V. Calmon

► **To cite this version:**

C. Riglet-Martial, P. Vitorge, V. Calmon. Electrochemical characterisation of the Ce(IV) limiting carbonate complex. *Radiochimica Acta*, R Oldenbourg Verlag GMBH, 1998, 82, pp.69-76. cea-02359293

HAL Id: cea-02359293

<https://hal-cea.archives-ouvertes.fr/cea-02359293>

Submitted on 12 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Electrochemical Characterisation of the Ce(IV) limiting carbonate complex

Chantal Riglet-Martial¹, Pierre Vitorge² and Véronique Calmon¹

¹CEA-Cadarache, DCC/DESD/SESD, 13108 Saint-Paul-lez-Durance cedex, France.chantal.martial(at)cea.fr

²CEA-Saclay, DCC/DESD/SESD, 91191 Gif-sur-Yvette Cedex, France.pierre.vitorge(at)cea.fr

Keywords : cerium (III/IV), actinides (III/IV), carbonate, complex, activity coefficients, cyclic voltametry

Abstract

The stoichiometry and the thermodynamic formation constant of the limiting complex of Ce(IV) were determined at $19.3 \pm 1.0^\circ\text{C}$ by using cyclic voltametry technique at a hanging mercury drop working electrode in concentrated bicarbonate/carbonate medium. The Ce(IV/III) redox potential was measured at pH varying from 9.3 to 10.6 and $[\text{CO}_3^{2-}]$ varying from 1.0 to 1.5 M by performing a CO_3^{2-} titration with CO_2 gas. The ionic strength and junction potential effects were taken into account for the potentiometric calibrations and measurements. Quantitative interpretation of the variations of the formal potential $E_{\text{IV/III}}$ showed that no polymerisation took place during the redox reaction, and that two CO_3^{2-} ligands, but no OH^- ligand, were exchanged. As the accepted stoichiometry for the limiting complex of Ce(III) is $\text{Ce}(\text{CO}_3)_4^{5-}$, the Ce(IV) species is $\text{Ce}(\text{CO}_3)_6^{8-}$. In a 3.06 molal Na^+ carbonate/bicarbonate medium (Ionic strength = 4.33 mol.kg^{-1}), $E_{\text{IV/III}}^0 = 0.161 \pm 0.008 \text{ V/SHE}$ (in molal units) was measured. This value, combined with the published $\text{Ce}(\text{CO}_3)_4^{4-}$ formation constant and the (re-evaluated) ($\text{Ce}^{4+}/\text{Ce}^{3+}$) standard potential, is used to calculate the $\text{Ce}(\text{CO}_3)_6^{8-}$ formation constant $\log_{10}(\beta_6^{\text{IV}}) = 42.2 \pm 0.5$ (defined in molal concentration except for Ce^{4+} in activity : see table 1) in the same medium. The values of $E_{\text{IV/III}}^0$ and $\log_{10}(\beta_6^{\text{IV}})$ are ionic strength dependant, e.g. $E_{\text{IV/III}}^0 = 0.182 \pm 0.009 \text{ V/SHE}$ and $\log_{10}(\beta_6^{\text{IV}}) = 41.8 \pm 0.5$ in molar units in a 2.67 M NaClO_4 medium (Na^+ molality = molal ionic strength = 3.06 mol.kg^{-1}). The possible formation of $\text{Ce}(\text{CO}_3)_5^{6-}$ is discussed under the experimental conditions used, $\log_{10}(\beta_5^{\text{IV}}) \leq 41.8 \pm 0.5$ (in molal units).

Introduction

The actinide chemistry in reducing aqueous solutions is important for many waste disposal issues. In a recent bibliographic review [1], we pointed out contradictory interpretations for the speciation of actinide(IV) elements in bicarbonate/carbonate media, and showed that the limiting complex can be a starting point to determine aqueous speciation in environmental conditions. From redox measurements involving the known U(VI), Np(V), Pu(VI) and Am(III) limiting carbonate complexes, the stoichiometry $\text{An}(\text{CO}_3)_5^{6-}$ has been proposed for the actinide(IV) limiting carbonate complex ([2, 3] for U(IV), [4] for Np(IV), [5] for Pu(IV) and [6, 7, 8] for Am(IV)). This stoichiometry has been confirmed from the published redox data and the stability of the limiting complex has been accurately determined for uranium only. Recently, this stoichiometry has been confirmed for the Pu(IV) limiting carbonate complex from EXAFS measurements [9]. Surprisingly, among analogue elements, both $\text{Ce}(\text{CO}_3)_5^{6-}$ and $\text{Ce}(\text{CO}_3)_6^{8-}$ [10, 11, 12, 13, 14, 15] have been proposed for the Ce(IV) limiting complex. As the chemistry of Ce(III) [16] and Am(III) [7, 8] in carbonate medium is well known, the reversible (or quasi-reversible) M(IV/III) redox couple may be used to study the chemistry of Ce(IV) and Am(IV) in concentrated bicarbonate/carbonate media. Convincing evidence, based on potentiometric measurements, of the esacarbonato cerium(IV) stoichiometry was proposed in reference [15] in a 8 molal Na^+ medium and in a carbonate concentration range from 2.084 to 4.00 mol/kg. But since the authors used an Na^+ specific electrode as a reference electrode, they had to correct their data for the Na^+ activity variations to determine the number of carbonate ions exchanged during the redox reaction. In the present work, the stoichiometry of the Ce(IV) limiting carbonate complex was checked by measuring the Ce(IV/III) formal potential in 3.06 m Na^+ solutions of much lower sodium carbonate concentrations (1.00 to 1.45 mol/kg) using cyclic voltametry with a different electrochemical cell and a different methodology in order to avoid the Na^+ activity coefficient corrections. The liquid junction potentials were measured and taken into account in the treatment of the data.

Experimental

All bicarbonate/carbonate solutions, prepared from crystallised NaHCO_3 (Merck, p.a.) and concentrated NaOH (Merck, p.a., $d=1.35$), were checked by acid-base titration using H_2SO_4 (Merck Titrisol). A 0.1 M Ce(III) stock solution was prepared from crystallised $\text{Ce}(\text{ClO}_4)_3 \cdot 6\text{H}_2\text{O}$ (Johnson Matthey, purity > 99.9%). The test solutions were obtained by adding an aliquot of the Ce stock solution to the previous sodium bicarbonate/carbonate aqueous solutions. The solutions used in the reference electrodes were prepared from crystallised NaCl and NaClO_4 (both Merck, p.a.). The response of the glass electrode was checked by using commercial buffer solutions (Merck Titrisol) of pH 2, 7, 9 and 12. Demineralized water delivered by a Millipore Milli-Q plus purifier was used for all the dilutions.

The redox potential of the Ce(IV/III) couple was measured by cyclic voltametry, using a classical three electrode device - including an Ag/AgCl reference electrode (Radiometer XM630/D8), a hanging mercury drop electrode (HMDE Metrom E410) as a working electrode and a platinum wire (Radiometer Pt11/CMP) as a counter electrode - connected to an electrochemical analyser (Radiometer Voltalab 32) including a programmable interface with the Voltmaster 2 software. In order to avoid the possible damage of the reference electrode by the carbonate ions, it was isolated from the test solution by means of a capillary extension (Radiometer FDL2/CMP) filled with a 0.02 M NaCl , (I-0.02) M NaClO_4 solution, of same ionic strength I as that of the test solution. The potential of this electrode (noted REF) was checked once a day. Differences higher than 0.2 mV were never observed. The formal potential of the REF electrode was calculated from Nernst law: $E_{\text{REF}} = E_{\text{AgCl/Ag}}^0 + A \log_{10}(a_{\text{Cl}^-})$. Activity coefficients were estimated by applying the SIT formula : $\log_{10}(\gamma_i) = -z_i^2 D(I_m) + \varepsilon(i, X) m_X$ for an ion i of charge z_i in a X medium, where X is an ion of charge opposite to that of i [3,8, 17]. Numerical values and definitions used are given in table 1.

The voltamograms were recorded between +0.240 and -0.020 V/SHE at a scanning speed of 0.002 V/s. The upper limit of +240 V/SHE was imposed by the oxidation wall of mercury. Ten millilitres of the c_0 M Na_2CO_3 , (I- $3c_0$) M NaHCO_3 test solution was first added to a thermostated (at $19.3 \pm 1.0^\circ\text{C}$) electrochemical cell (of type Tacussel RM06-C + CRSR) and then deoxygenated with argon (HP45 Carboxyque) for one hour. After deoxygenation, a voltamogram of the electrolyte was recorded in order to verify the absence of any electroactive impurities.

In order to determine whether the redox reaction under study involved a polymer or not, a first set of experiments was first carried out with a cerium total concentration $[\text{Ce}]_t$ varying from $2.5 \cdot 10^{-4}$ to $2 \cdot 10^{-3}$ M. Two compositions of test solutions P1 (1.42 M Na_2CO_3 , 0.13 M NaHCO_3 , pH = 10.36) and P2 (1.33 M Na_2CO_3 , 0.37 M NaHCO_3 , pH = 9.86) were investigated. Five determinations of the $E_{1/2}$ potential (the mean value of the oxidation and reduction peak potentials) were performed for each Ce concentration. In order to determine the stoichiometry of the Ce(IV) complex, a second set of experiments was made as follows : $[\text{Ce}]_t$ was kept constants but pH and $[\text{CO}_3^{2-}]$ were varied simultaneously by stepwise acidification of the initial (P1 + 10^{-3} M Ce) test solution by means of CO_2 (Air Products, quality 4.5). The gas was previously passed through a NaClO_4 solution of same molar ionic strength as that of P1. At each step of acidification, the test solution was left for stabilisation with an argon cover at the surface until constant pH (variation < 0.05 pH unit during 10 minutes) was obtained. The voltamogram of the solution was then recorded and the pH was checked again. Five determinations of the $E_{1/2}$ potential were performed for each pH step.

For both sets of experiments (with test solutions of general initial composition c_0 M Na_2CO_3 , (I- $3c_0$) M NaHCO_3), the junction potential arising between the test solution and the REF electrode was measured by means of the cell REF || 0.02 M NaCl , c_0 M Na_2CO_3 , (I-0.02- $3c_0$) M NaHCO_3 || REF', where REF = Ag/AgCl | 0.02 M NaCl , (I-0.02) M NaClO_4 |, REF' is an Ag/AgCl wire, || denotes a liquid-liquid junction (capillary extension) and | denotes liquid-solid contact. The variations of the junction potential as a function of pH were investigated by stepwise acidification of an initial 0.02 M NaCl , c_0 M Na_2CO_3 , (I-0.02- $3c_0$) M NaHCO_3 solution with CO_2 gas, and measuring the electromotive force of the cell rapidly after stabilisation (to avoid the damage to the REF' electrode in direct contact with carbonate ions). After each measurement, the REF electrode was controlled by means of the cell REF' || 0.02 M NaCl , (I-0.02) M NaClO_4 || REF₀, where REF₀ is an electrode of same type as REF. Differences higher than 0.2 mV were never observed. A linear variation of the junction potentials versus pH (with a slope of about 1 mV/pH unit) was found in the pH range under study. The measured $E_{1/2}$ potentials of the Ce(IV/III) couple in carbonate media were corrected for the junction potentials calculated from the experimental regression straight line at the pH of the corresponding test solutions.

The pH was measured with a Radiometer XC161 combined glass electrode connected to a Tacussel ISIS 20000 pH-meter. The Ag/AgCl reference electrode compartment was filled with a 0.02 M NaCl , (I-0.02) M

NaClO₄ solution of same molar ionic strength I as that of the initial NaHCO₃/Na₂CO₃ test solution. As the acidification by CO₂ gas involved a gradual variation of the ionic strength of the test solution, the glass electrode was standardised in activity units (pH = -log₁₀ a_{H⁺} and not -log₁₀ [H⁺]) by means of two NaHCO₃/Na₂CO₃ buffer solutions of same Na⁺ concentration (3.0 M) as that of the initial NaHCO₃/Na₂CO₃ test solution. The theoretical pH of these buffer solutions were calculated on the basis on their chemical compositions, determined by acid-base titration, and by using the equilibrium constants and SIT coefficients from table 1. The precision of the pH measurement for the buffer solutions was about 0.05 pH unit. In addition, the accuracy of the pH electrode response was controlled by means of a standardisation between pH 7 to 12. The pH range from 8.2 to 12 was checked by a titration of a 1 M NaHCO₃ solution with a 1 M NaOH solution, while the pH range under 8.2 was controlled by measuring the pH of a 1 M NaHCO₃ solution in equilibrium with CO₂ gas. The slope of the glass electrode (figure 1) was found to be the theoretical one (58.03 mV/pH unit at 19.3°C) within a precision of 0.05 pH units in the pH range between 7.5 to 10. Above pH=10, a systematic error varying between 0.05 to 0.6 pH unit was observed. Because of the limited solubility of NaHCO₃ (about 1 M), the control of the accuracy of the pH electrode could not be carried out with concentrated carbonate solutions (about 1.5 M) similar to that used for the electrochemical study of cerium.

Treatment of the data

Preliminary experiments showed that the interval between the oxidation and reduction peaks of Ce(IV/III) increased with the potential scanning rate, which was an indication of a quasi-reversible redox system. For reversible or quasi-reversible systems, the cyclic voltametry curves are theoretically symmetrical around the (constant) half-wave potential E_{1/2}, whatever the potential scanning rate. This feature was observed for scanning rates varying from 0.002 to 0.008 V.s⁻¹ (at higher scanning rates, the shift of the oxidation peak towards the oxidation wall of mercury was such that the oxidation peak potential could not be measured any more). In addition, the current intensity of the oxidation and reduction peaks varied linearly as a function of the square root of the scanning rate, with similar absolute values of the slope during oxidation (+0.16) and in reduction (-0.11). We then assumed that the Ce(III) and Ce(IV) diffusion coefficients were similar and that the experimentally measured E_{1/2} potential could be used as an estimate of the formal redox potential E_{IV/III} of the Ce(IV/III) couple.

The speciation at varying pH and [CO₃²⁻] was calculated from the experimental conditions (c_{Na₂CO₃}, c_{NaHCO₃}, pH) using equations (1 to 8) and the definitions and numerical values listed in table 1. The Na₂CO₃ and NaHCO₃ initial (molal) total concentrations are denoted m_{Na₂CO₃} and m_{NaHCO₃}. Since m_{H⁺} could always be neglected, the approximation γ_{H⁺} = 1 was used in the mass balance equation (4) and in the definition of the ionic strength I (Table 1).

$$\log_{10} p \approx q_{\text{Na}_2\text{CO}_3} m_{\text{CO}_3^{2-}} + q_{\text{NaHCO}_3} m_{\text{HCO}_3^-} \quad (1)$$

$$\log_{10} a_{\text{H}_2\text{O}} \approx r_{\text{Na}_2\text{CO}_3} m_{\text{CO}_3^{2-}} + r_{\text{NaHCO}_3} m_{\text{HCO}_3^-} \quad (2)$$

$$m_{\text{Na}^+} = (2 m_{\text{Na}_2\text{CO}_3} + m_{\text{NaHCO}_3}) \frac{1 + m_{\text{HCO}_3^-} M_{\text{H}_2\text{O}} / 2}{1 + m_{\text{NaHCO}_3} M_{\text{H}_2\text{O}} / 2} \quad (3)$$

$$m_{\text{Na}^+} + m_{\text{H}^+} = 2 m_{\text{CO}_3^{2-}} + m_{\text{HCO}_3^-} + m_{\text{OH}^-} \quad (4)$$


$$K'_{\text{m1}} = K_{\text{m1}} / K_{\text{me}} = \frac{m_{\text{CO}_3^{2-}}}{m_{\text{HCO}_3^-} m_{\text{OH}^-}} \quad (5)$$

$$D(I_{\text{m}}) = \frac{0.5050 \sqrt{I_{\text{m}}}}{1 + 1.5 \sqrt{I_{\text{m}}}} \quad (6)$$

$$\log_{10} K_{\text{me}}(I_{\text{m}}) = \log_{10} K_{\text{me}}(0) + D(I_{\text{m}}) - \varepsilon(\text{Na}^+, \text{OH}^-) m_{\text{Na}^+} + \log_{10} a_{\text{H}_2\text{O}} \quad (7)$$

$$\log_{10} K_{\text{m1}}(I_{\text{m}}) = \log_{10} K_{\text{m1}}(0) + 3 D(I_{\text{m}}) - ([\varepsilon(\text{Na}^+, \text{CO}_3^{2-}) - \varepsilon(\text{Na}^+, \text{HCO}_3^-)]) m_{\text{Na}^+} \quad (8)$$

Previous studies of the Ce(III) – CO₃²⁻ – H₂O system indicate that the limiting carbonate complex of Ce(III), of stoichiometry Ce(CO₃)₄⁵⁻, is the major soluble species in media of [CO₃²⁻] > 1.0 M [16]. The electrochemical equilibrium between Ce(III) and Ce(IV), and its equilibrium potential are :



$$E = E_{IV/III}^0 + A \log_{10} \frac{a_{IV}^{1/n} a_{H^+}^{i/n}}{a_{III} a_{CO_3^{2-}}^{j/n-4} a_{H_2O}^{i/n}} \quad (10)$$

Introducing the expression of activity coefficient (Table 1) into equation (10) :

$$E = E_{IV/III} + A \log_{10} \frac{m_{IV}^{1/n}}{m_{III}} \quad (11)$$

$$\text{where } E_{IV/III} = E_{IV/III}^0(I) + A \left[\frac{i}{n} \log_{10} m_{H^+} - \left(\frac{j}{n} - 4 \right) \log_{10} m_{CO_3^{2-}} \right] \quad (12)$$

$$\text{and } E_{IV/III}^0(I) = E_{IV/III}^0(0) + A \log_{10} \frac{\gamma_{IV}^{1/n} \gamma_{H^+}^{i/n}}{\gamma_{III} \gamma_{CO_3^{2-}}^{j/n-4} a_{H_2O}^{i/n}} \quad (13)$$

In cyclic voltametry, the half-wave potential $E_{1/2}$ corresponds to equal concentrations of the two oxidation states at the working electrode. As the difference between the diffusion coefficients is usually negligible (as we shall verify that the Ce(IV) complex is mononuclear, this approximation is certainly valid), $E_{1/2}$ is assumed to be equal to $E_{IV/III}$ (equation 11).

Similarly, if Ce^{IV} polynuclear species are formed, we have :

$$(m_{III})_{1/2} = n (m_{IV})_{1/2} = m_{Ce} / 2 \quad (14)$$

where m_{Ce} is the Ce total molal concentration. The introduction of equation (14) into the Nernst relation (11) gives :

$$E_{1/2} = E_{IV/III} + A \left[\left(1 - \frac{1}{n}\right) \log_{10} 2 - \frac{1}{n} \log_{10} n \right] - \left(1 - \frac{1}{n}\right) A \log_{10} m_{Ce} \quad (15)$$

The plot of the experimental data $E_{1/2}$ as a function of $\log_{10} m_{Ce}$ gives (equation 15) a straight line with slope $(1-1/n)$ which we shall use for the determination of the stoichiometric coefficient n of the Ce(IV) complex. For $n = 1$, $E_{1/2} = E_{IV/III}$ (equation 15). When n is known, equation 12 and the data ($E_{IV/III}$, $m_{CO_3^{2-}}$, pH) give the stoichiometric coefficients i and j , and the formal potential $E_{IV/III}^0(I)$ at ionic strength I .

Results and discussion

As $[Ce]_t$ had no influence on $E_{1/2}$ (figure 2), the Ce(IV) complex is mononuclear (that is $n = 1$). The experimental data were then interpreted with the simplified equation (12a) :

$$E_{IV/III} = E_{IV/III}^0(I) + A (i \log_{10} m_{H^+} - (j - 4) \log_{10} m_{CO_3^{2-}}) \quad (12a)$$

The effects of the ionic strength variations during the experiments were estimated to be relatively small : among all the chemical conditions investigated during the redox measurements, the Debye-Hückel term $D(I_m)$ varied from 0.255 to 0.259 and the Na^+ molal concentration varied from 3.03 to 3.10 mol/kg. This induced a maximum variation of $\log_{10}(\gamma_{CO_3^{2-}})$ of 0.015 (from -1.280 to -1.265) which corresponds to a variation of the potential of less than 1mV (which is negligible). In addition, all the attempts to fit the experimental data according to equations (12) and (13) by taking into account the variations of the activity coefficients (by using the SIT formula) proved to be unsuccessful because the ionic strength changes were not large enough. Hence, all activity coefficients and the $E_{IV/III}^0(I)$ potential, could be considered as constant. The small ionic strength variations were neglected in the qualitative slope analysis (figures 3 and 4) but not for the following calculations. All the experimental data could be interpreted with the simplified equation (12a) where the i and $(j - 4)$ coefficients had clearly to be set to 0 and 2 respectively (figures 3 and 4). Nevertheless, other sets of (i, j) coefficients were tried e.g. the $Ce(CO_3)_5^{6-}$ species which gave poorer standard deviation and some systematic deviations, while species such $CeHCO_3(CO_3)_4^{5-}$, $CeHCO_3(CO_3)_5^{7-}$, $CeOH(CO_3)_5^{7-}$ or $CeOH(CO_3)_6^{9-}$ gave theoretical curves completely erroneous when compared with experimental data. Finally the following interpretation is proposed :



$$\text{and } E_{IV/III} = E_{IV/III}^0(I) - A 2 \log_{10} m_{CO_3^{2-}} \quad (12b)$$

A value of $E_{IV/III}^0(I) = 0.155_3 \pm 0.002_3$ V/SHE (where the uncertainty is 1.96 times the standard deviation) was fitted. Assuming partial dissociation of the Ce(IV) limiting complex did not significantly improve the fitting, which gave $\log_{10} k_{6m}^{IV} = 0.459$ and $E_{IV/III}^0(I) = 0.161_5 \pm 0.001_6$ V/SHE. In order to make $E_{IV/III}^0(I)$

consistent with the first estimation, the uncertainty was increased and k_{6m}^{IV} was considered as a maximum possible value (table 1).

The Ce(IV/III) redox potential shift between carbonate and non complexing media is determined by the ratio of the Ce(III) and Ce(IV) complexing constants. To avoid ionic strength correction we used the $\beta_{m4}^{III}(I_m)$ and $\beta_{m6}^{IV}(I_m)$ definitions given in table 1. For these constants, as for $E_{IV/III}^0(I_m)$, the numerical values are valid for a given ionic strength and a given Na^+ concentration (here 3.06 mol.kg^{-1}), whatever the major anion in the aqueous solution. The thermodynamical cycle is calculated in the following way using the definitions given in table 1.

$$\begin{aligned} E &= E_{IV/III} + A \log_{10} \frac{m_{IV}}{m_{III}} & (11a) \\ &= E_{Ce^{4+}/Ce^{3+}}^0(0) + A \log_{10} \frac{a_{Ce^{4+}}}{a_{Ce^{3+}}} \\ &= E_{Ce^{4+}/Ce^{3+}}^0(0) + A [\log_{10} \beta_{m4}^{III}(I_m) - \log_{10} \beta_{m6}^{IV}(I_m) - 2 \log_{10} m_{CO_3^{2-}} + \log_{10} \frac{m_{IV}}{m_{III}}] \end{aligned}$$

Comparing with equation (12b) the potential shift is

$$E_{IV/III}^0(I_m) - E_{Ce^{4+}/Ce^{3+}}^0(0) = A [\log \beta_{m4}^{III}(I_m) - \log \beta_{m6}^{IV}(I_m)] \quad (16)$$

This last formula was used to estimate the formation constant of the Ce(IV) limiting carbonate complex (table 1).

Cyclic voltametry was successfully used at high ionic strength but some difficulties arose at lower ionic strength, probably due to precipitation when the Ce(IV) limiting carbonate complex is dissociated. This work confirms that Ce(IV) forms the $Ce(CO_3)_6^{8-}$ complex, while the U and transU limiting carbonate complexes are rather $An(CO_3)_5^{6-}$ in similar chemical conditions ($0.7M < [CO_3^{2-}] < 2M$). This result corroborates the conclusions of Salvatore et al. [15] obtained at higher ionic strength, but the interpretation of our data is more straightforward since our method avoided any Na^+ activity coefficient correction. However, our methodology did not allow the determination of the Na^+ content of the $Ce(CO_3)_5^{6-}$ complex, as suggested in reference [15]. The estimation of the $Ce(CO_3)_5^{6-}$ formation constant value (table 1) is one order of magnitude or slightly higher than for actinide(IV) [1]. The chemical form, $Ce(CO_3)_6^{8-}$, of the limiting carbonate complex of Ce(IV) is consistent with the known coordination chemistry of Ce(IV) and Th(IV), e.g. the NO_3^- group, which is isoelectronic with the CO_3^{2-} group, gives rise to 12-coordinated nitrate complexes like $Ce(NO_3)_6^{2-}$ and $Th(NO_3)_6^{2-}$ in which the 6 bidentate groups define a nearly regular icosahedron [22,23] (In that respect, it seems that the size of the metal is not necessarily a dominant factor [22]). In addition, the same icosahedral coordination geometry was found for the $Th(CO_3)_6^{8-}$ anion in the solid state structure of tuliokite $Na_6BaTh(CO_3)_6.6H_2O$ [14]. Further information about the molecular and solid state structure of the limiting carbonate complex of Ce(IV) could be obtained by using XRD and EXAFS spectroscopy techniques, which have already been used successfully to clarify the structure of the limiting complex $Pu(CO_3)_5^{6-}$ [9]. The ratio $\beta_6^{IV}/\beta_4^{III}$ of 28.6 calculated in the present work at $I_m = 4.33 \text{ mol.kg}^{-1}$ seems consistent with the reported estimation of 25 (using very approximate ionic strength corrections) at $I_m = 12 \text{ mol.kg}^{-1}$ [15] : the difference is in the order of magnitude of the activity coefficient (or ion-pairing, as proposed in [15]) corrections. As for Ce(IV), no evidence of any mixed $OH^-CO_3^{2-}$ or polynuclear Ce(IV) complex was found. For quantitative comparison with actinides(IV), further work could consist in studying the $Ce(CO_3)_6^{8-}$ dissociation into $Ce(CO_3)_5^{6-}$ (probably), and performing a similar investigation on the Am(IV)/Am(III) system [6] with proper junction potential measurements and I corrections [7, 8]. Similar electrochemical studies referring to the Np(IV) [4] and Pu(IV) [5, 18] limiting carbonate complexes would be more tedious as those species are involved in irreversible redox couples only.

References

- [1] Vitorge, P. : CEA-BIB-246 (1995).
- [2] Ciavatta, L., Ferri, D., Grenthe, I., Salvatore, F., Spahiu, K. : Inorg. Chem., 22 (1983) 2088-2092.

- [3] Grenthe, I., Fuger, J., Konings, R.J.M., Lemire, R.J., Muller, A.B., Nguyen-Trung, C., Wanner, H. : *NEA-TDB Chemical thermodynamics of uranium*. Elsevier Science Publishers B.V., 1992, 715p.
- [4] Delmau, L.H., Vitorge, P., Capdevila, H. : CEA-N-2807 (1995).
- [5] Capdevila, H., Thèse de l'Université de Paris-Sud (05/06/1992).
- [6] Bourges, J.Y., Guillaume, B., Koehly, G., Hobart, D.E., Peterson, J.R. : *Inorg. Chem.*, 22 (1983) 1179-1184.
- [7] Robouch, P. : Thèse de l'Université Louis Pasteur, Strasbourg (13/11/1987).
- [8] Silva, R., Bidoglio, G., Rand, M., Robouch, P., Wanner, H., Puigdomenech, I. : *NEA-TDB Chemical thermodynamics of americium*. Elsevier Science Publishers B.V., 1995, 374p.
- [9] Clark, D.L., Conradson, S.D., Keogh, D.W., Neu, M.P., Palmer, P.D., Rogers, R.D., Runde, W., Scott, B.L., Tait, C.D. : communication T3-A1 and Neu, M.P., Clark, D.L., Conradson, S.D., Scott, B.L., Palmer, P.D., Reilly, S.D., Runde, W.H., Tait, C.D. : communication T3-B5, *Actinides '97 International Conference, Baden-Baden, Germany, September 21-26, 1997*.
- [10] Dervin, J. : Thèse de l'Université de Reims, France (1973).
- [11] Dervin, J., Faucherre, J. : *Bull. soc. chim. France* n°542 - 11 (1973) 2926-2929.
- [12] Dervin, J., Faucherre, J. : *Bull. soc. chim. France* n°543 - 11 (1973) 2930-2933.
- [13] Haltier, E., Fourest, B., David, F. : *Radiochim. Acta* 51 (1990) 107-112
- [14] Clark, D. L., Hobart, D. E., Neu, M. P. : *Chem. Rev.* 95, (1995) 25-48.
- [15] Salvatore, F., Vasca, E. : *J. Coord. Chem.*, 1990, Vol.21, 237-246.
- [16] Ferri, D., Grenthe, I., Hietanen, S., Salvatore, F. : *Act. Chem. Scand.*, A37 (1983) 359-365
- [17] Grenthe, I., Plyasunov, A.V., Spahiu, K. : *Modelling in Aquatic Chemistry*. pages 325-426 edited by Grenthe, I. and Puigdomenech, I. OECD-NEA Paris (1997).
- [18] Capdevila, H., Vitorge, P. : *Radiochim. Acta* 68, 51-62 (1995).
- [19] Pitzer, K.S. : *Activity Coefficients in Electrolyte Solutions*, 2nd edition, edited by Pitzer K.S., pages 73-153 (1991).
- [20] Bard, A.J., Parsons, R., Jordan J. : *Standard Potential in Aqueous Solutions*, Dekker M. Inc., New York (1985), NY 10016, p. 304.
- [21] Sillén, L.G., Martell, A.E. : *Stability constants of metal ion complexes*. Special Publ. No.17, 1964, 754p and Suppl. No. 1, Special Publ. No. 25, 1971, 865p. The Chemical Society, London ; Högfeltd, E. : *Stability constants of metal ion complexes, part A : inorganic ligands*. IUPAC chemical data series (21), 1982, 310p, Oxford.
- [22] Greenwood, N.N., Earnshaw, A. : *Chemistry of the Elements*, Pergamon Press, 1984.
- [23] Cotton, S., *Lanthanides and Actinides*, Macmillan Physical Science Series, 1991.

Table 1. Physical and chemical parameters used or determined in this study.

For a pure electrolyte : $\log_{10} a_{\text{H}_2\text{O}} \approx r m$, $p = m / c$ and $\log_{10} p \approx q m$; where c and m denote the molar (mol.l^{-1} or M) and molal (mol.kg^{-1}) concentrations of the salt in solution. When the electrolyte is a mixture of two salts (Na_2CO_3 , NaHCO_3), the approximations $\log_{10} p \approx q_1 m_1 + q_2 m_2$ and $\log_{10} a_{\text{H}_2\text{O}} \approx r_1 m_1 + r_2 m_2$ are used, where q_x and r_x are the q and r parameters for the pure salt x , and m_x is the molal concentration of the salt x in the mixture. p values (l.kg^{-1}) are taken from [8]. The q and r values are calculated in this work. The $\log_{10} a_{\text{H}_2\text{O}}$ values (to obtain r) are calculated from Pitzer parameters [19]. c_i , m_i , a_i and γ_i denote respectively, the molar (mol.l^{-1} or M) concentration, the molal (mol.kg^{-1}) concentration, the molal activity and the activity coefficient of the species i : $a_i = m_i \gamma_i$ and $p = m_i / c_i$. The coefficients γ_i are calculated by using the SIT formula [3, 8, 1]. The SIT coefficients $\epsilon(\text{M}^{\text{Z}_M}, \text{A}^{\text{Z}_A})$ (kg.mol^{-1}) are taken from [8]; the uncertainty is increased for the SIT coefficients which are estimated by analogy.

Electrolyte X	q_x	r_x	$\epsilon(i,j)$	$j = \text{ClO}_4^-$	$j = \text{OH}^-$	$j = \text{CO}_3^{2-}$	$j = \text{HCO}_3^-$	$j = \text{Cl}^-$	$j = \text{Ce}(\text{CO}_3)_4^{5-}$
NaHCO₃	0.01249	-0.0148 ± 0.0015	i = Na⁺		0.04 ± 0.01	-0.08 ± 0.03	0.00 ± 0.02	0.03 ± 0.01	-0.20 $\pm 0.25^a$
Na₂CO₃	0.00426	-0.0136 ± 0.0002	i = Ce³⁺	0.51 $\pm 0.04^b$					
M_{H2O} = 0.0180 kg.mol.⁻¹			i = Ce⁴⁺	0.76 $\pm 0.20^c$					

^aEstimated by analogy with $\text{U}(\text{CO}_3)_4^{4-}$ and $\text{U}(\text{CO}_3)_4^{6-}$. ^bestimated from [19] by analogy with La and Pr, 0.47 ± 0.03 is tabulated in [3, 8] ^cEstimated by analogy with U^{4+} .

Table 2. Thermodynamic equilibrium constants and redox potentials used or determined in this work.

The equilibrium constants are defined in molal units (as in the text, and when nothing else is stated for numerical values). Subscript m is used for molal units (only when needed in order to make the difference with molar units). $I_m = \frac{1}{2} \sum z_i^2 m_i$ is molal ionic strength. $A = R T \ln 10 / F$, typically 0.05916 and 0.05803 V at 25 and 19.3°C. IV and III denote the Ce(IV) and Ce(III) limiting carbonate complexes. For each constant, the first value is usually the numerical datum taken from the corresponding reference, while values at other ionic strengths are calculated in this work by means of the SIT formula [3] by using the ϵ values listed in table 1. The $E_{IV/III}^0(4.33m)$ (V/SHE) value was measured in this work in a 0.46 m NaHCO₃ 1.29 m Na₂CO₃ aqueous solution at 19.3°C. $\beta_{m6}^{IV}(4.33m)$ is calculated with equation (16) by using the necessary auxiliary data at about 25°C (the temperature corrections were assumed to be less than uncertainty).

Name	Definition	Numerical value	Reference
$K_{mw}(I_m)$	$= m_{OH^-} a_{H^+}$	$\log_{10} [K_w(0)] = -14.001 \pm 0.14^{(1)}$	[17] page 371
$K_{m1}(I_m)$	$= \frac{m_{CO_3^{2-}} a_{H^+}}{m_{HCO_3^-}}$	$\log_{10} [K_1(0)] = -10.329 \pm 0.020^{(1)}$	[8]
$E_{AgCl/Ag}^0(I_m)$	$\approx E_{AgCl/Ag}^0(0) - A \log_{10} a_{Cl^-}$	$E_{AgCl/Ag}^0(0) \approx 0.2263^{(1)}$ at 19.3°C	[20]
$E_{Ce^{+4}/Ce^{+3}}^0(I)$	$\approx E_{1/2}$ in acidic media of ionic strength I	$E_{Ce^{+4}/Ce^{+3}}^0(I) = 1.72 \pm 0.02^{(2)}$ $1.821 \pm 0.023^{(1)}$ at I = 0	[21]
$\beta_4^{III}(I)$	$= \frac{[Ce(CO_3)_4^{5-}]}{[Ce^{3+}][CO_3^{2-}]^4}$	$\beta_4^{III}(I) = 13.7 \pm 0.1^{(3)}$ at $I_m = 3.5$ $11.67 \pm 0.98^{(1)}$ at I = 0	[16]
$\beta_{m4}^{III}(I_m)$	$= \frac{m_{Ce(CO_3)_4^{5-}}}{a_{Ce^{3+}} m_{CO_3^{2-}}^4}$	$\beta_{m4}^{III}(I_m) = 13.51 \pm 0.16^{(4)}$ at $I_m = 3.06$ $13.61 \pm 0.16^{(5)}$ at $I_m = 4.33$	
$E_{IV/III}^0(I_m)$	$\approx E_{1/2} + 2 A \log_{10} m_{CO_3^{2-}}$	$E_{IV/III}^0(I_m) = 0.182 \pm 0.009^{(4)}$ at $I_m = 3.06$ $0.161 \pm 0.008^{(5)}$ at $I_m = 4.33$	this work
$\beta_{m6}^{IV}(I_m)$	$= \frac{m_{Ce(CO_3)_6^{8-}}}{a_{Ce^{4+}} m_{CO_3^{2-}}^6}$	$\lg \beta_6^{IV}(I_m) = 41.8 \pm 0.5^{(4)}$ at $I_m = 3.06$ $42.2 \pm 0.5^{(5)}$ at $I_m = 4.33$	this work
$k_{m6}^{IV}(I_m)$	$= \frac{m_{Ce(CO_3)_6^{8-}}}{m_{Ce(CO_3)_5^{6-}} m_{CO_3^{2-}}}$	$\lg k_6^{IV}(I_m) \leq 0.03 \pm 0.47^{(4)}$ at $I_m = 3.06$ $0.03 \pm 0.19^{(5)}$ at $I_m = 4.33$	this work
$\beta_{m5}^{IV}(I_m)$	$= \frac{m_{Ce(CO_3)_5^{8-}}}{a_{Ce^{4+}} m_{CO_3^{2-}}^5}$	$\lg \beta_5^{IV}(I_m) \leq 41.6 \pm 0.5^{(4)}$ at $I_m = 3.06$ $41.8 \pm 0.5^{(5)}$ at $I_m = 4.33$	this work

Footnotes :

⁽¹⁾Standard value (at 0 ionic strength).⁽²⁾1 M HClO₄ ; this value may be in error since the hydrolysis correction does not seem to be validated, this possible error would propagate on $\beta_{m6}^{IV}(I_m)$ and $\beta_{m5}^{IV}(I_m)$ values. Still our E^0 value calculated at I=0 is consistent with the value selected in [15].⁽³⁾3 M NaClO₄.⁽⁴⁾2.67 M NaClO₄ which corresponds to Na⁺ molality = $I_m = 3.06$ mol.kg⁻¹.⁽⁵⁾Na₂CO₃ / NaHCO₃ media as used in the present work ; Na⁺ molality = 3.06 ± 0.04 mol.kg⁻¹ and $I_m = 4.33 \pm 0.37$ mol.kg⁻¹.

Table 3: Ce(IV/III) formal potential measured in concentrated carbonate/bicarbonate media

From the Na_2CO_3 and NaHCO_3 initial concentrations (noted $[\text{Na}_2\text{CO}_3]$ and $[\text{NaHCO}_3]$) and the pH values measured while the acidic titration with CO_2 gas, the speciation is calculated by solving the system of equations (1 to 8) with the parameters listed in tables 1 and 2. $[i]$ and m_i denote the molar (mol.l^{-1} or M) and molal (mol.kg^{-1}) concentrations of the species i . The glass electrode is calibrated in activity unit, with buffers at the same molar ionic strength as that of the initial test solutions. The potential $E_{1/2}$, mean value of the oxidation and reduction peaks, is measured on the voltamograms. It is then corrected for the junction potential E_j (mV), and the reference electrode potential E_{REF} (mV/SHE) (calculated with the parameters of table 1 as indicated in the text) to give $E_{\text{IV/III}}$ (mV/SHE).

Initial conditions			Measurements		Values calculated from initial conditions						$E_{\text{IV/III}}$
$[\text{Na}_2\text{CO}_3]$	$[\text{NaHCO}_3]$	$[\text{Ce}]_t$	E_j	pH	I_m	m_{Na^+}	$m_{\text{HCO}_3^-}$	$m_{\text{CO}_3^{2-}}$	m_{OH^-}	E_{REF}	
1.468	0.084	0.001	12.5	10.552	4.564	3.071	0.085	1.493	4.88E-04	329.8	136.1
1.468	0.084	0.001	12.3	10.276	4.531	3.073	0.157	1.458	2.58E-04	329.8	138.3
1.420	0.120	0.001	13.0	10.388	4.487	3.032	0.122	1.455	3.35E-04	329.8	137.7
1.420	0.120	0.001	12.3	9.766	4.334	3.041	0.457	1.292	7.97E-05	329.8	143.9
1.420	0.120	0.001	12.1	9.563	4.235	3.047	0.673	1.187	4.98E-05	329.8	147.3
1.420	0.120	0.001	11.9	9.388	4.125	3.054	0.912	1.071	3.32E-05	329.8	150.9
1.420	0.120	0.001	11.8	9.294	4.057	3.058	1.059	0.999	2.67E-05	329.8	153.4
1.422	0.127	0.001	13.8	10.361	4.473	3.026	0.130	1.448	3.15E-04	329.8	138.1
1.422	0.127	0.001	13.3	9.844	4.354	3.033	0.391	1.321	9.55E-05	329.8	141.5
1.422	0.127	0.001	13.1	9.616	4.253	3.039	0.609	1.215	5.64E-05	329.8	144.4
1.422	0.127	0.001	12.9	9.440	4.150	3.045	0.835	1.105	3.75E-05	329.8	149.0
1.422	0.127	0.001	12.8	9.329	4.073	3.049	1.002	1.024	2.90E-05	329.8	152.6
1.422	0.127	0.001	12.8	9.257	4.019	3.053	1.119	0.967	2.45E-05	329.8	155.7
1.462	0.087	0.001	14.4	10.536	4.551	3.063	0.088	1.487	4.70E-04	329.8	134.6
1.462	0.087	0.001	14.1	9.943	4.445	3.070	0.320	1.375	1.20E-04	329.8	138.7
1.462	0.087	0.001	13.9	9.671	4.338	3.076	0.552	1.262	6.38E-05	329.8	143.0
1.462	0.087	0.001	13.8	9.494	4.241	3.082	0.764	1.159	4.24E-05	329.8	147.7
1.462	0.087	0.001	13.7	9.305	4.111	3.090	1.047	1.022	2.73E-05	329.8	153.4
1.326	0.369	2.49E-04 to 1.96E-03	13.3	9.864	4.452	3.094	0.378	1.358	9.95E-05	329.8	141.2
1.422	0.127	2.49E-04 to 1.96E-03	13.8	10.361	4.473	3.026	0.130	1.448	3.15E-04	329.7	137.0
1.484	0.078	0.001	13.2	10.584	4.608	3.099	0.079	1.509	5.23E-04	329.7	134.5




Fig. 1. Glass electrode calibration. The calibration was checked by two titrations of a 1 M NaHCO_3 solution with a 1 M NaOH solution at $T = 19.3 \pm 1.0^\circ\text{C}$. The theoretical pH ($-\log_{10} a_{\text{H}^+}$ and not $-\log_{10} [\text{H}^+]$) values were calculated by using the equilibrium constants and SIT coefficients listed in tables 1 and 2. The observed deviation at $\text{pH} > 10$ was taken into account in the treatment of the experimental data of the Ce(IV)/Ce(III) redox potential. No deviation (of the glass electrode slope) was observed at $\text{pH} < 8.2$ where $\text{NaHCO}_3/\text{CO}_{2(\text{g})}$ buffers were used for the pH calibration.




Fig. 2. Influence of the cerium total concentration on the Ce(IV)/Ce(III) redox potential in concentrated carbonate/bicarbonate media. Two sets of experiments were performed: $[\text{Na}^+] = 3.0$ (or 3.1) mol.kg^{-1} , $[\text{CO}_3^{2-}] = 1.45$ (or 1.36) mol.kg^{-1} , $[\text{HCO}_3^-] = 0.13$

(or 0.38) mol.kg⁻¹ and pH = 10.36 (or 9.86) respectively, T = 19.3 ± 1.0°C, A = 58.03 mV/log₁₀ unit. This figure shows that the measured redox potential E_{IV/III} is independent of the Ce total concentration. Since in these conditions the Ce(III) major aqueous species is mononuclear [16], the Ce(IV) species is also mononuclear (equation 15). Theoretical curves are plotted for n = 1 (solid line) and n = 2 (dotted line), where n is the Ce stoichiometric coefficient in the Ce(IV) aqueous species (for n > 2 the deviation between the experimental data and the theoretical lines are even greater than for n = 2).




Fig. 3. Influence of the CO₃²⁻ concentration on the Ce(IV/III) formal potential in concentrated carbonate/bicarbonate media. [Na⁺] = 3.06 mol.kg⁻¹, 1.00 < [CO₃²⁻] < 1.45 mol.kg⁻¹, [Ce]_t = 0.001 M, 9.2 < pH < 10.6, T = 19.3 ± 1.0°C, A = 58.03 mV/log₁₀ unit. This figure shows that, within experimental uncertainties, the measured data falls into a single straight line (dotted line), with a slope corresponding to the theoretical one assuming that two CO₃²⁻ ions are exchanged during the redox reaction. Since in these conditions the Ce(III) major species is Ce(CO₃)₄⁵⁻ [16], the Ce(IV) species is Ce(OH)_i(CO₃)₆⁸⁻ⁱ; As the pH influence plot (figure 4) confirms that i=0, the major Ce(IV) complex is then Ce(CO₃)₆⁸⁻. That species is possibly partially dissociated into Ce(CO₃)₅⁶⁻ (bolded curve) as proposed in this work (see text).




Fig. 4.

Fig. 4. pH influence on the Ce(IV/III) formal potential in concentrated carbonate/bicarbonate media. $[\text{Na}^+] = 3.06 \text{ mol.kg}^{-1}$, $1.00 < [\text{CO}_3^{2-}] < 1.45 \text{ mol.kg}^{-1}$, $[\text{Ce}]_t = 0.001 \text{ M}$ at $T = 19.3 \pm 1.0^\circ\text{C}$, $A = 58.03 \text{ mV}/\log_{10} \text{ unit}$. The experimental data are the same as in figure 3. This figure shows that the corrected redox potential $E_{\text{IV/III}} - E^0_{\text{IV/III}}(I) + A 2 \log_{10} m_{\text{CO}_3^{2-}}$ (equation 12a) is independent of pH. The Ce(IV) and Ce(III) aqueous species have then the same OH^- stoichiometric coefficient ($i = 0$): since the Ce(III) and Ce(IV) major species are $\text{Ce}(\text{CO}_3)_4^{5-}$ [16] and $\text{Ce}(\text{OH})_i(\text{CO}_3)_6^{8-i}$ (Figure 3), the formula of the the Ce(IV) complex is finally $\text{Ce}(\text{CO}_3)_6^{8-}$, which is in agreement with the interpretation of the figure 3.