

Aryl-Alkyl Cross-coupling *versus* Homo-coupling: A Mechanistic Study On The Catalytic Performance Of A Bulky Silylamide Iron(II) Complex

Lidie Rousseau, Guillaume Lefèvre

► **To cite this version:**

Lidie Rousseau, Guillaume Lefèvre. Aryl-Alkyl Cross-coupling *versus* Homo-coupling: A Mechanistic Study On The Catalytic Performance Of A Bulky Silylamide Iron(II) Complex. GreenCat 2020 - 14th International Green Catalysis Symposium, Mar 2020, Rennes, France. cea-02482667

HAL Id: cea-02482667

<https://hal-cea.archives-ouvertes.fr/cea-02482667>

Submitted on 18 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Aryl-Alkyl Cross-coupling *versus* Homo-coupling: A Mechanistic Study On The Catalytic Performance Of A Bulky Silylamide Iron(II) Complex


Lidie Rousseau ^{a,b}, Guillaume Lefèvre*^a

^a CSB2D, Institute of Chemistry for Life and Health Sciences, Chimie ParisTech, PSL University, CNRS, 75005 Paris, France ;

^b LCMCE, NIMBE, CEA, CNRS, Univ. Paris-Saclay, 91191 Gif-sur-Yvette, France ;

lidie.rousseau@chimieparistech.psl.eu

Iron catalysis has raised a great interest in the organometallic field since the 1970s and the first successful Fe-catalyzed cross-coupling reactions developed by Kochi,¹ as it is a cheaper, more earth-abundant and less toxic alternative to many other transition metals. Kumada-type reactions between aryl Grignard reagents and organic electrophiles are of high importance, but a significant issue remains in the competition between effective cross-coupling, and homocoupling of the Grignard reagent affording unwanted symmetrical biaryl byproducts. Nakamura² proposed an explanation for the large presence of this byproduct in the presence of several aryl groups on the iron center resulting from multiple Mg-to-Fe transmetallations, and proposed the addition of a source of fluoride anions to block this possibility. However, our study of the catalytic behavior of the bulky silylamide iron(II) complex³ [Fe(N(SiMe₂Ph)₂)₂] in an aryl-alkyl cross-coupling reaction refutes the hypothesis that the multi-transmetallations are the only cause of homocoupling. The steric hindrance of the ligands effectively prevents more than one transmetallation on the iron atom, affording a stable Ar–Fe bond, but it does not quite impede the homocoupling. Stoichiometric and catalytic tests involving ¹H paramagnetic NMR, EPR spectroscopy and cyclic voltammetry were lead in order to discuss the mechanism of the reaction, evidencing transient Fe^{III} species responsible for the formation of the homocoupling product.



¹ Kochi, J. K.; Tamura, M.; *J. Am. Chem. Soc.* **1971**, *93* (6), 1483–1485.

² Hatakeyama, T.; Nakamura, M.; *J. Am. Chem. Soc.* **2007**, *129* (32), 9844–9845.

³ Chen, H.; Bartlett, R. A.; Dias, H. V. R.; Olmstead, M. M.; Power, P. P.; *J. Am. Chem. Soc.* **1989**, *111* (12), 4338–4345.