



HAL
open science

Water radiolysis quantification upon soft X-rays exposure using a microfluidic cell

Lucie Huart, Christophe Nicolas, Marie-Anne Herve Du Penhoat, Jean Philippe Renault, Corinne Chevallard, Jean-Michel Guigner

► **To cite this version:**

Lucie Huart, Christophe Nicolas, Marie-Anne Herve Du Penhoat, Jean Philippe Renault, Corinne Chevallard, et al.. Water radiolysis quantification upon soft X-rays exposure using a microfluidic cell. RD11 - 11th International Workshop on X-ray Radiation Damage to Biological Samples, Mar 2020, Villigen, Switzerland. cea-02491898

HAL Id: cea-02491898

<https://cea.hal.science/cea-02491898>

Submitted on 26 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Contribution ID: 23

Type: Poster

Water radiolysis quantification upon soft X-rays exposure using a microfluidic cell

Since the discovery of X-rays, their use has grown steadily. The best example are synchrotron radiation sources. They provide flexible, powerful methods for learning about the structure and behaviour of matter at the molecular and atomic levels. However, they can induce sample damages, intended or not(1,2). Soft X-rays (keV range) are particularly interesting because of their wide application for structural studies (microscopy, spectroscopy, XFEL), but also because they were shown efficient at inducing complex DNA damage leading to cell death(3).

Combining microfluidic, that offers an adaptable and well-defined sample environment, and sodium benzoate, as a fluorescent dosimeter, the study proposes a sensitive way to measure radical production upon soft X-rays exposure, despite their very low penetration in liquids. The study specifically focuses on ionizing the K-shell electrons of metal ions. Indeed, theoretical studies predict that fast electron transfer from the molecule toward the metallic ion, leaving behind ionized water molecules, which then could dissociate into OH radical (4). Therefore, the OH scavenging method was applied to solutions containing Mg or Na ions. By varying the X-ray energy below and above the K-shell of these atoms (1317 eV for Mg and 1071 for Na), the OH radicals production was measured in order to probe an eventual threshold effect. Our results will be compared with standard gamma-rays exposures and differences will be highlighted.

[1] Gianoncelli et al., Scientific Report (2015)

[2] A. Boissière et al., Radiation Research, 167 (4), 493-500 (2007)

[3] D.T. Goodhead et al., Int. J. Radiat. Biol. (1977).

[4] V. Stumpf, K. Gokhberg, L. S. Cederbaum, Nature Chemistry (2016).

Auteur principal: HUART, Lucie (CEA Saclay)

Co-auteurs: Dr NICOLAS, Christophe (Synchrotron Soleil); Prof. HERVÉ DU PENHOAT, Marie-Anne (IMPMC, Sorbonne Université); Prof. RENAULT, Jean-Philippe (Nimbe, CEA Saclay); Prof. CHEVALLARD, Corinne (NIMBE, CEA Saclay); Dr GUIGNER, Jean-Michel (IMPMC, Sorbonne Université)

Orateur: HUART, Lucie (CEA Saclay)

Session Classification: Poster