

# Modélisation et validation expérimentale de l'interaction rayonnement-milieu cellulaire en radiothérapie par photo-activation d'éléments lourds

R. Delorme, Christophe Champion, Mehdi Gmar, Mathieu Agelou, Hélène Elleaume, Jean-Luc Ravanat

### ▶ To cite this version:

R. Delorme, Christophe Champion, Mehdi Gmar, Mathieu Agelou, Hélène Elleaume, et al.. Modélisation et validation expérimentale de l'interaction rayonnement-milieu cellulaire en radiothérapie par photo-activation d'éléments lourds. 15èmes Journées d'Etudes de la Chimie sous Rayonnement (JECR 2010), May 2010, Bordeaux, France., pp.Affiche A5, 2010. cea-02557645

## HAL Id: cea-02557645 https://cea.hal.science/cea-02557645

Submitted on 1 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

### Modeling and experimental validation of the radiation - DNA interaction in radiotherapy by photon activation of heavy elements

#### DELORME Rachel<sup>1</sup>

Supervisors: Christophe Champion<sup>2</sup>, Mehdi Gmar<sup>1</sup>, Mathieu Agelou<sup>1</sup>, Hélène Elleaume<sup>3</sup>, Jean-Luc Ravanat<sup>4</sup>

CEA, DRT, LIST, DCSI, LM2S : Laboratoire Modélisation, Simulation et Systèmes, F-91191 Gif-Sur-Yvette, France. <sup>2</sup>Laboratoire de Physique Moléculaire et des Collisions, Université Paul Verlaine de Metz , France. 3INSERM U836 / European Synchrotron Radiation Facility (ESRF) ID17, Grenoble, France. 4CEA, DSM, INAC, SCIB, Laboratoire des Lésions des Acides Nucléiques, Grenoble, France

Contact: rachel.delorme@cea.fr

#### CONTEXT

- Malignant brain tumors represent a few percents of adult cancers and the most frequent of children cancers. Because of the delicate location and the radio sensitivity of healthy brain, the current treatments are not efficient for resistant tumors as high-grade gliomas.
- Promising treatment: An innovative approach using X-rays in addition with heavy elements, as iodine or gold nanoparticles.
- Principle: heavy elements introduced in the target volume allow to increase the photoelectric cross section what has for consequence to enhance the dose effect into the tumor.
- Limitation: physical processes and biological impact of this technique are not well understood. The experimental results can not be explained from macroscopic dose calculations.
- Proposition: to consider the radio-induced damages at the cell level.


#### **OBJECTIVES**

- To model and simulate, with a Monte Carlo code, the photon interactions in an inhomogeneous media as cells or DNA, in presence of heavy elements. This will be done at CEA Saclay1 with help of C. Champion<sup>2</sup>.
- To compare the calculations with the Geant4DNA code.
- To validate the code with experimental measurements, on cells or plasmids, carried out in CEA of Grenoble<sup>4</sup> and in the beam line dedicated to medical studies at ESRF3.

#### STATE OF THE ART


- Photon activation therapy with injection of iodinated contrast agent (ICA)
  - First proposition of this technique by Norman and co-workers (1-2), using a modified CT scanner. Performed the first and lonely

human clinical trial on 8 patients bearing brain metastasis and Showed that the method was sure and potentially beneficial.



Development of the technique at the medical beam line of ESRF using monochromatic X-rays in the 50-100 keV range. Several experiments were performed on animals bearing highgrade gliomas to optimize the treatment (3-4).


→ Human clinical trial in preparation.



### Photon activation therapy in presence of gold nanoparticles (GNP)

Hainfeld 2004 (5): first in vivo experiment which proves the advantage of gold nanoparticles in radiotherapy for cancer treatment.

Study: mice bearing a carcinoma on leg. Survival at 1 year : increase until a factor 4.3 for GNP compared to irradiation alone.



- · In vitro studies (cells, protein or DNA) allow to optimize the vectorisation and the radiosensitivity of GNP (6-7).
- Current limitation of the use of GNP in therapy: the cost, the lack of in vivo toxicity studies of GNP and the lack of clinical trials.

#### **Monte Carlo Simulations**


- Macroscopic dose calculation in presence of GNP were made (8). Conclusion: interactions need to be considered at the cellular level to reproduce the in vivo results.
- Monte Carlo codes developed to simulate the radiation--induced damages to cells and DNA following the "track-structure" and "microdosimetry" formalisms (9 -11).
- Geant4DNA: implementation of a new package into the code Geant4 which aims to simulated the physical interactions for low energies (12), the chemical process, the DNA damages and the molecular geometries.

#### FIRST EXPERIMENT

Aim of the study: to show an enhancement of the relative dose-effect in presence of heavy elements with a fractionated dose according to previous results which showed a

Study: Mars 2010 at ESRF. 50 fisher rats bearing F98 gliomas.

such effect for small doses (4).



- Groups: 50 rats separate in 6 groups:
- · irradiation only
- ICA only
- irradiation + ICA
- GNP only
- irradiation + GNP.
- Irradiation: 10 fractions of 4Gy on the right cerebrum from an X-Ray tube (150 kVp, 20 mA).
- Results under way.



#### **BIBLIOGRAPHY**

- Norman et al. (1997). X-ray phototherapy for canine brain masses. Radiat. Oncol. Investig. 5 8-14.
- Rose et al. (1999). A first radiotherapy of human metastatic brain tumors delivered by a computerized tomography scanner(CTRx). Int. J. Radiat. Oncol. Biol. Phys. 45 1127-32
- ADAM et al. (2003). Synchrotron radiation therapy of malignant brain glioma loaded with an iodinated contrast agent First trial on rats bearing F98 gliomas. Int. J. of rad. Oncol. Biol. Phy.. vol. 57, p. 1413-1426.
- 4. Adam JF et al (2006). Prolonged survival of fisher rats bearing F98 glioma after iodine-enhanced synchrotron stereotactic radiotherapy. Int. J. Radiation Oncology Biol. Phys., Vol. 64, No. 2, pp. 603-611, 2006
- 5. J. F. Hainfeld et al(2004). The use of gold nanoparticles to enhance radiotherapy in mice. Phys. Med. Biol. 49 N309-315
- Chithrani et al (2006). Determining the Size and Shape Dependence of Gold Nanoparticle Uptake into Mammalian Cells. Nano letters 2006 Vol. 6, No. 4 662-668
- Emilie Brun et al (2009). Parameters governing gold nanoparticle X-ray radiosensitization of DNA in solution. Colloids and Surfaces B: Biointerfaces 72 128–134
- Cho et al. (2005). Estimation of tumour dose enhancement due to gold nanoparticles during typical ratreatments: a preliminary Monte Carlo study. Physics in Medicine and biology. Vol 50 p N163-N173.
- 9. Pomplun et al (2005). A computer-generated supercoiled model of the pUC19 plasmid. Europ. Biophys. Journal.
- 10.Terrissol et al (2008). On the biological efficiency of I-123 and I-125 decay on the molecular level. Int. J. Radiat. Biol., Vol. 84, No. 12, December 2008, pp. 1063–1068
- 11.Nikjoo et al (2008). Can Monte Carlo track structure codes reveal reaction mechanism in DNA damage and tion therapy? Radiation Physics and Chemistry 77 (2008) 1270– 1279
- 12. Chauvie et al (2007). Geant4 Physics Processes for Microdosimetry Simulation: Design Foundation and Implementation of the First Set of Models. IEEE transactions on nuclear sciences, Vol. 54, N°. 6, Dec 2007.