

**Development of a method to identify
alpha-emitter-bearing- particles in soil samples collected
in the Fukushima Dai-ichi Nuclear Power Plant
(FDNPP) area**

Aurélie Diacre, Anne-Laure Faure, Agnès Moureau, Olivier Marie, Nina Griffiths, O. Evrard, Fabien Pointurier

► **To cite this version:**

Aurélie Diacre, Anne-Laure Faure, Agnès Moureau, Olivier Marie, Nina Griffiths, et al.. Development of a method to identify alpha-emitter-bearing- particles in soil samples collected in the Fukushima Dai-ichi Nuclear Power Plant (FDNPP) area. EGU General Assembly 2020, May 2020, Vienne, Austria. cea-02569292

HAL Id: cea-02569292

<https://hal-cea.archives-ouvertes.fr/cea-02569292>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Development of a method to identify alpha-emitter-bearing-particles in soil samples collected in the Fukushima Dai-ichi Nuclear Power Plant (FDNPP) area

Aurélie Diacre¹, Anne-Lare Faure¹, Agnès Moureau², Olivier Marie¹, Nina Griffiths², Olivier Evrard³, and Fabien Pointurier¹

¹Commissariat à l'Énergie Atomique et aux énergies renouvelables (CEA, DAM, DIF), F-91297 Arpajon, France

²Laboratoire de Radio Toxicologie, CEA, Université Paris-Saclay, 91297 Bruyères-le-Châtel, France

³Laboratoire des Sciences du Climat et de l'Environnement (LSCE/IPSL), Unité Mixte de Recherche 8212 (CEA/ CNRS/UVSQ), Université Paris-Saclay, Gif-sur-Yvette, France

The Fukushima Dai-ichi Nuclear Power Plant (FDNPP) accident that occurred in March 2011 released significant quantities of radionuclides in the environment. So far, most of the research focused on radio-cesium and rarely on actinides. Until now, most of the studies on uranium and plutonium released by FDNPP were conducted on bulk environmental samples (soil, sediment, biota, etc.) and rarely on individual particles. The investigation of individual particles allows working on the FDNPP signature alone compared to studies of bulk material which may also incorporate the signature of global fallout. Accordingly, the objective of the current research is to identify and characterize actinide-bearing particles in soil samples collected in the vicinity of FDNPP to get a better understanding of their formation mechanisms and of their fate in the environment. In order to identify and locate actinide-bearing particles in soil samples, we developed and implemented a method based on alpha-autoradiography (Jaegler et al., 2019), which allows identifying and locating specifically alpha-emitters, including plutonium isotopes ²³⁹Pu, ²⁴⁰Pu and ²⁴¹Pu using a CR39 SSNTD device (Solid State Nuclear Tracks Detector).

Before alpha-autoradiography samples were dried, grinded with an agate mortar and sieved to several cutoffs: 1mm, 700µm, 400µm, 200µm, 100µm and 63µm. Cutoffs were chosen according to the sizes of actinide-bearing particles detected by Satou et al (2018). The absence of significant loss of uranium particles by the sieving process was demonstrated by the analysis of a test soil sample with a ¹³⁷Cs activity below 650 Bq/kg spiked with depleted uranium reference particles. After sieving, the test sample was deposited onto a carbon planchet for secondary ion mass spectrometry analysis in order to determine the particle recovery yield.

Then, the detection of actinide-bearing particles by means of alpha autoradiography was performed on various soil samples collected in the vicinity of FDNPP. Longer exposure times logically improve the detection efficiency of alpha-emitters-bearing particles. However, the exposure time should not exceed two months to limit the impact of emissions from naturally-occurring alpha-emitters.

We will present here the first results of detection of alpha-emitting particles in the analyzed soil samples. Relative impacts of naturally-occurring alpha-emitters ($^{234}\text{U}/^{238}\text{U}$, ^{235}U , ^{232}Th and daughter nuclides) and of plutonium isotopes on alpha-radiography trace observation will be discussed.

The next steps of this study will be to develop and implement methods to sample and isolate alpha-emitting particles from the soil matrix and to characterize them in size, morphology, elemental and isotopic compositions. Full characterization of individual particles will be very helpful to determine their origin and to provide an understanding of their formation process and to determine their mobility and life-duration in the environment.

References:

- H. Jaegler, F. Pointurier, et al., 2019. Method for detecting and characterising actinide-bearing micro-particles in soils and sediment of the Fukushima Prefecture, Japan. *Journal of Radioanalytical and Nuclear Chemistry*, 321 (2019), 57–69. doi:10.1007/s10967-019-06575-w
- Y. Satou, K. Sueki, et al., 2018. Analysis of two forms of radioactive particles emitted during the early stages of the Fukushima Dai-ichi Nuclear Power Station accident. *Geochemical Journal*, 52 (2018), 1-7. doi:10.2343/geochemj.2.0514.