

# Relative Contribution of Rill/Interrill and Gully/Channel Erosion to Small Reservoir Siltation in Mediterranean Environments

Abir Ben Slimane, Damien Raclot, O. Evrard, Mustapha Sanaa, Irene Lefevre, Yves Le Bissonnais

► **To cite this version:**

Abir Ben Slimane, Damien Raclot, O. Evrard, Mustapha Sanaa, Irene Lefevre, et al.. Relative Contribution of Rill/Interrill and Gully/Channel Erosion to Small Reservoir Siltation in Mediterranean Environments. *Land Degradation and Development*, Wiley, 2015, 27 (3), pp.785-797. 10.1002/ldr.2387 . cea-02610586

**HAL Id: cea-02610586**

**<https://hal-cea.archives-ouvertes.fr/cea-02610586>**

Submitted on 18 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1  
2  
3  
4  
5 **1 RELATIVE CONTRIBUTION OF RILL/INTERRILL AND GULLY/CHANNEL**  
6 **2 EROSION TO SMALL RESERVOIR SILTATION IN MEDITERRANEAN**  
7 **3 ENVIRONMENTS**

4 Abir Ben Slimane<sup>a,b,c</sup>, Damien Raclot<sup>c</sup>, Olivier Evrard<sup>d</sup>, Mustapha Sanaa<sup>b</sup>, Irène Lefèvre<sup>d</sup>,  
5 Yves le Bissonnais<sup>e</sup>

6  
7 <sup>a</sup>Abir Ben Slimane  
8 Institut National de Recherche en Génie Rural, Eaux et Forêts.  
9 Rue Hédi EL Karray El Menzah IV - BP N°10, 2080 Ariana, Tunisia  
10 e-mail: [abir.ben.slimane@gmail.com](mailto:abir.ben.slimane@gmail.com)

11  
12 <sup>b</sup>Abir Ben Slimane. Mustapha Sanaa  
13 Institut National Agronomique de Tunis,  
14 43, Avenue Charles Nicolle 1082 -Tunis- Mahrajène (Tunisia)  
15 e-mail: [mustaphasanaa2005@yahoo.fr](mailto:mustaphasanaa2005@yahoo.fr)

16  
17 <sup>c</sup>Abir Ben Slimane. Damien Raclot  
18 IRD-UMR LISAH (INRA-IRD-SupAgro),  
19 2 Place Viala, F-34060 Montpellier (France)  
20 e-mail: [damien.raclot@ird.fr](mailto:damien.raclot@ird.fr)

21  
22 <sup>d</sup>Olivier Evrard. Irène Lefèvre  
23 Laboratoire des Sciences du Climat et de l'Environnement (LSCE/IPSL) – Unité Mixte de  
24 Recherche 8212 (CEA, CNRS, UVSQ),  
25 91198-Gif-sur-Yvette Cedex (France)  
26 e-mail: [Olivier.Evrard@lsce.ipsl.fr](mailto:Olivier.Evrard@lsce.ipsl.fr)

1  
2  
3  
4  
5 27 [Irene.Lefevre@Isce.ipsl.fr](mailto:Irene.Lefevre@Isce.ipsl.fr)  
6  
7

8 28  
9

10 29 <sup>e</sup>Yves le Bissonais

11 30 *INRA-UMR LISAH (INRA-IRD-SupAgro),*

12 31 *2 Place Viala, F-34060 Montpellier (France)*

13 32 e-mail: [lebisson@supagro.inra.fr](mailto:lebisson@supagro.inra.fr)  
14  
15  
16  
17

18 33  
19

20 34 The corresponding author: Damien Raclot

21 35 e-mail: [damien.raclot@ird.fr](mailto:damien.raclot@ird.fr)  
22  
23

24 36 Phone: +33 (0)4 99 61 21 39  
25  
26

27 37 Fax: +33 (0)4 67 63 26 14  
28  
29

30 38  
31

32 39 ABSTRACT  
33  
34

35 40 Reservoir siltation due to water erosion is an important environmental issue in Mediterranean  
36 41 countries where storage of clear surface water is crucial for their economic and agricultural  
37 42 development. The high density of gully systems observed in Mediterranean regions raises  
38 43 the question of their contribution to reservoir siltation. In this context, this study quantified the  
39 44 absolute and relative contributions of rill/interrill and gully/channel erosion in sediment  
40 45 accumulation at the outlet of small Tunisian catchments (0.1-10 km<sup>2</sup>) during the last 15 years  
41 46 (1995-2010). To this end, a fingerprinting method based on measurements of cesium-137  
42 47 and Total Organic Carbon combined with long-term field monitoring of catchment sediment  
43 48 yield was applied to five catchments in order to cover the diversity of environmental  
44 49 conditions found along the Tunisian Ridge and in the Cape Bon region. Results showed the  
45 50 very large variability of erosion processes among the selected catchments, with rill/interrill  
46 51 erosion contributions to sediment accumulated in outlet reservoirs ranging from 20 to 80%.

1  
2  
3  
4  
5 52 Overall, rill/interrill erosion was the dominant process controlling reservoir siltation in three  
6  
7 53 catchments whereas gully/channel erosion dominated in the other two catchments. We  
8  
9 54 identified the presence of marly gypsum substrates and the proportion of catchment surface  
10  
11 55 covered by soil management/conservation measures as the main drivers of erosion process  
12  
13 56 variability at the catchment scale. These results provided a sound basis to propose  
14  
15 57 guidelines for erosion mitigation in these Mediterranean environments and suggested to  
16  
17 58 apply models simulating both rill/interrill and gully/channel erosion in catchments of the  
18  
19 59 region.  
20  
21  
22  
23

24  
25 61 **Keywords:** Erosion. Gully. Rill and interrill. Sediment fingerprinting. Small Mediterranean  
26  
27 62 reservoirs.  
28  
29  
30  
31

## 64 INTRODUCTION

32  
33  
34 65 Water erosion is considered to be one of the main causes of land degradation and is a major  
35  
36 66 threat to the soils worldwide (Cerdan et al., 2010; Lal, 2001; Mendal & Sharda, 2013; Zhao et  
37  
38 67 al., 2013). The Mediterranean region is particularly prone to erosion and higher sediment  
39  
40 68 yields were reported in this environment than in many other regions across the world  
41  
42 69 (Woodward, 1995; Vanmaercke et al., 2011, 2012). This situation is due to the specific  
43  
44 70 Mediterranean context (Cerdà et al., 2010; Cantón et al., 2011; García-Ruiz et al., 2013)  
45  
46 71 characterised by an erosive climate affecting steep catchments composed of poor soils and a  
47  
48 72 very long history of intense cultivation including some inappropriate farming practices (Cerdà  
49  
50 73 et al., 2009; Laudicina et al., 2014; Raclot et al., 2009). The severe soil losses observed in  
51  
52 74 this region have direct on-site effects by affecting both soil sustainability and agricultural  
53  
54 75 productivity (De Vente & Poesen, 2005) and off-site effects by increasing flood risk, and  
55  
56 76 affecting water quality and quantity. The rapid siltation of numerous artificial reservoirs built  
57  
58 77 during the last decades to mitigate water scarcity represents an important societal issue in  
59  
60

1  
2  
3  
4  
5 78 North African countries, as the decline of surface water storage capacity may greatly impact  
6  
7 79 their agricultural, economic and social development (Ayadi et al., 2010; Ben Mammou &  
8  
9 80 Louati, 2007; Hentati et al., 2010; Habi & Morsli, 2011; Lahlou, 2000). The prediction of the  
10  
11 81 main erosion processes occurring in catchments is therefore essential for guiding the  
12  
13 82 implementation of erosion control measures adapted to the catchment context.  
14  
15

16 83 The dominance of gully erosion as the main source of sediment in Mediterranean  
17  
18 84 environments has been underlined by many authors. Studies conducted at the local scale, i.e.  
19  
20 85 plot or gully scale, in Algeria (Collinet & Zante, 2005; Roose et al., 2000) showed that gully  
21  
22 86 erosion could produce ten to one hundred times more sediment than sheet erosion. Similarly,  
23  
24 87 when moving up to the catchment scale, several studies highlighted the important role of  
25  
26 88 gullies controlling the reservoir siltation in Spain (De Vente et al., 2008) and in Italy (De  
27  
28 89 Vente et al., 2006). In a review on gully contribution to total catchment erosion (Poesen et al.,  
29  
30 90 2003), this feature appeared to provide the dominant source of sediment in all catchments  
31  
32 91 located in Mediterranean regions. For example, in Spanish reservoirs, gully erosion supplied  
33  
34 92 83% (Poesen et al., 1996) of sediment and largely dominated compared to rill and inter-rill  
35  
36 93 erosion. However, similar measurements detailing the respective contribution of individual  
37  
38 94 erosion processes at the catchment scale are rarely available (Porto et al., 2014). Existing  
39  
40 95 estimations are often given for a limited number of erosive events as the collection of  
41  
42 96 required data through detailed topographic surveys is very difficult and time consuming to  
43  
44 97 cover the entire catchment area. As a result, there is a lack of information regarding the  
45  
46 98 contribution of gully erosion to sediment fluxes at the catchment scale over the mid to long-  
47  
48 99 term. In addition, the factors explaining the differences of erosive behaviour (i.e., absolute  
49  
50 100 and relative contribution of the rill/interrill and gully/channel erosion processes) between  
51  
52 101 catchments are poorly understood.  
53  
54

55 102 In this context, sediment fingerprinting may provide an alternative technique to quantify the  
56  
57 103 relative contribution of individual erosion processes at the catchment scale. This approach  
58  
59  
60

1  
2  
3  
4  
5 104 aims to provide useful quantitative information on sources delivering sediment to rivers that is  
6  
7 105 very difficult, time-consuming and expensive to assess using classic monitoring surveys in  
8  
9 106 the field. This technique has been successfully applied to floodplain deposit samples (Collins  
10  
11 107 et al., 2010; Wasson et al., 2010; Wilkinson et al., 2009); to suspended sediment (Collins et  
12  
13 108 al., 1998; Devereux et al., 2010; Mukundan et al., 2010; Nagle et al., 2007; Walling, 2005), or  
14  
15 109 to core reservoir samples (Ben Slimane et al., 2013; Juracek & Ziegler, 2009; Mourrier, 2008)  
16  
17 110 to apportion the respective contributions of surface and subsurface soils as sediment  
18  
19 111 sources. In these studies, the use of fallout radionuclides (in particular, caesium-137) alone,  
20  
21 112 or in combination with other tracers has proven to be powerful in discriminating between  
22  
23 113 subsoil and topsoil sources. Several recent papers (Guzmán et al., 2013; Haddadchi et al.,  
24  
25 114 2013; Walling, 2013) have reviewed the current approaches, advantages and challenges of  
26  
27 115 this technique.

28  
29  
30 116 The objectives of this paper are to: 1/ quantify the mean inter-annual absolute (in  $\text{Mg ha}^{-1} \text{yr}^{-1}$ )  
31  
32 117 and relative (in %) contribution of rill/interrill and gully/channel erosion processes to the outlet  
33  
34 118 sediment yield of several Mediterranean catchments; 2/ investigate the main driving factors  
35  
36 119 that may explain reservoir siltation, and check whether these driving factors change when  
37  
38 120 considering total sediment yield (i.e., sediment providing from rill/interrill plus gully/channel  
39  
40 121 erosion processes) or individual soil erosion processes (either rill/interrill or gully/channel)  
41  
42 122 contribution to catchment sediment yield. To this end, we combined catchment sediment  
43  
44 123 yield measurements derived from field monitoring with a fingerprinting approach applied to  
45  
46 124 reservoir deposits accumulated at the outlet of five catchments covering a large range of  
47  
48 125 environmental conditions found in Maghreb countries.

49  
50  
51  
52 126  
53  
54  
55  
56  
57  
58  
59  
60

## 127 MATERIAL AND METHODS

128 *Study sites*

129 Five rural catchments were selected among a pilot network of more than 30 small hillside  
130 catchments with reservoirs (0.1-10 km<sup>2</sup>) distributed across the Tunisian Ridge and the Cape  
131 Bon (Figure 1). These five catchments (El Hnach, El Melah, Fidh Ali, Kamech, and Sbahia)  
132 were chosen because of their contrasted landscape characteristics as described in Table 1  
133 and their continuous monitoring by the Tunisian Direction of Soil and Water Conservation  
134 (DGAFTA-CES) and the French Research Institute for Development (IRD).

135 The surface area of these catchments ranges from 0.61 to 3.67 km<sup>2</sup>. They drain into small  
136 reservoirs built between 1991 and 1994 with an initial storage capacity ranging from 20,000  
137 to 150,000 m<sup>3</sup> where sediment has accumulated for more than 15 years. In 2009-2010, El  
138 Melah, El Hnach and Fidh Ali reservoirs were completely filled with sediment, but Kamech  
139 and Sbahia lakes were still operational. They are all associated with very low nutrient levels  
140 as generally found in North African rural environments. The drainage density including both  
141 wadis (i.e., dry creeks) and gullies ranges from 48.2 to 158.3 m ha<sup>-1</sup> (Rebai et al., 2012). The  
142 catchments are distributed along an annual rainfall gradient comprised between 285 and 650  
143 mm. The lithology mainly consists of marls (soft substrate), sandstones and limestones (hard  
144 substrates), but their relative surface cover varies from one basin to another. Marly gypsum  
145 was the most widely found in Fidh Ali catchment. Cropland occupied 10 to 70% of the total  
146 surface depending on the catchment. The rest of the surface was covered by scrubland  
147 devoted to grazing, by forests, by gullies/badlands and by a few houses. The main active  
148 erosion processes in these catchments are related to either rill/interrill or gully/channel  
149 (including bank) processes. Contribution of other processes such as mass movements is  
150 negligible. Two catchments were equipped with a large panel of erosion control measures  
151 covering between 30 and 40% of the surface area. They consist in contour bench terraces,  
152 tree planting, small contour stone bunds and a few small stone check-dams installed across

1  
2  
3  
4  
5 153 gullies in upstream positions. The main climate, lithology, topography and land cover  
6  
7 154 characteristics of the studied catchments and their erosion control measures are synthesized  
8  
9 155 in Table 1. An aerial view derived from Google Earth and ground-based photographs  
10  
11 156 showing gullies, land cover, reservoirs and erosion mitigation measures are also provided for  
12  
13 157 each catchment as supplementary material.  
14  
15

16 158

### 19 159 *Hydrological and total sediment yield measurements*

20  
21 160 Monitoring in the reservoir of the five studied catchments was undertaken since the  
22  
23 161 construction of the small reservoirs. This continuous monitoring consisted of instantaneous  
24  
25 162 measurements of rainfall using a tipping bucket rain gauge (0.5 mm) and water levels with a  
26  
27 163 1 cm precision water level gauge recorder. Between 3 and 10 precise bathymetric surveys  
28  
29 164 were also conducted in each reservoir to establish up-to-date depth/volume and  
30  
31 165 depth/surface curves. Reservoir siltation volumes were quantified between successive  
32  
33 166 bathymetric surveys. Sediment concentrations during overflow through the spillway or  
34  
35 167 emptying through bottom drain valve were also measured through manual sampling. All  
36  
37 168 these measurements enable the calculation of i) mean annual rainfall depth and mean  
38  
39 169 rainfall erosivity by calculating the EI30 index (Wischmeier & Smith, 1958); ii) continuous  
40  
41 170 variation of water level within the reservoir; iii) variation of sediment deposits between  
42  
43 171 successive bathymetric surveys and iv) continuous outputs of water and sediment from the  
44  
45 172 reservoir.  
46  
47

48  
49 173 Continuous runoff input into the reservoir was then quantified by drawing a hydrologic budget  
50  
51 174 of the reservoir. In addition, sediment input into the reservoir was computed by drawing  
52  
53 175 sediment budgets for successive bathymetric surveys following the method described in  
54  
55 176 Albergel et al. (1998, 2005). Mean annual runoff (in mm yr<sup>-1</sup>) and sediment yield (in Mg yr<sup>-1</sup>)  
56  
57 177 were finally quantified for each catchment and for the longest period of records available.  
58  
59  
60



1  
2  
3  
4  
5 178 In this study, catchment sediment yield is expressed as an area-specific yield (SSY in Mg ha<sup>-1</sup>  
6  
7 179 yr<sup>-1</sup>) by dividing the reservoir sediment input by the contributing surface area of the  
8  
9 180 catchment. The precision on the SSY evaluation then mainly depends on: (i) the precision of  
10  
11 181 the DEM as derived from bathymetric surveys (estimated to about 10 %); (ii) the precision on  
12  
13 182 volumes discharged over the spillway (5%) and on measured sediment concentrations (30%);  
14  
15 183 (iii) the precision on average silt density (10%). As a result, global error on SSY was  
16  
17 184 estimated to reach about 20% for these small Tunisian reservoirs (Raclot & Albergel, 2006).

#### 185 *Field sampling*

186 Sampling of representative source material was conducted in 2009-2010 within each of the  
187 five catchments. In total, between 10 and 17 samples representative of rill/interrill and  
188 gully/channel source types were collected within each study site. Each sample was  
189 composed by at least five subsamples collected within an approximate 5-m radius around the  
190 sampling point to increase the representativeness of the sample. During sampling, attention  
191 was paid to document the entire range of geomorphological and pedological conditions  
192 observed within the catchments (see supplementary material for sampling source location).  
193 Gully and channel source sampling was restricted to freshly cut sections in the bottom or the  
194 banks (when those features were deeper than 40 cm). The sampling depth of topsoil material  
195 for rill/interrill source was 0-10 cm in tilled cropland (as soil is homogenized in the entire  
196 ploughed layer), and only 0-2 cm in untilled scrubland environments.

197 Two to four sediment cores were also collected in 2009-2010 in each reservoir  
198 simultaneously to the source material sampling (see supplementary material for sampling  
199 core location). Each core covered the entire layer of sediment at the sampling locations in  
200 the reservoirs and ranged from 0.60 to 2.50 m. This was confirmed by the presence of a  
201 more compact soil layer at the base of the core. In addition, the core depths were consistent  
202 with data provided by topographical surveys conducted immediately after the reservoir  
203 construction. The number and location of cores in reservoir deposits were chosen in order to

1  
2  
3  
4  
5 204 collect one core in the vicinity of the dam and additional cores at the outlet of each main  
6  
7 205 tributary delivering material to the reservoir. Ben Slimane et al. (2013) showed that this type  
8  
9 206 of sampling scheme where a limited number of sediment cores are collected at strategic  
10  
11 207 locations within the reservoir offered a good compromise to reduce the cost of laboratory  
12  
13 208 analyses while providing relevant and representative sediment fingerprinting results. Cores  
14  
15 209 were then described at the laboratory before and after drying. A single composite sample of  
16  
17 210 each core was then prepared.

#### 211 *Laboratory analysis*

212 Chemical and radionuclide analyses were conducted on all samples. Total Organic Carbon  
213 (TOC) contents and caesium-137 ( $^{137}\text{Cs}$ ) activities were measured as they were shown to  
214 provide relevant information to apportion the sources that delivered sediment material  
215 accumulated in North African reservoirs built after 1986 (Ben Slimane et al., 2013).  
216 Arguments supporting the selection of these two tracers in the context of this study are  
217 presented in the discussion section. Samples were described, air-dried, hand-disaggregated  
218 and sieved to 2 mm at the *Environmental and Soil Science Laboratory (INAT, Tunisia)*.

219 Activities in  $^{137}\text{Cs}$  were quantified by gamma-spectrometry using the very low-background  
220 coaxial N- and P-types GeHP detectors (Canberra / Ortec) available at the *Laboratoire des*  
221 *Sciences du Climat et de l'Environnement* (Gif-sur-Yvette, France). The detectors were  
222 periodically controlled with internal and IAEA soil and sediment standards and decay-  
223 corrected to the date of sampling (Evrard et al., 2010). Uncertainties on results were  
224 estimated by combining counting statistics and calibration uncertainties. Summing and self-  
225 absorption effects were taken into account by analysing standards with similar densities and  
226 characteristics as the collected samples.

227 Total Organic Carbon (TOC) was measured by high temperature combustion (NF ISO 10694)  
228 at the *Bioemco Laboratory* (Paris, France) for samples collected in Kamech catchment, and  
229 at the *Soil Analysis Laboratory* (Arras, France) for samples collected at the other study sites.

1  
2  
3  
4  
5 230 After the preliminary destruction of organic matter and dispersion of soil and sediment  
6  
7 231 particles, the grain-size distribution was determined based on the principle of laser diffraction  
8  
9 232 using a Beckman Coulter LS 13320 particle size analyser at the *Laboratoire Géosciences*  
10  
11 233 *Montpellier* (Montpellier, France). This device is equipped with an agitator and an adjustable  
12  
13 234 ultrasonicator to maintain uniform suspension, which enables the analysis of particles with  
14  
15 235 diameters comprised between 0.375 and 2,000  $\mu\text{m}$ . Specific Surface Areas (SSA, square  
16  
17 236 metres per cubic metre) were derived from these laser diffraction data.  
18  
19  
20  
21

### 22 238 *Fingerprinting main steps*

23  
24 239 The fingerprinting properties were first corrected in order to take into account the grain size  
25  
26 240 effects on their adsorption onto particles, as  $^{137}\text{Cs}$  and TOC are known to be enriched in the  
27  
28 241 finest (clay to loam-sized) particle fractions (Motha et al., 2003). He and Walling (1996)  
29  
30 242 tested the particle size effects on the adsorption of  $^{137}\text{Cs}$  on soils and sediments and showed  
31  
32 243 that  $^{137}\text{Cs}$  content can be closely represented by a power function of SSA values calculated  
33  
34 244 for the samples, with exponent values varying between 0.6 and 0.8. In this study, the  
35  
36 245 correction was performed using this power function with an exponent value of 0.7 and  
37  
38 246 applied to both  $^{137}\text{Cs}$  activities and TOC content values. Each soil source (i.e. surface topsoil  
39  
40 247 and gully/channel bank) was subsequently characterised by its mean concentration/activity  
41  
42 248 and the standard deviation of each of its fingerprint properties.  
43  
44

45 249 The ability of individual fingerprinting properties to discriminate between the potential soil  
46  
47 250 sources was then confirmed by conducting a non-parametric Kruskal-Wallis *H*-test as  
48  
49 251 suggested by Collins & Walling (2002). A detailed description of this procedure is provided by  
50  
51 252 Evrard et al. (2011). To characterise the properties of both groups of sources, we assumed  
52  
53 253 that their concentrations ( $c_{i,j}$ ) could be represented by a normal distribution (Eq. 1).  
54

$$55$$
$$56$$
$$57 \quad c_{i,j} \approx N(\mu, \sigma^2) \quad (1)$$
$$58$$
$$59$$
$$60$$

255 Where  $j$  is a specific group of sources;  $i$  is a specific fingerprinting property;  $\mu$  is the average  
 256 concentration in fingerprint property  $i$  measured in source  $j$ ; and  $\sigma^2$  (Eq. 2) is the variance of  
 257 the probability distribution of the mean of property  $i$  in source  $j$ .

$$258 \quad \hat{\sigma}^2 = \left( \frac{S.D.}{\sqrt{d}} \right)^2 \quad (2)$$

259 Where  $d$  is the number of independent samples and S.D. is the standard deviation  
 260 associated with the values of the fingerprinting properties measured in the samples.

261 A multivariate mixing model was then used to estimate the relative contribution (in %) of the  
 262 potential sediment sources in each core sediment sample (Eq. 3).

$$263 \quad \begin{bmatrix} \bar{c}_{1,1} & \bar{c}_{1,2} & \dots & \dots & \bar{c}_{1,S} \\ \bar{c}_{2,1} & \bar{c}_{2,2} & \dots & \dots & \bar{c}_{2,S} \\ \dots & \dots & \bar{c}_{i,j} & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \bar{c}_{V,1} & \dots & \dots & \dots & \bar{c}_{V,S} \end{bmatrix} \begin{bmatrix} \hat{\beta}_1 \\ \dots \\ \hat{\beta}_j \\ \dots \\ \hat{\beta}_S \end{bmatrix} = \begin{bmatrix} y_1 \\ \dots \\ y_j \\ \dots \\ y_V \end{bmatrix} \quad (3)$$

264 where  $\bar{c}_{i,j}$  is the mean value of fingerprinting property  $i$  measured in source  $j$ ;  $\hat{\beta}_j$  is the  
 265 coefficient representing the contribution of source  $j$  to river sediment; S corresponds to the  
 266 number of potential sediment sources and V represents the fingerprinting properties selected  
 267 by the Wilk's lambda procedure.

268 The following physical constraints were applied to  $\hat{\beta}_j$  (Eq. 4):

$$269 \quad \sum_{j=1}^S \hat{\beta}_j = 1; \quad 0 \leq \hat{\beta}_j \leq 1 \quad (4)$$

270 These additional constraints ensured that the sum of all source contributions in the sediment  
 271 was equal to one and that each fraction of these contributions lied between zero and one,  
 272 inclusive.

273 Based on the Monte Carlo method, a series of  $p=10,000$  random positive numbers was then  
 274 generated for each fingerprinting property and for each source. The robustness of the source  
 275 ascription solutions  $\beta_j$  was then assessed using a mean 'goodness of fit' (*GOF*) index (Eq. 5;  
 276 Motha et al., 2003).

$$277 \quad GOF = 1 - \left\{ \frac{1}{p} \times \left( \sum_{i=1}^v \frac{|y_i - \sum_{j=1}^S \hat{\beta}_j \bar{c}_{i,j}|}{y_i} \right) \right\} \quad (5)$$

278 We only used the sets of simulated random numbers that obtained a *GOF* index value higher  
 279 than 0.80 in the subsequent steps. The use of the Monte Carlo method allowed the  
 280 calculation of 95% confidence intervals.

281

## 282 RESULTS

### 283 *Analysis of catchment sediment yield*

284 Sediment yield was evaluated for each catchment from field measurements by adding  
 285 sediment stored in the reservoir to sediment exported from the reservoirs during overflow  
 286 through the spillway or emptying through bottom drain valve. Sediment exported from the  
 287 reservoirs during overflow or emptying represented less than 10% of the total sediment  
 288 inputs to the reservoir in all five catchments. This indicates their very high sediment trapping  
 289 efficiency -more than 90%- confirming previous results found for several similar reservoirs  
 290 across the Maghreb region (Albergel et al., 1998).

291 Sediment yield corresponds to reservoir sediment inputs due to the combination of all active  
 292 erosion processes in the catchments. The corresponding monitoring periods are provided in  
 293 Table 2 together with additional hydrological characteristics.

294 Three of the five catchments were characterised by very similar catchment sediment yields

1  
2  
3  
4  
5 295 (between 15 and 17 Mg ha<sup>-1</sup> yr<sup>-1</sup>), whereas Fidh Ali catchment showed a significantly larger  
6  
7 296 sediment yield (38 Mg ha<sup>-1</sup> yr<sup>-1</sup>) and Sbahia catchment a significantly lower sediment yields  
8  
9 297 (10 Mg ha<sup>-1</sup> yr<sup>-1</sup>). The lifetime of the reservoir (number of years to be completely silted) is  
10  
11 298 about 11 years for Fidh Ali and El Hnach; 15 years for El Melah, and about 30 years for  
12  
13 299 Kamech and Sbahia.

14  
15  
16 300

17  
18  
19 301 *Quantification of the relative contribution of rill/interrill and gully/channel erosion to catchment*  
20  
21 302 *sediment yield for the five catchments*

22  
23  
24 303 Both <sup>137</sup>Cs and TOC passed the Kruskal-Wallis test and were used in combination for  
25  
26 304 discriminating sediments sources in the five studied catchments. Their values and the related  
27  
28 305 specific surface areas (SSA) in both source material and composite sediment core samples  
29  
30 306 collected in the different study sites are presented in Table 3.

31  
32  
33 307 A significant difference in <sup>137</sup>Cs and TOC values between both potential sediment sources  
34  
35 308 was observed in all study catchments, with systematically higher values in the topsoil source  
36  
37 309 than in the gully/channel material. Mean <sup>137</sup>Cs activities in the topsoil samples varied from 3.5  
38  
39 310 Bq kg<sup>-1</sup> to 10.1 Bq kg<sup>-1</sup> whereas they were systematically lower than 0.7 Bq kg<sup>-1</sup> in  
40  
41 311 gully/channel samples. Values of <sup>137</sup>Cs exceeding 1 Bq kg<sup>-1</sup> in the topsoil samples indicate  
42  
43 312 that there has been significant caesium atmospheric fallout during the last decades across  
44  
45 313 the entire Tunisian Ridge and Cape Bon. This confirmed previous <sup>137</sup>Cs measurements  
46  
47 314 conducted across the Maghreb region (Damnati et al., 2012; Faleh et al., 2005). The mean  
48  
49 315 TOC value varied from 0.3% and 0.9% in gully/channel material and from 0.5% to 2.5% in  
50  
51 316 topsoil. A recent study conducted in 25 soil samples collected under cropland and 10  
52  
53 317 additional samples collected under forests in Tunisia showed that the organic carbon content  
54  
55 318 of the topsoil ranged from 0.8 to 3.2% for all soils, with a median TOC value of 2.4% for  
56  
57 319 forest soils and 1.4% for cultivated soils (Annabi et al., 2009). The topsoil samples analysed

1  
2  
3  
4  
5 320 in the 5 study sites were characterised by TOC values that are usually found in Tunisian  
6  
7 321 cultivated soils located in similar bioclimatic zones. Compared to TOC contents found in soils  
8  
9 322 of many other parts of the world, the low values measured in Mediterranean environments  
10  
11 323 can be explained by the low precipitation amounts and the high temperatures prevailing in  
12  
13 324 this region, which are favourable to C mineralization. Among the five studied catchments,  
14  
15 325 samples collected in both El Melah and Fidh Ali sites had TOC values < 1% which indicates  
16  
17 326 a poor soil quality, even for Mediterranean areas (Jones et al., 2004). The very low TOC  
18  
19 327 values measured in topsoil samples of these 2 catchments may also be explained by their  
20  
21 328 coarser texture (i.e., lower SSA values in Table 3) that is known to be less effective to store  
22  
23 329 soil organic carbon than fine-textured soils (Meersmans et al., 2012).

24  
25  
26 330 The SSA values measured in core and source material samples of a given catchment  
27  
28 331 remained very similar for 4 of the 5 study sites, which means that enrichment and depletion  
29  
30 332 effects caused by selective mobilisation and transport of sediment were of limited magnitude.  
31  
32 333 A significant enrichment of fine-grained sediment during erosion and transportation between  
33  
34 334 soils and the reservoir was only observed in El Melah catchment.

35  
36  
37 335 Fingerprinting results obtained for the different cores collected in each reservoir were  
38  
39 336 extrapolated to the entire catchment scale by attributing to each core a weighting factor  
40  
41 337 corresponding to its representativeness in terms of sediment volume accumulated in the  
42  
43 338 reservoir as proposed by Ben Slimane et al. (2013). Figure 2 illustrates the results of the  
44  
45 339 source apportionment for each catchment.

46  
47  
48 340 Results showed the contrasted contribution of erosion processes within the five selected  
49  
50 341 catchments, as the soil surface relative contribution to reservoir sediment ranged between 20%  
51  
52 342 and 80% depending on the site. The surface topsoil was the dominant source of sediment in  
53  
54 343 Fidh Ali, El Melah and Kamech sites, whereas gully/channel material dominated in El Hnach  
55  
56 344 and Sbaihia catchments.

345

346 *Rill/interrill and gully/channel erosion contributions to catchment sediment yield*

347 Rill/interrill and gully/channel erosion contributions to catchment sediment yield were  
348 calculated by applying the relative source contribution to the catchment sediment yields  
349 measured in each catchment (Figure 2). Rill/interrill erosion contribution varied from 2.2 Mg  
350 ha<sup>-1</sup> yr<sup>-1</sup> in Sbaihia catchment to 26.5 Mg ha<sup>-1</sup> yr<sup>-1</sup> in Fidh Ali catchment, whereas  
351 gully/channel erosion contribution varied between 3 Mg ha<sup>-1</sup> yr<sup>-1</sup> for Kamech and 11.5 Mg ha<sup>-1</sup>  
352 yr<sup>-1</sup> in Fidh Ali. The lowest rill/interrill erosion contribution of 2.2 Mg ha<sup>-1</sup> yr<sup>-1</sup> obtained in  
353 Sbaihia catchment remained significantly higher than the tolerable soil loss estimated to 1.4  
354 Mg ha<sup>-1</sup> yr<sup>-1</sup> (Verheijen et al., 2009), indicating the severe levels reached by soil erosion along  
355 the Tunisian Ridge and in the Cape Bon region.

356 The range of variation of rill/interrill erosion contribution was larger (from 1 to 12) than  
357 catchment sediment yields and gully/channel erosion contribution (from 1 to 4) considering  
358 the five investigated catchments. Quantification of individual erosion process contributions  
359 therefore provided important information that was not well reflected by catchment sediment  
360 yields.

361 Overall, Fidh Ali catchment was clearly characterized by the highest catchment sediment  
362 yields and the highest rill/interrill and gully/channel erosion contributions. When considering  
363 the other four catchments, the ranking was modified depending of the erosion type  
364 considered. This is illustrated for instance by the fact that Kamech catchment had the second  
365 most important rill/interrill erosion contributions but the lowest gully/channel erosion  
366 contributions.

367

368

## DISCUSSION



1  
2  
3  
4  
5 369 *Selection of <sup>137</sup>Cs and TOC as sediment tracers in this study*

6  
7 370 One of the main challenges associated with the choice of tracers in fingerprinting technique  
8  
9 371 still lies in the need for better accounting for the selectivity/conservative behaviour of  
10  
11 372 sediment and fingerprint properties between the sources of sediments and the collected  
12  
13 373 material (Koiter et al., 2013). A recent extensive review on sediment tracers in water erosion  
14  
15 374 studies (Guzmán et al., 2013) showed that fallout radionuclides are the most extensively  
16  
17 375 used soil redistribution tracers reported in the scientific literature and that <sup>137</sup>Cs is by far the  
18  
19 376 dominant one, especially for medium time scales (tens of years) across a broad range of  
20  
21 377 spatial scales from hillslope and small catchments to large basins. Its use in fingerprinting  
22  
23 378 techniques -alone or in combination with other tracers- has proven to be effective in  
24  
25 379 discriminating between subsoil and topsoil sources in several studies (Owens et al. 1999;  
26  
27 380 Zhang & Walling, 2005, Juracek & Ziegler 2009; Smith et al. 2012). The conservative  
28  
29 381 behaviour of <sup>137</sup>Cs in soil environments has been demonstrated in many studies and detailed  
30  
31 382 arguments may be found in Guzmán et al. (2013) for instance. If biochemical properties have  
32  
33 383 the potential to provide better spatial constraints for sediment sources compared to other  
34  
35 384 fingerprint properties (Koiter et al., 2013), the use of organic constituents as soil redistribution  
36  
37 385 tracers is less usual. One reason is that the behaviour of organic constituents may often be  
38  
39 386 suspected as poorly conservative. Many arguments have been formulated by Ben Slimane et  
40  
41 387 al (2013) regarding the ability of TOC to be used as an additional tracer to <sup>137</sup>Cs in the  
42  
43 388 specific context of recent North African reservoirs (less than 20 years old) : i) autochthonous  
44  
45 389 source of organic constituents in oligotrophic North African reservoir is negligible as proved  
46  
47 390 in a series of more than 20 modern Tunisian reservoirs (Rahaingomanana 1998); ii) TOC  
48  
49 391 degradation in the deposits is also negligible for recent reservoir as the kinetics of TOC  
50  
51 392 degradation were proved to be very slow in oligotrophic lake deposits as demonstrated by  
52  
53 393 Patience et al. (1995) for the Lac du Bouchet (France); iii) terrestrial organic residues  
54  
55 394 probably did not experience major changes via bacterial alteration during their transport,  
56  
57 395 settling and incorporation into the sediment because terrestrial higher plant debris had  
58  
59  
60

1  
2  
3  
4  
5 396 already been submitted to strong biotic as well as abiotic degradation under oxic conditions  
6  
7 397 in soils (Vandenbroucke & Largeau 2007); iv) degradation during mobilisation and transport  
8  
9 398 is also likely to be limited due to the very short sediment transport distances within the small  
10  
11 399 studied catchment. The conservative behaviour of TOC was also corroborated by previous  
12  
13 400 studies conducted in recent North African reservoirs. Albergel et al. (2006) demonstrated for  
14  
15 401 instance that the majority of the organic matter found at two Tunisian reservoirs (El Gouazine  
16  
17 402 and Fidh Ali) originated from upstream soil sources, and that this organic matter was not  
18  
19 403 transformed in the recently accumulated sediment (approximately 10 years old in their  
20  
21 404 studies). Ben Slimane et al. (2013) confirmed the terrestrial origin of TOC in the Kamech  
22  
23 405 reservoir deposits using Rock-Eval analysis and  $\delta^{13}\text{C}$  measurements to demonstrate the  
24  
25 406 very low kinetics of TOC degradation through analysis of the entire sediment deposit  
26  
27 407 sequence.

28  
29  
30 408 Previous research and the very different TOC contents found in topsoil and subsoil sources  
31  
32 409 (Table 3) support the relevance of using TOC in combination with  $^{137}\text{Cs}$  as potential sediment  
33  
34 410 tracers in the context of this study. It was verified by comparing results from fingerprinting  
35  
36 411 approach using either the combination of  $^{137}\text{Cs}$  and TOC or  $^{137}\text{Cs}$  alone. Figure 3 shows that  
37  
38 412 very similar results are obtained by both methods. As the use of multiple tracers in a mixing  
39  
40 413 model allows for more reliable source apportionment than the use of only one tracer (Walling  
41  
42 414 et al. 1993; Small et al. 2002; Martinez-Carreras et al. 2008) we finally focused our  
43  
44 415 discussion on the fingerprinting results derived from the combined use of  $^{137}\text{Cs}$  and TOC as  
45  
46 416 tracers.

47  
48  
49  
50 417 *On the driving factors of reservoir siltation*

51  
52 418 Although Fidh Ali catchment is characterized by the lowest mean rainfall amount and the  
53  
54 419 lowest rainfall erosivity index values, it clearly showed the highest erosion contributions  
55  
56 420 whatever the erosion processes considered (Figure 2). The extreme intensity of erosion in  
57  
58 421 Fidh Ali catchment can be explained by its soil composition that contains a very high  
59  
60

1  
2  
3  
4  
5 422 percentage of marly gypsum showing intense swelling/shrinkage processes (clayey soils).  
6  
7 423 They are known to increase soil sensitivity to erosion (Cantón et al., 2001; Sfar Felfoul et al.,  
8  
9 424 1996, 1999) as they facilitate both aggregate dispersion and gully initiation. In this catchment,  
10  
11 425 this specific lithological feature directly appears as the first-order factor driving the very high  
12  
13 426 erosion rates. Areas covered with a large marly gypsum surface should therefore be  
14  
15 427 considered as priority zones for implementing erosion protection measures in the Tunisian  
16  
17 428 Ridge.

18  
19  
20 429 A simple statistical analysis by means of linear regressions was performed in order to  
21  
22 430 investigate the relationships between dominant erosion processes contributing to reservoir  
23  
24 431 siltation and the following characteristics: total catchment area, mean annual rainfall, mean  
25  
26 432 annual erosivity index, mean annual runoff, mean annual runoff coefficient, global slope  
27  
28 433 index, drainage density, percentage of catchment surface covered by (i) soft lithological  
29  
30 434 substrate types, (ii) badlands, (iii) cropland, (iv) equipped with soil conservation measures.  
31  
32 435 Table 4 summarizes the correlation coefficient values obtained for each linear relationship  
33  
34 436 tested when considering the 5 studied catchments. The significant correlations at the 0.05  
35  
36 437 level are mentioned when considering the 5 catchments but also when excluding Fidh Ali as  
37  
38 438 its extreme erosion values may greatly affect some correlations. As this analysis is restricted  
39  
40 439 to a set of 4 or 5 catchments, its statistical significance is relatively low. Consequently only  
41  
42 440 correlation values significant at the 0.05 level are discussed in the remainder of the text and  
43  
44 441 further investigations will be required to confirm these preliminary findings regarding the  
45  
46 442 identification of first-order factors controlling reservoir siltation.

47  
48  
49 443 First, correlation coefficient values between potential driving factors and the contribution of a  
50  
51 444 single individual erosion process (related to either rill/interrill or gully/channel sediment  
52  
53 445 sources) were generally higher than the ones obtained between driving factors and SSY.  
54  
55 446 This may indicate that the tested factors explain individual erosion process contributions  
56  
57 447 rather than the combined effect of all erosion phenomena. Second, several correlations  
58  
59  
60

1  
2  
3  
4  
5 448 values were significant at the 0.05 level when considering the five studied catchments but  
6  
7 449 non-significant when excluding Fidh Ali. This means that the significance of the correlations  
8  
9 450 is mainly due to the extreme erosion values recorded in Fidh Ali catchment and not to the  
10  
11 451 tested factor. On the contrary, a significant negative correlation was obtained between the  
12  
13 452 percentage of catchment area under cropland and the gully/channel erosion contribution (at  
14  
15 453 the 0.05 level) by both including and excluding the Fidh Ali site. This result can be interpreted  
16  
17 454 as the fact that gully/channel erosion took place in a highly degraded environment unsuitable  
18  
19 455 for cultivation. Finally, the percentage of total catchment area equipped with erosion control  
20  
21 456 measures (% of total managed area) appeared to be the most relevant driving factor of  
22  
23 457 reservoir siltation in the five studied catchments. It was significantly correlated with the  
24  
25 458 relative contribution of individual erosion processes (with or without Fidh Ali) and with  
26  
27 459 rill/interrill erosion rate (when excluding the Fidh Ali catchment), both in a negative way. This  
28  
29 460 confirms that soil protection measures have a significant impact on surface erosion by  
30  
31 461 limiting either soil detachment or sediment transportation from hillslopes to catchment outlet.  
32  
33 462 This is a major finding of this study as this factor is anthropogenic and can therefore be  
34  
35 463 controlled.  
36

37  
38  
39 464

40  
41  
42 465 *Main erosion processes contributing to reservoir siltation*

43  
44 466 Jebari et al. (2010) have proposed to use a rough direct relationship between rainfall  
45  
46 467 characteristics (maximum 15-min duration rainfall intensity) and the respective contribution of  
47  
48 468 rill, interrill and gully erosion processes to sediment siltation in 28 Tunisian small reservoirs.  
49  
50 469 Their results indicated that rill/interrill erosion was largely dominant in the major part of the  
51  
52 470 Tunisian Ridge as gully erosion contribution exceeded 20% in only 5 of the 28 studied  
53  
54 471 catchments, and 50% in one single catchment. Considering the same sites of investigation  
55  
56 472 used in this study, they found a dominant contribution of rill/interrill erosion with values  
57  
58 473 between 82% and 90% for the five catchments. There is therefore a contradiction between  
59  
60

1  
2  
3  
4  
5 474 the findings provided by Jebari et al. (2010) and our results as we found that two of the five  
6  
7 475 investigated catchments were in fact dominated by gully/channel erosion processes. This  
8  
9 476 inconsistency may arise from the fact that rainfall characteristics are important but not  
10  
11 477 sufficient to explain the dominant erosion processes in a catchment and that we need to take  
12  
13 478 into account human activities such as soil conservation measures as major driving factors to  
14  
15 479 understand and characterize properly the erosive processes in a catchment.  
16  
17

18 480 To a wider extent, the predominance of gully/channel erosion contribution in two of the five  
19  
20 481 studied catchments confirms the significant contribution of subsuperficial erosion processes  
21  
22 482 (other than sheet and rill erosion) to catchment sediment yield in the Mediterranean zone as  
23  
24 483 recently underlined by Vanmaercke et al. (2012) who compared sheet/rill erosion rates and  
25  
26 484 sediment yield in a large number (n = 1794) of European catchments. This result is likely  
27  
28 485 valid in other regions of the world as for the Alpine zone (Vanmaercke et al., 2012) or in  
29  
30 486 Northern Ethiopia (Haregeweyn et al., 2013).  
31  
32

33 487  
34  
35

#### 36 488 *Implications for catchment management*

37

38 489 A simple sediment source apportionment method conducted at the catchment scale showed  
39  
40 490 that the variability observed for individual (i.e., rill/interrill or gully/channel) erosion process  
41  
42 491 magnitude was two or three times higher than the variability observed for total erosion. The  
43  
44 492 explicit consideration of different erosion processes provided a more efficient way to outline  
45  
46 493 the high variability of erosion processes between catchments than the simple calculation of  
47  
48 494 specific sediment yields (SSY). Moreover, the ranking of catchments following an order of  
49  
50 495 increasing/decreasing erosion contributions proved to be significantly modified depending on  
51  
52 496 the type of erosion processes considered. Identification and quantification of the dominant  
53  
54 497 sediment sources is crucial for our understanding of human impacts on catchment SY and  
55  
56 498 for the design of efficient management strategies to reduce SY at the catchment scale  
57  
58  
59  
60

1  
2  
3  
4  
5 499 (Vanmaercke et al., 2012). In Tunisia, implementation of conservation farming practices  
6  
7 500 should be encouraged in catchments characterised by similar features as Kamech, Fidh Ali  
8  
9 501 and El Melah sites. In these environments, maintaining a minimal vegetation cover of the soil  
10  
11 502 during autumn may provide a good solution to protect it from sheet and rill erosion (Crosaz,  
12  
13 503 1995; Maetens et al., 2012; Menashe, 1998; Rey & Berger, 2002). In contrast, measures  
14  
15 504 dealing with gully and channel erosion should be targeted on sites similar to El Hnach and  
16  
17 505 Sbaihia catchments. To a wider extent, installation of any erosion protection measure must  
18  
19 506 be carefully designed as it is well-known that complex interactions exist between these  
20  
21 507 different types of erosion processes at the catchment scale (Bryan, 2000; de Figueiredo &  
22  
23 508 Fonseca, 1997; Romkens et al., 2001; Roose et al., 2010). In our study, we showed that the  
24  
25 509 use of conservation measure was efficient in decreasing both relative (in %) and absolute (in  
26  
27 510  $\text{Mg ha}^{-1} \text{ yr}^{-1}$ ) rill/interrill erosion contributions to reservoir siltation. These measures that were  
28  
29 511 targeted to reduce rill/interrill erosion therefore fulfilled their objective. However, the  
30  
31 512 significant correlation between the percentage of total managed area and the relative  
32  
33 513 contributions of individual erosion processes (including or excluding Fidh Ali) also means that  
34  
35 514 the implementation of soil conservation measures has a negative effect on the contribution of  
36  
37 515 gully/channel erosion processes. This result suggests that measures aimed to control topsoil  
38  
39 516 erosion must be implemented in association with complementary measures aimed to gully  
40  
41 517 stabilization. Such a combination is especially required when topsoil conservation measures  
42  
43 518 aim to reduce sediment detachment or to enhance sediment trapping without decreasing  
44  
45 519 runoff as it is the case for contour stone bunds for instance. This confirms the need for  
46  
47 520 combining several types of measures (vegetative and structural) to increase sediment  
48  
49 521 trapping and protect reservoir from siltation (Mekonnen et al., 2014).  
50  
51  
52  
53 522 Identification and quantification of the dominant sediment sources are also crucial to identify  
54  
55 523 the most appropriate models to simulate erosion at the catchment scale (De Vente et al.,  
56  
57 524 2013; Morgan, 2011; Vanmaercke et al., 2012) and avoid their misapplication (Govers, 2011).  
58  
59 525 Poor modelling results are frequently obtained when a single model is applied to a wide  
60

1  
2  
3  
4  
5 526 series of catchments, as most models are generally designed for describing a limited number  
6  
7 527 of processes (either sheet and rill erosion or gully and bank erosion) but they are rarely  
8  
9 528 designed to simulate the complete range of phenomena (De Vente & Poesen, 2005; De  
10  
11 529 Vente et al., 2013, Haregeweyn et al., 2013). In Mediterranean environments, the occurrence  
12  
13 530 of severe catchment sediment yield was explained by a complex combination of rill/interrill  
14  
15 531 and gully/channel erosion processes. In the light of these results, we suggest that  
16  
17 532 interactions between both processes need to be further investigated and integrated into soil  
18  
19 533 erosion models.  
20  
21  
22 534  
23  
24  
25 535

## CONCLUSIONS

26  
27 536 This work confirmed the relevance of using a fingerprinting method based on the  
28  
29 537 measurement of  $^{137}\text{Cs}$  and TOC to estimate the relative contribution of rill/interrill vs.  
30  
31 538 gully/channel erosion processes to reservoir siltation in Northern Africa catchments. Among  
32  
33 539 the five investigated catchments, three of them were characterized by dominant rill/interrill  
34  
35 540 erosion contribution, whereas gully/channel erosion contribution prevailed in the two other  
36  
37 541 sites. The results also showed that a very large variability of erosion processes contributed to  
38  
39 542 reservoir siltation in catchments located along the Tunisian Ridge even when similar  
40  
41 543 catchment specific sediment yields were recorded. The presence of a large surface cover of  
42  
43 544 marly gypsum material was confirmed to enhance both rill/interrill and gully/channel erosion  
44  
45 545 processes. Furthermore, the implementation of soil protection measures was also identified  
46  
47 546 as a main factor driving erosion in this region. This work also corroborated the existence of  
48  
49 547 complex interactions between rill/interrill and gully processes. Consequently, implementation  
50  
51 548 of soil conservation measures must be carefully designed at the catchment scale. Indeed soil  
52  
53 549 conservation measures aimed to prevent particle detachment on cultivated hillslopes and  
54  
55 550 their downstream transportation seemed to have negative feedback by amplifying  
56  
57 551 downstream gully/channel erosion. In catchments where rill/interrill erosion contribution  
58  
59  
60

1  
2  
3  
4  
5 552 dominates, we therefore suggest to implement topsoil protection measures that significantly  
6  
7 553 reduce sediment concentration and runoff at the same time. To a wider extent, this work also  
8  
9 554 corroborated the interest to combine different types of conservation measures (vegetative  
10  
11 555 and structural, in both the fields and gullies/channels) to limit reservoir siltation.  
12

13  
14 556 Overall, our results also suggest the need to develop erosion models simulating interactions  
15  
16 557 between rill/interrill and gully/channel processes in order to guide the implementation of  
17  
18 558 erosion control measures in Mediterranean catchments.  
19

20  
21 559  
22

## 23 24 560 ACKNOWLEDGEMENTS

25  
26  
27 561 This study was financially supported by the IRD-DSF, SCAC of French embassy and a CNRS/DGRS  
28  
29 562 exchange agreement (No. 24443) between France and Tunisia. It was undertaken in the framework of  
30  
31 563 the OMERE Observatory funded by INRA and IRD. The authors would like to acknowledge the field  
32  
33 564 staff -and especially Radhouane Hamdi- for the help in data collection. We are also indebted to the  
34  
35 565 Tunisian Ministry of Agriculture (DG-ACTA) that participated to the hydrological and sediment  
36  
37 566 measurements in the five studied catchments.  
38

39 567  
40

## 41 42 568 REFERENCES

43  
44 569 Albergel J, Boufaroua M, Pepin Y. 1998. Bilan de l'érosion sur les petits bassins versants des lacs  
45  
46 570 collinaires en climat semi-aride Tunisien. In : L'eau et la fertilité des sols : deux ressources à gérer  
47  
48 571 ensemble. *Bulletin - Réseau Erosion* **18**: 67-75.

49  
50  
51 572 Albergel J, Collinet J, Pépin Y, Nasri S, Boufaroua M, Droubi A, Merzouk A. 2005. Sediment budgets  
52  
53 573 on hill reservoirs of small catchments in North Africa and the Middle East. *IASH Pub.* **291 (1)**: 323–  
54  
55 574 331.

56  
57 575 Albergel J, Mansouri T, Zante P, Ben Mamou A, Abdeljaoued S. 2006. Organic carbon in the  
58  
59  
60



- 1  
2  
3  
4  
5 576 sediments of hill dams in a semiarid Mediterranean area. In: Roose E, Lal R, Feller C, Barthès B,  
6  
7 577 Stewart BA (eds) Soil erosion and carbon dynamics. Taylor and Francis, Boca Raton, pp 289–299.  
8  
9  
10 578 Annabi M, Bahri H, Latiri K. 2009. Statut organique et respiration microbienne des sols du nord de la  
11  
12 579 Tunisie. *Biotechnologie Agronomie Société et Environnement* **13(3)**: 401-408.  
13  
14 580 Ayadi I, Abida H, Djebbar Y, Mahjoub R. 2010. Sediment yield variability in central Tunisia: a  
15  
16 581 quantitative analysis of its controlling factors. *Hydrological Sciences Journal* **55(3)**: 446- 458.  
17  
18  
19 582 Ben Mammou A, Louati MH. 2007. Évolution temporelle de l'envasement des retenues de barrages de  
20  
21 583 Tunisie. *Revue des sciences de l'eau / Journal of Water Science* **20(2)**: 201-210. DOI:  
22  
23 584 10.7202/015813ar.  
24  
25 585 Ben Slimane A, Raclot D, Evrard O, Sanaa M, Lefèvre I, Ahmadi M, Tounsi M, Rumpel C, Ben  
26  
27 586 Mammou A, Le Bissonnais Y. 2013. Fingerprinting sediment sources in the outlet reservoir of a  
28  
29 587 hilly cultivated catchment of Tunisia. *Journal of Soils and Sediments* **13**: 801-815. DOI:  
30  
31 588 10.1007/s11368-012-0642-6.  
32  
33 589 Bryan RB. 2000. Soil erodibility and processes of water erosion on hillslope. *Geomorphology* **32**: 385-  
34  
35 590 415.  
36  
37  
38 591 Cantón Y, Solé-Benet A, Queralt I, Pini R. 2001. Weathering of a gypsum-calcareous mudstone under  
39  
40 592 semi-arid environment at Tabernas, SE Spain: laboratory and field-based experimental approaches.  
41  
42 593 *Catena* **44**: 111-132.  
43  
44 594 Cantón Y, Solé-Benet A, de Vente J, Boix-Fayos C, Calvo-Cases A, Asensio C, Puigdefábregas J.  
45  
46 595 2011. A review of runoff generation and soil erosion across scales in semiarid south-eastern Spain.  
47  
48 596 *J Arid Environ* **75(12)**: 1254-1261.  
49  
50 597 Cerdà A, Giménez-Morera A, Bodí MB. 2009. Soil and water losses from new citrus orchards growing  
51  
52 598 on sloped soils in the western Mediterranean basin. *Earth Surface Processes and Landforms* **34**,  
53  
54 599 1822-1830. DOI: 10.1002/esp.1  
55  
56  
57 600 Cerdà A, Lavee H, Romero-Díaz A, Hooke J, Montanarella L. 2010. Preface of the special Issue on  
58  
59 601 Soil Erosion and Degradation on Mediterranean Type Ecosystems. *Land Degradation &*  
60

- 1  
2  
3  
4  
5 602 *Development* **21**: 71-74.  
6  
7  
8 603 Cerdan O, Govers G, Le Bissonnais Y, Van Oost K, Poesen J, Saby N, Gobin A, Vacca A, Quinton J,  
9  
10 604 Auerswald K, Klik A, Kwaad FJPM, Raclot D, Ionita I, Rejman J, Rousseva S, Muxart T, Roxo MJ,  
11  
12 605 Dostal T. 2010. Rates and spatial variations of soil erosion in Europe: A study based on erosion  
13  
14 606 plot data. *Geomorphology* **122**: 167–177.
- 15  
16 607 CNEA (Centre National des Etudes Agricoles). 2007. Elaboration d'une étude sur l'état de  
17  
18 608 désertification pour une gestion durable des Ressources Naturelles. Centre National des Etudes  
19  
20 609 Agricoles, Tunisie. (in French, available at  
21  
22 610 [http://www.environnement.nat.tn/envir/sid/dmdocuments/desertification\\_tunisie/rapport\\_cnea\\_esd.](http://www.environnement.nat.tn/envir/sid/dmdocuments/desertification_tunisie/rapport_cnea_esd.pdf)  
23  
24 611 pdf)
- 25  
26 612 Collinet J, Zante P. 2005. Analyse du ravinement de bassins versants à retenues collinaires sur sols à  
27  
28 613 fortes dynamiques structurales (Tunisie). *Géomorphologie : relief, processus, environnement* **1**: 61-  
29  
30 614 74.
- 31  
32 615 Collins AL, Walling DE. 2002. Selecting fingerprint properties for discriminating potential suspended  
33  
34 616 sediment sources in river basins. *Journal of Hydrology* **261**: 218-244.
- 35  
36  
37 617 Collins AL, Walling DE, Leeks GJL. 1998. Fingerprinting the origin of fluvial suspended sediment in  
38  
39 618 larger river basins: combining assessment of spatial provenance and source type. *Geografiska*  
40  
41 619 *Annaler: Series A, Physical Geography* **79**: 239-254. DOI: 10.1111/j.0435-3676.1997.00020.x
- 42  
43 620 Collins AL, Walling DE, Webb L, King P. 2010. Apportioning catchment scale sediment sources using  
44  
45 621 a modified composite fingerprinting technique incorporating property weightings and prior  
46  
47 622 information. *Geoderma* **155**: 249-261.
- 48  
49 623 Crosaz Y. 1995. Le matériel végétal: un outil pour la protection des sols. *Bulletin - Réseau Erosion* **15**:  
50  
51 624 449-460.
- 52  
53  
54 625 Damnati B, Ibrahimi S, Radakovitch O. 2012. Quantifying Erosion Using <sup>137</sup>Cs And <sup>210</sup>Pb in  
55  
56 626 Cultivated Soils In Three Mediterranean Watershed: Synthesis Study From El Hachef, Raouz And  
57  
58 627 Nakhla (North West Morocco). *Journal of African Earth Sciences* **79**: 50-57. DOI:  
59  
60

- 1  
2  
3  
4  
5 628 <http://dx.doi.org/10.1016/j.jafrearsci.2012.10.006>.
- 6  
7  
8 629 De Figueiredo T, Fonseca F. 1997. Les sols, les processus d'érosion et l'utilisation de la terre en  
9  
10 630 montagne au Nord-Est du Portugal: Approche cartographique sur quelques zones du Parc Naturel  
11  
12 631 de Montesinho. *Bulletin - Réseau Erosion* **17**: 205–217.
- 13  
14 632 De Vente J, Poesen J. 2005. Predicting soil erosion and sediment yield at the basin scale: Scale  
15  
16 633 issues and semi-quantitative models. *Earth-Science Reviews* **71**: 95–125.
- 17  
18  
19 634 De Vente J, Poesen J, Bazzofi P, Van Rompaey A, Verstaeten G. 2006. Predicting catchment  
20  
21 635 sediment yield in Mediterranean environments: the importance of sediment sources and  
22  
23 636 connectivity in Italian drainage basins. *Earth Surf Proc Land* **31**: 1017-1034.
- 24  
25 637 De Vente J, Poesen J, Verstraeten G, Van Rompaey A, Govers G. 2008. Spatially distributed  
26  
27 638 modelling of soil erosion and sediment yield at regional scales in Spain. *Global and Planetary*  
28  
29 639 *Change* **60**: 393-415.
- 30  
31 640 De Vente J, Poesen J, Verstraeten G, Govers G, Vanmaercke M., Van Rompaey A, Arabkhedri M.,  
32  
33 641 Boix-Fayos C. 2013. Predicting soil erosion and sediment yield at regional scales: Where do we  
34  
35 642 stand? *Earth-Science Reviews* **127**: 16-29.
- 36  
37  
38 643 Devereux OH, Prestegard KL, Needelman BA, Gellis AC. 2010. Suspended sediment sources in an  
39  
40 644 urban watershed, Northeast Branch Anacostia River, Maryland. *Hydrological Processes* **24**: 1391–  
41  
42 645 1403.
- 43  
44 646 Evrard O, Némery J, Gratiot N, Duvert C, Ayrault S, Lefèvre I, Poulénard J, Prat C, Bonté P, Esteves  
45  
46 647 M. 2010. Sediment dynamics during the rainy season in tropical highland catchments of central  
47  
48 648 Mexico using fallout radionuclides. *Geomorphology* **124**: 42–54.
- 49  
50 649 Evrard O, Navratil O, Ayrault S, Ahmadi M, Némery J, Legout C, Lefèvre I, Poirel A, Bonté P, Esteves  
51  
52 650 M. 2011. Combining suspended sediment monitoring and fingerprinting to determine the spatial  
53  
54 651 origin of fine sediment in a mountainous river catchment. *Earth Surf Proc Land* **36**: 1072–1089.
- 55  
56  
57 652 Faleh A, Navas A, Sadiki A. 2005. Erosion and dam siltation in a Rif catchment (Morocco). *IAHS-AISH*  
58  
59 653 *Publication* **292**: 58-64.
- 60

- 1  
2  
3  
4  
5 654 García-Ruiz JM, Nadal-Romero E, Lana-Renault N, Beguería S. 2013. Erosion in Mediterranean  
6  
7 655 landscapes: changes and future challenges. *Geomorphology* **198**: 20-36.  
8  
9 656 DOI:10.1016/j.geomorph.2013.05.023  
10  
11 657 Govers G. 2011. Misapplications and Misconceptions of Erosion Models, in: Morgan, R.P.C., Nearing,  
12  
13 658 M.A. (Eds.), *Handbook of Erosion Modelling*. John Wiley & Sons, Ltd, Chichester, UK, 117-134.  
14  
15 659 DOI:10.1002/9781444328455.ch7.  
16  
17  
18 660 Guzmán G, Quinton JN, Nearing MA, Mabit L, Gómez JA. 2013. Sediment tracers in water erosion  
19  
20 661 studies: Current approaches and challenges. *J Soils Sediments* **13**: 816–833.  
21  
22 662 Habi M, Morsli B. 2011. Contraintes et perspectives des retenues collinaires dans le Nord-Ouest  
23  
24 663 algérien. *Sécheresse* **22**: 49-56. DOI : 10.1684/sec.2011.0293.  
25  
26  
27 664 Haddadchi A, Ryder DS, Evrard O, Olley J. 2013. Sediment fingerprinting in fluvial systems: Review of  
28  
29 665 tracers, sediment sources and mixing models. *International Journal of Sediment Research* **28**: 560-  
30  
31 666 578.  
32  
33 667 Haregeweyn N, Poesen J, Verstraeten G, Govers G, de Vente J, Nyssen J, Deckers J, Moeyersons J.  
34  
35 668 2013. Assessing the performance of a spatially distributed soil erosion and sediment delivery  
36  
37 669 model (WATEM/SEDEM) in Northern Ethiopia. *Land Degradation & Development* **24**: 188- 204.  
38  
39 670 DOI 10.1002/ldr.1121.  
40  
41 671 He Q, Walling DE. 1996. Interpreting particle size effects in the adsorption of <sup>137</sup>Cs and unsupported  
42  
43 672 <sup>210</sup>Pb by mineral soils and sediments. *Journal Environ Radioactivity* **30**: 117-137.  
44  
45  
46 673 Hentati A, Kawamura A, Amaguchi H, Iseri Y. 2010. Evaluation of sedimentation vulnerability at small  
47  
48 674 hillside reservoirs in the semi-arid region of Tunisia using the Self-Organizing Map. *Geomorphology*  
49  
50 675 **122**: 56-64.  
51  
52 676 Jebari S, Berndtsson R, Bahri A, Boufaroua M. 2010. Spatial soil loss risk and reservoir siltation in  
53  
54 677 semi-arid Tunisia. *Hydrolog. Sci. J.* **55(1)**: 121-137.  
55  
56  
57 678 Jones RJA, Hiederer R, Rusco E, Loveland PJ, Montanarella L. 2004. The map of organic carbon in  
58  
59 679 topsoils in Europe, Version 1.2, September 2003. Office for Official Publications of the European  
60

- 1  
2  
3  
4  
5 680 Communities, Luxembourg.  
6  
7  
8 681 Juracek KE, Ziegler AC. 2009. Estimation of sediment sources using selected chemical tracers in the  
9  
10 682 Perry lake basin, Kansas, USA. *International Journal of Sediment Research* **24(1)**: 108-125.  
11  
12 683 Koiter AJ, Owens PN, Petticrew EL, Lobb DA. 2013. The behavioural characteristics of sediment  
13  
14 684 properties and their implications for sediment fingerprinting as an approach for identifying sediment  
15  
16 685 sources in river basins. *Earth-Science Reviews* **125**: 24–42.  
17  
18 686 Lahlou A. 2000. Quelques aspects environnementaux dans les pays du Maghreb, ed. ISESCO, Rabat,  
19  
20 687 Maroc.  
21  
22  
23 688 Lal R. 2001. Soil degradation by erosion. *Land Degradation & Development* **12**: 519-539. DOI:  
24  
25 689 10.1002/ldr.472  
26  
27  
28 690 Laudicina VA, Novara A, Barbera V, Egli M, Badalucco L. 2014. Long-term tillage and cropping  
29  
30 691 system effects on chemical and biochemical characteristics of soil organic matter in a  
31  
32 692 Mediterranean semiarid environment. *Land Degradation & Development*. DOI: 10.1002/ldr.2293  
33  
34 693 Maetens W, Poesen J, Vanmaercke M. 2012. How effective are soil conservation techniques in  
35  
36 694 reducing plot runoff and soil loss in Europe and the Mediterranean? *Earth-Science Reviews* **115(1-**  
37  
38 695 **2)**: 21-36. DOI:10.1016/j.earscirev.2012.08.003.  
39  
40 696 Mandal D, Sharda VN. 2013. Appraisal of soil erosion risk in the Eastern Himalayan region of India for  
41  
42 697 soil conservation planning. *Land Degradation & Development* **24**: 430–437. DOI: 10.1002/ldr.1139  
43  
44  
45 698 Martinez-Carreras N, Gallart F, Iffly JF, Pfister L, Walling DE, Krein A. 2008. Sediment dynamics in  
46  
47 699 changing environments. In: Schmidt J, Cochrane T, Phillips C, Elliot S, Davies T, Basher L (eds),  
48  
49 700 Uncertainty assessment in suspended sediment fingerprinting based on tracer mixing models: a  
50  
51 701 case study from Luxembourg. IAHS Publ 325, Wallingford, pp 94–105.  
52  
53 702 Meersmans J, Martin MP, Lacarce E, De Baets S, Jolivet C, Boulonne L, Lehmann S, Saby NPA,  
54  
55 703 Bispo A, Arrouays D. 2012. A high resolution map of French soil organic carbon. *Agron. Sustain.*  
56  
57 704 *Dev.* **32**: 841–851. DOI 10.1007/s13593-012-0086-9.  
58  
59  
60

- 1  
2  
3  
4  
5 705 Mekonnen M, Keesstra SD, Stroosnijder L, Baartman JE, Maroulis J. 2014. Soil conservation through  
6 sediment trapping: *A review. Land Degradation & Development*. DOI: 10.1002/ldr.2308  
7  
8  
9  
10 707 Menashe E. 1998. Vegetation and Erosion: A literature Survey. Proceedings of the Native Plants  
11 Symposium, Oregon State University, Forestry Sciences Lab., Corvallis, OR. 130- 135.  
12  
13  
14 709 Morgan RC. 2011. Model Development: A user's perspective, in: Morgan, R.P.C., Nearing, M.A. (Eds.),  
15 Handbook of Erosion Modelling. John Wiley & Sons, Ltd, Chichester, UK, 9-32.  
16  
17 710 DOI:10.1002/9781444328455.ch2.  
18  
19  
20 712 Motha JA, Wallbrink PJ, Hairsine PB, Grayson RB. 2003. Determining the sources of suspended  
21 sediment in a forested catchment in southeastern Australia. *Water Resour Res* **39**(3):1056.  
22  
23 713 DOI10.1029/2001WR000794  
24  
25  
26 715 Mourrier B. 2008. Contribution de l'approche sédimentologique à la reconstitution de l'histoire des sols.  
27 Définition de traceurs pédologiques et application sur des sédiments lacustres de montagne  
28 (Maurienne, Savoie, France). Thèse de doctorat, France.  
29  
30 716  
31  
32  
33 718 Mukundan R, Radcliffe DE, Ritchie JC, Risse LM, McKinley RA. 2010. Sediment Fingerprinting to  
34 Determine the Source of Suspended Sediment in a Southern Piedmont Stream. *J. Environ. Qual.*  
35  
36 719 **39**: 1328–1337.  
37  
38  
39 721 Nagle GN, Fahey TJ, Ritchie JC, Woodbury PB. 2007. Variations in sediment sources and yields in  
40 the Finger Lakes and Catskills regions of New York. *Hydrol. Process.* **21**: 828–838.  
41  
42 722  
43  
44 723 Owens PN, Walling DE, Leeks GJL. 1999. Use of floodplain sediment cores to investigate recent  
45 historical changes in overbank sedimentation rates and sediment sources in the catchment of the  
46 River Ouse, Yorkshire, UK. *Catena* **36**:21–47.  
47  
48 724  
49  
50 726 Patience AJ, Lallier-Vergès E, Sifeddine A, Albéric P, Guillet B. 1995. Organic fluxes and early  
51 diagenesis in the lacustrine environment: the superficial sediments of the Lac du Bouchet (Haute  
52 Loire, France). In: Lallier-Vergès E, Tribouvillard NP, Bertrand P (Eds.), Organic Matter  
53 Accumulation: The Organic Cyclicities of the Kimmeridge Clay Formation (Yorkshire, GB) and the  
54 Recent Maar Sediments (Lac du Bouchet, France). Springer-Verlag, Heidelberg, pp. 145–  
55  
56 729  
57  
58 730  
59  
60

- 1  
2  
3  
4  
5 731 156.Poesen J, Vandaele K, Van Wesemael B. 1996. Contribution of gully erosion to sediment  
6 732 production in cultivated lands and rangelands. *IAHS Publications* **236**: 251– 266.
- 7  
8  
9 733 Poesen J, Nachtergaele J, Verstraeten G, Valentin C. 2003. Gully erosion and environmental change:  
10 734 importance and research needs. *Catena* **50 (2/4)**: 91-133.
- 11  
12  
13  
14 735 Porto P, Walling DE, Capra A. 2014. Using  $^{137}\text{Cs}$  and  $^{210}\text{Pb}$  measurements and conventional  
15 736 surveys to investigate the relative contributions of interrill/rill and gully erosion to soil loss from a  
16 737 small cultivated catchment in Sicily. *Soil & Tillage Research* **135**: 18–27.
- 17  
18  
19  
20 738 Raclot D, Albergel J. 2006. Runoff and water erosion modelling using WEPP on a Mediterranean  
21 739 cultivated catchment. *Phys. Chem. Earth* **31(17)**: 1038-1047.
- 22  
23  
24  
25 740 Raclot D, Le Bissonnais Y, Louchart X, Andrieux P, Moussa R, Voltz M. 2009. Soil tillage and scale  
26 741 effects on erosion from fields to catchment in a Mediterranean vineyard area. *Agriculture,*  
27 742 *Ecosystems and Environment* **134**: 201–210.
- 28  
29  
30  
31 743 Rahaingomanana N. 1998. Caractérisation géochimique des lacs collinaires de la Tunisie semi-aride  
32 744 et régulation géochimique du phosphore. PhD thesis, Univ I, Montpellier.
- 33  
34  
35  
36 745 Rebai H, Raclot D, Ben OueddouH. 2012. Facteurs du ravinement dans la dorsale tunisienne et le cap  
37 746 bon. *Rev. Mar. Sci. Agron. Vét.* **1**: 18-22.
- 38  
39  
40 747 Rey F, Berger F. 2002. Interactions végétation-érosion et génie écologique pour la maîtrise de  
41 748 l'érosion en Montagne. Colloque international « L'eau en montagne : gestion intégrée des hauts  
42 749 bassins versants», Megève, France. <http://www.inbo-news.org/IMG/pdf/Rey-Berger-2.pdf>
- 43  
44  
45  
46 750 Romkens MJM, Helming K, Prasad SN. 2001. Soil erosion under different rainfall intensities, surface  
47 751 roughness, and soil water regimes. *Catena* **46**: 103-123.
- 48  
49  
50  
51 752 Roose E, Chebbani R, Bourougaa L. 2000. Ravinement en Algérie, facteurs de contrôle, quantification  
52 753 et réhabilitation. *Science et changements planétaires / Sécheresse* **11(4)**: 317-326.
- 53  
54  
55  
56 754 Roose E, Sabir M, Laouina A. 2010. Gestion durable des eaux et des sols au Maroc: Valorisation des  
57 755 techniques traditionnelles méditerranéennes, IRD ed., Marseille.

- 1  
2  
3  
4  
5 756 Sfar Felfoul M, Snane MH, Mlaouihi A, Megdiche MF. 1996. Intégration de certains facteurs  
6  
7 757 biophysiques dans l'espace pour l'étude du ravinement dans le sous bassin versant d'Oued Maiz  
8  
9 758 dans la région de Haffouz (Tunisie centrale). *Bulletin - Réseau Erosion* **16**: 457-470.
- 10  
11 759 Sfar Felfoul M, Snane MH, Mlaouihi A, Megdiche MF. 1999. Importance du facteur lithologique sur le  
12  
13 760 développement des ravins du bassin versant d'oued Maiez en Tunisie centrale. *Bull. Eng. Geol.*  
14  
15 761 *Environ.* **57**: 285-293.
- 16  
17  
18 762 Small IF, Rowan JS, Franks SW. 2002. Structure, function and management implications of fluvial  
19  
20 763 sedimentary systems. In: Dyer FJ, Thoms MC, Olley JM (eds), Quantitative sediment fingerprinting  
21  
22 764 using a Bayesian uncertainty estimation framework. IAHS Publ 276, Wallingford, pp 443–450.
- 23  
24 765 Smith HG, Sheridan GJ, Nyman P, Child DP, Lane PNJ, Hotchkis MAC, Jacobsen GE. 2012.  
25  
26 766 Quantifying sources of fine sediment supplied to post-fire debris flows using fallout radionuclide  
27  
28 767 tracers. *Geomorphology* **139-140**:403-415.
- 29  
30 768 Temple-Boyer E, Richard JF, Arnould P. 2007. Segmenter les paysages de l'eau : une méthode pour  
31  
32 769 l'interprétation hydrodynamique des paysages (Dorsale tunisienne). *Sécheresse* **18(3)**: 149-60.
- 33  
34  
35 770 Vandembroucke M, Largeau C. 2007. Kerogen origin, evolution and structure. *Organic Geochemistry*  
36  
37 771 **38**:719–833.
- 38  
39 772 Vanmaercke M, Poesen J, Verstraeten G, de Vente J, Ocakoglu F. 2011. Sediment yield in Europe:  
40  
41 773 Spatial patterns and scale dependency. *Geomorphology* **130**: 142-161.
- 42  
43  
44 774 Vanmaercke M, Maetens W, Poesen J, Jankauskas B, Jankauskiene G, Verstraeten G, de Vente J.  
45  
46 775 2012. A comparison of measured catchment sediment yields with measured and predicted hillslope  
47  
48 776 erosion rates in Europe. *Journal of Soils and Sediments* **12**: 586-602.
- 49  
50 777 Verheijen FGA, Jones RJA, Rickson RJ, Smith CJ. 2009. Tolerable versus actual soil erosion rates in  
51  
52 778 Europe. *Earth Science Reviews* **94**(1-4): 23-38. DOI: 10.1016/j.earscirev.2009.02.003.
- 53  
54  
55 779 Walling DE, Woodward JC, Nicholas AP. 1993. Tracers in hydrology. In: Peters NE, Hoehn E,  
56  
57 780 Leibundgut C, Tase N, Walling DE (eds), A multi-parameter approach to fingerprint suspended-  
58  
59 781 sediment sources. IAHS Publ 215, Wallingford, pp 329–338.



- 1  
2  
3  
4  
5 782 Walling DE. 2005. Tracing suspended sediment sources in catchments and river systems. *Sci Total*  
6  
7 783 *Environ* **344**: 159-184.  
8  
9  
10 784 Walling DE. 2013. The evolution of sediment source fingerprinting investigations in fluvial systems. *J.*  
11  
12 785 *Soils Sediments* **13**: 1658–1675.  
13  
14 786 Wasson RJ, Furlonger L, Parry D, Pietsch T, Valentine E, Williams D. 2010. Sediment sources and  
15  
16 787 channel dynamics, Daly River, Northern Australia. *Geomorphology* **114**: 161-174.  
17  
18 788 Wilkinson SN, Wallbrink PJ, Hancock GJ, Blake WH, Shakesby RA, Doerr SH. 2009. Fallout  
19  
20 789 radionuclide tracers identify a switch in sediment sources and transport-limited sediment yield  
21  
22 790 following wildfire in a eucalypt forest. *Geomorphology* **110**: 140-151.  
23  
24  
25 791 Wischmeier WH, Smith DD. 1958. Rainfall energy and its relation to soil loss. *Trans. Am. Geophys.*  
26  
27 792 *Union* **39**: 285-291.  
28  
29 793 Woodward JC. 1995. Patterns of erosion and suspended sediment yield in Mediterranean river basins,  
30  
31 794 in: Foster, I. D. L., Gurnell, A. M., Webb, B.W., (Eds.), *Sediment and Water Quality in River*  
32  
33 795 *Catchments*. Wiley, Chichester, 365–389.  
34  
35  
36 796 Zhang X, Walling DE. 2005. Landscape and watershed processes: characterizing land surface erosion  
37  
38 797 from cesium-137 profiles in lake and reservoir sediments. *J Environ Qual* **34**:514–52.  
39  
40 798 Zhao G, Mu X, Wen Z, Wang F, Gao P. 2013. Soil erosion, conservation, and eco-environment  
41  
42 799 changes in the loess plateau of China. *Land Degrad. Dev.* **24**: 499–510. DOI: 10.1002/ldr.2246  
43  
44  
45 800

1  
2  
3  
4  
5 801 **Figure legends**  
6  
7

8 802  
9  
10

11 803 Figure 1. Location of the selected catchments within a pilot network of monitored catchment  
12 804 reservoirs along the Tunisian Ridge and the Cape Bon region (modified from Temple-Boyer  
13 805 et al., 2007). Rainfall information (isohyet) has been superimposed.  
14  
15

16  
17 806 Figure 2. Catchment sediment yield, rill/interrill and gully/channel erosion contribution to  
18 807 reservoir siltation for the five study catchments.  
19  
20

21  
22 808 Figure 3. Sediment source apportionments for the five studied catchments using the mixing  
23 809 model either with the combination of  $^{137}\text{Cs}$  and TOC or with  $^{137}\text{Cs}$  only.  
24  
25

26  
27 810  
28  
29

30 811  
31  
32

33 812  
34  
35

36 813  
37  
38

39 814  
40  
41

42 815  
43  
44

45 816  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

817 Table 1. Characteristics of the five studied catchments as of 2010. The bioclimatic zones were extracted from the bioclimatic map of Tunisia  
 818 (CNEA, 2007). The global slope index ( $m\ km^{-1}$ ) is defined by the ratio between altitude difference (m) of approximately 5% and 95% of the  
 819 catchment surface and the length of the equivalent rectangle (km).

820

| Catchments      | Description |                    |   |                  | Characteristics / Factors      |                      |  |                    |                       |  |   |                      |                                 |
|-----------------|-------------|--------------------|---|------------------|--------------------------------|----------------------|--|--------------------|-----------------------|--|---|----------------------|---------------------------------|
|                 | Name        | Area<br>( $km^2$ ) | Drainage<br>density<br>( $m\ ha^{-1}$ ) | Reservoir        |                                | Regional climate     |  | Lithology          | Topography            |  | Land cover  | Catchment management |                                 |
|                 |             |                    |   | Building<br>year | Initial<br>volume<br>( $m^3$ ) | Bioclimatic<br>zones | Mean<br>annual<br>rainfall<br>( $mm$ ) |                    | Altitude<br>range (m) | Global slope<br>index<br>( $m\ km^{-1}$ )                                    |   | Type of management   | Total<br>managed<br>area<br>(%) |
| <b>El Hnach</b> | 3.67        | 97.3               | 1992                                    | 77 400           | Higher Semi<br>arid            | 436                  | 25% limestone,<br>75% Marls            | Max:745<br>Min:500 | 104                   | 47% cropland,<br>7% limestone<br>outcrops,<br>7% badlands,<br>39% scrubland. | contour bench terraces +<br>small stone check-dams+<br>small contour stone<br>bunds + tree planting | 32                   |                                 |
| <b>El Melah</b> | 0.61        | 48.2               | 1991                                    | 20 000           | Higher Semi<br>arid            | 450                  | 50% Marls 50%<br>sandstone             | Max:190<br>Min:90  | 36                    | 60% cropland,<br>40% scrubland.  | Without management  | 0                    |                                 |
| <b>Fid Ali</b>  | 2.38        | 158.3              | 1991                                    | 134 700          | Lower semi-<br>arid            | 285                  | 85% gypsum<br>marl, 15%<br>limestone   | Max:460<br>Min:360 | 38                    | 35% cropland,<br>53% scrubland,<br>12% badlands.                             | Small weir dry stone  | 6                    |                                 |
| <b>Kamech</b>   | 2.63        | 79                 | 1994                                    | 142 500          | Sub-humid                      | 650                  | 80% Marls, 20%<br>sandstone            | Max:203<br>Min:95  | 40                    | 70% cropland,<br>30% scrubland.  | Without management  | 0                    |                                 |
| <b>Sbaihia</b>  | 3.57        | 53.3               | 1993                                    | 135 500          | Higher Semi<br>arid            | 450                  | 70% Marls, 30%<br>limestone            | Max:480<br>Min:200 | 77                    | 50% cropland,<br>47% forest,<br>3% badlands.                                 | contour bench terraces  | 41.3                 |                                 |

821 Table 2. Mean annual hydrologic and sediment yield measurements in the five studied  
822 catchments.

823

| Catchment | Periods            | Area-specific sediment yield (SSY) (Mg ha <sup>-1</sup> yr <sup>-1</sup> ) | Annual rainfall (mm yr <sup>-1</sup> ) | Erosivity index (MJ mm ha <sup>-1</sup> h yr <sup>-1</sup> ) | Runoff (mm yr <sup>-1</sup> ) | Runoff coefficient (%) |
|-----------|--------------------|--|--|--|-------------------------------|------------------------|
| El Hnach* | 09/1992 – 02/2007* | 15   | 372                                    | 525  | 28                            | 7.5                    |
| El Melah  | 10/1995 – 08/1999  | 17   | 578                                    | 1201   | 38                            | 6.6                    |
| Fidh Ali  | 06/1993 – 09/1999  | 38   | 279                                    | 469  | 18                            | 6.5                    |
| Kamech    | 09/1994 – 11/2008  | 15   | 561                                    | 1063   | 114                           | 20.3                   |
| Sbailia   | 09/1993 – 11/2006  | 10   | 454                                    | 798  | 72                            | 15.7                   |

824

825 \* Rainfall and runoff measurements for El Hnach catchment were only available for the period between 09/1994  
826 and 08/1999 because of technical failures.

827

828

829 Table 3. Mean and Standard deviation of potential fingerprinting properties and specific  
 830 surface area (SSA) measured in sediment source and reservoir sediment samples (n=85)  
 831 collected in the five studied catchments.

|                 | Locations                 | Number of samples /cores | $^{137}\text{Cs}$ ( $\text{Bq kg}^{-1}$ ) | TOC (%)        | SSA ( $\text{m}^{-1}$ ) |
|-----------------|---------------------------|--------------------------|---|----------------|-------------------------|
| <b>El Hnach</b> | Topsoil                   | 6                        | $7.8 \pm 10.3$                            | $1.2 \pm 0.4$  | $25823 \pm 5360$        |
|                 | Gully/channel bank        | 5                        | $0 \pm 0.1$                               | $0.5 \pm 0.1$  | $24253 \pm 2945$        |
|                 | Upstream reservoir area   | 2                        | $1.4 \pm 0.1$                             | $0.7 \pm 0.3$  | $17035 \pm 846$         |
|                 | Downstream reservoir area | 1                        | 2.3                                       | 0.8            | 29329                   |
| <b>El Melah</b> | Topsoil                   | 7                        | $3.9 \pm 2.5$                             | $0.7 \pm 0.2$  | $13083 \pm 6783$        |
|                 | Gully/channel bank        | 4                        | $0 \pm 0.0$                               | $0.4 \pm 0.1$  | $13925 \pm 8076$        |
|                 | Upstream reservoir area   | 3                        | $4.9 \pm 2.7$                             | $1.5 \pm 0.1$  | $18982 \pm 6362$        |
|                 | Downstream reservoir area | 1                        | 5.7                                       | 1.17           | 23876                   |
| <b>Fidh Ali</b> | Topsoil                   | 6                        | $3.5 \pm 2.8$                             | $0.5 \pm 0.1$  | $12065 \pm 3513$        |
|                 | Gully/channel bank        | 4                        | $0.1 \pm 0.2$                             | $0.3 \pm 0.04$ | $15290 \pm 3458$        |
|                 | Upstream reservoir area   | 1                        | 2.1                                       | 0.7            | 11443                   |
|                 | Downstream reservoir area | 1                        | 3.1                                       | 0.5            | 14880                   |
| <b>Kamech</b>   | Topsoil                   | 8                        | $3.7 \pm 1.5$                             | $1.1 \pm 0.1$  | $29043 \pm 3921$        |
|                 | Gully/channel bank        | 9                        | $0.2 \pm 0.3$                             | $0.5 \pm 0.1$  | $23540 \pm 2146$        |
|                 | Upstream reservoir area   | 8                        | $1.4 \pm 0.4$                             | $0.7 \pm 0.2$  | $15695 \pm 2607$        |
|                 | Downstream reservoir area | 5                        | $3.1 \pm 0.6$                             | $1.0 \pm 0.1$  | $26253 \pm 1725$        |
| <b>Sbaihia</b>  | Topsoil                   | 6                        | $10.1 \pm 4.2$                            | $2.5 \pm 1.0$  | $25968 \pm 4484$        |
|                 | Gully/channel bank        | 6                        | $0.7 \pm 1.0$                             | $0.9 \pm 0.8$  | $25713 \pm 3392$        |
|                 | Upstream reservoir area   | 1                        | 2.2                                       | 0.8            | 20338                   |
|                 | Downstream reservoir area | 1                        | 2.9                                       | 0.8            | 26137                   |

832

833 *Table 4. Correlation coefficient values (R) between catchment sediment yield, rill/interrill and*  
 834 *gully/channel erosion contribution to reservoir siltation and the tested factors in the 5 studied*  
 835 *catchments.*

836

|                                | <b>Catchment sediment yield</b> | <b>Rill/interrill relative contribution</b> | <b>Rill/interrill erosion contribution</b> | <b>Gully/channel erosion contribution</b> |
|--------------------------------|---------------------------------|---|--|---|
| <b>Area</b>                    | -0.25                           | -0.66                                       | -0.41                                      | 0.35                                      |
| <b>Mean Annual Rainfall</b>    | -0.66                           | 0.27  | -0.43                                      | -0.95*                                    |
| <b>Erosivity Index</b>         | -0.50                           | 0.39  | -0.25                                      | -0.91*                                    |
| <b>Runoff</b>                  | -0.55                           | 0.20  | -0.34                                      | -0.83                                     |
| <b>Runoff coefficient</b>      | -0.51                           | 0.05  | -0.35                                      | -0.68                                     |
| <b>Global slope index</b>      | -0.47                           | -0.86                                       | -0.69                                      | 0.40                                      |
| <b>% of tender lithology</b>   | 0.45                            | 0.09  | 0.39                                       | 0.36                                      |
| <b>% of badlands</b>           | 0.77                            | -0.12                                       | 0.56                                       | 0.93*                                     |
| <b>Drainage density</b>        | 0.89*                           | 0.28  | 0.78                                       | 0.68                                      |
| <b>% of cropland</b>           | -0.64                           | 0.35  | -0.39                                      | -0.98* <sup>#</sup>                       |
| <b>% of total managed area</b> | -0.45                           | -0.98* <sup>#</sup>                         | -0.68 <sup>#</sup>                         | 0.42                                      |

\* Correlation is significant at the 0.05 level considering the 5 study catchments.

<sup>#</sup> Correlation is significant at the 0.05 level considering 4 catchments only (Fidh Ali excluded).

837

838

839




Figure 1. Location of the selected catchments within a pilot network of monitored catchment reservoirs along the Tunisian Ridge and the Cape Bon region (modified from Temple-Boyer et al., 2007). Rainfall information (isohyet) has been superimposed.  
258x238mm (96 x 96 DPI)






Figure 2. Catchment sediment yield, rill/interrill and gully/channel erosion contribution to reservoir siltation for the five study catchments.  
417x256mm (96 x 96 DPI)






Figure 3. Sediment source apportionments for the five studied catchments using the mixing model either with the combination of 137Cs and TOC or with 137Cs only.  
235x165mm (96 x 96 DPI)