

# Quantifying the main sediment sources in agricultural landscapes of Southern Brazil cultivated with conventional and conservation practices

Olivier Evrard, Marion Le Gall, Tales Tiecher, Jean P.G. Minella, J. Laceby, Sophie Ayrault

► **To cite this version:**

Olivier Evrard, Marion Le Gall, Tales Tiecher, Jean P.G. Minella, J. Laceby, et al.. Quantifying the main sediment sources in agricultural landscapes of Southern Brazil cultivated with conventional and conservation practices. EGU General Assembly 2017, Apr 2017, Vienne, Austria. pp. EGU2017-12040. cea-02667070

**HAL Id: cea-02667070**

**<https://hal-cea.archives-ouvertes.fr/cea-02667070>**

Submitted on 31 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





## **Quantifying the main sediment sources in agricultural landscapes of Southern Brazil cultivated with conventional and conservation practices**

Olivier Evrard (1), Marion Le Gall (1), Tales Tiecher (2), Jean Paolo Gomes Minella (3), J. Patrick Laceby (1), and Sophie Ayrault (1)

(1) Laboratoire des Sciences du Climat et de l'Environnement (LSCE), CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France (olivier.evrard@lsce.ipsl.fr), (2) Departamento de Solos, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil, (3) Departamento de Solos, Universidade Federal de Santa Maria, Santa Maria, Brazil

Agricultural expansion that occurred in the 1960s in Southern Brazil significantly increased soil erosion and sediment supply to the river networks. To limit the deleterious impacts of soil erosion, conservation practices were progressively implemented in the 1990s, including the direct sowing of crops on a soil densely covered with plant residues, contour farming, the installation of ponds to trap sediment in the landscape and the use of crop rotations. However, there remains a lack of observational data to investigate the impact of these conservation practices on soil erosion and sediment supply. This data is crucial to protect soil resources and maintain the sustainability of food production systems in this region of the world characterized by a rapidly increasing population.

Accordingly, sediment sources were investigated in the Guaporé catchment (2,032 km<sup>2</sup>) representative of the cultivated environments found in this part of the world. In the upper catchment, the landscape is characterized by gentle slopes and deep soils (Ferralsols, Nitisols) corresponding to the edge of the basaltic plateau. Soybean, corn and wheat under direct sowing are the main crops in this area. In contrast, steep and shallow soils (Luvisols, Acrisols, Leptosols) highly connected to the rivers are found in the lower catchment, where tobacco and corn fields are cultivated with conventional ploughing.

These soil types were characterized by elemental geochemistry and <sup>87</sup>Sr/<sup>86</sup>Sr ratios. Sediment sources were then modelled using the optimal suite of properties (<sup>87</sup>Sr/<sup>86</sup>Sr ratios, K, Ti, Co, As, Ba, and Pb). The results demonstrate that sediment collected at the catchment outlet during two hydrological years (2012–2014) mainly originated from downstream soils (Luvisols, Acrisols, Leptosols; 92±3%), with this proportion remaining stable throughout the monitoring period. This research indicates that conservation practices implemented in the upper catchment are effective and that similar methods should be applied to downstream soils in order to conserve soil resources and limit the degradation of freshwater environments.