

Temporal evolution of sediment source contributions in a mining catchment of New Caledonia by using a tracing approach combining radionuclides and geochemistry

Virginie Sellier, Olivier Evrard, Oldrich Navratil, J. Laceby, Michel Allenbach, Irène Lefèvre

► To cite this version:

Virginie Sellier, Olivier Evrard, Oldrich Navratil, J. Laceby, Michel Allenbach, et al.. Temporal evolution of sediment source contributions in a mining catchment of New Caledonia by using a tracing approach combining radionuclides and geochemistry. EGU General Assembly 2019, Apr 2019, Vienne, Austria. pp. EGU2019-8097. cea-02667198

HAL Id: cea-02667198

<https://hal-cea.archives-ouvertes.fr/cea-02667198>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Temporal evolution of sediment source contributions in a mining catchment of New Caledonia by using a tracing approach combining radionuclides and geochemistry

Virginie Sellier (1), Olivier Evrard (1), Oldrich Navratil (2), Patrick Laceby (3), Michel Allenbach (4), and Irène Lefèvre (1)

(1) CNRS, LSCE, Gif-sur-Yvette Cedex, France (virginie.sellier@lsce.ipsl.fr), (2) Laboratoire Environnement-Ville-Société (UMR 5600), Université Lumière Lyon 2, Lyon, France, (3) Environmental Monitoring and Science Division (EMSD), Alberta Environment and Parks (AEP), 3115 – 12 Street NE, Calgary, Alberta (Canada), (4) Université de la Nouvelle-Calédonie, PPME-EA 3325, BP R4, 98 800 Nouméa (Nouvelle-Calédonie)

New Caledonia is an island located in the south-west of the Pacific Ocean, containing large nickel resources. It is currently the world's sixth largest nickel producer. Although the mining sector has contributed to the economic growth in New Caledonia since the 1870s, it is considered to be the main driver of environmental degradation. Indeed, the excessive sediment input generated by soil erosion from open-cast mining areas has strongly modified the morphology of rivers (hyper-sedimentation) and impacted the island's ecosystems (flooding, lagoon siltation, water pollution).

Nevertheless, the past and ongoing contributions of suspended sediment sources have not yet been quantified. The objective of the current research is to characterize the evolution of sediment source contributions over the last 40 years to identify the mining activities impacts (showing a peak in the 1970s) and the sediment pollution.

To this end, a pilot sediment tracing study has been conducted in one of the first areas exploited for nickel mining, the 400-km² Thio River catchment. Mining (ie. overburden, bare mining soil and access road erosion) and no-mining sources (ie. landslides, bushfires) were identified with an approach combining radionuclide and geochemical measurements. In addition, a 1.60 m long sediment core was collected in the Thio river deltaic floodplain in April 2016. Sediment dating was carried out on the chronology of flooding based on a rainfall-discharge statistical analysis in the Thio region. A distribution modelling approach was used to reconstruct sediment source contribution temporal variability in the successive sediment layers.

Results demonstrate that the contribution of mining sources largely dominated with a mean contribution of 65 % over the entire profile. The fifteen flooding events identified by the statistical analysis generated the deposition of around 1 m of sediment in the floodplain at that location over the last forty years. An increase in the mining source contribution was observed along the sediment profile and particularly at 1-m depth, i.e. a depth corresponding to the 1970s nickel production peak.

In the future, similar studies should be carried out in other catchments draining mines in New Caledonia to investigate whether the temporal changes in the mining source contribution were similar to those observed in the Thio river catchment in order to guide the implementation of control measures.