

## Effects of soil relocation on soil organic carbon stocks in hilly agricultural landscapes

Sebastian Doetterl, Olivier Evrard, Antoine Stevens, Bas van Wesemael,  
Kristof Oost

► **To cite this version:**

Sebastian Doetterl, Olivier Evrard, Antoine Stevens, Bas van Wesemael, Kristof Oost. Effects of soil relocation on soil organic carbon stocks in hilly agricultural landscapes. EGU General Assembly 2010, May 2010, Vienne, Austria. pp. EGU2010-15192. cea-02667342

**HAL Id: cea-02667342**

**<https://hal-cea.archives-ouvertes.fr/cea-02667342>**

Submitted on 31 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





## **Effects of soil relocation on soil organic carbon stocks in hilly agricultural landscapes**

Sebastian Doetterl (1), Olivier Evrard (2), Antoine Stevens (1), Bas van Wesemael (1), and Kristof Van Oost (1)

(1) Geography Department, Université catholique de Louvain, Belgium, (2) Laboratoire des Sciences du Climat et de l'Environnement, CNRS, Gif-sur-Yvette, France

Agricultural land use is well known to have a major impact on altering physical and chemical soil parameters. Among these, soil organic matter (SOM) is recognized as an important parameter for soil fertility and must be considered for modeling global carbon fluxes due to its magnitude and high content of soil organic carbon (SOC) in many soils. Former investigations on arable land concentrated on the plough layers of soils, ignoring redistribution of SOM due to erosional and depositional processes. It is widely discussed whether soil erosion can function as a carbon sink due to carbon burial on sedimentary sites, or as a source due to accelerated decomposition of SOC during transportation.

Beside land use, we assume topography to be the controlling unit for modeling total SOC. We postulate that the implication of vertical and lateral soil fluxes will largely improve our understanding of carbon dynamics on a landscape scale. We conducted a sampling campaign in different hilly agricultural regions in Central Europe and performed depth dependent analysis of SOC contents from agricultural soils with different soil substrates and land use following geomorphic gradients. We accompanied this study with an assessment of recent (since approx. 1950) lateral soil fluxes on our research sites using  $^{137}\text{Cs}$  radioisotopes and measured depth dependent bulk density for carbon stock estimations.

Our data gives clear evidence that traditional perceptions on SOC dynamics leave out major slope related controls and suffer from a lack of understanding landscape dynamics. An integrative approach from point to landscape scale of pedogenic and geomorphic processes and its effects on SOC dynamics is needed to improve our understanding of its role in the global carbon cycle.