



**HAL**  
open science

## Tracking the dispersion of particle-borne radioactive contamination along coastal rivers of Fukushima Prefecture (2011-2014)

Olivier Evrard, Yuichi Onda, Hugo Lepage, Fabien Pointurier, Caroline Chartin, J. Patrick Lacey, Anne-Claire Pottin, Amélie Hubert, Irène Lefevre, Sophie Ayrault, et al.

### ► To cite this version:

Olivier Evrard, Yuichi Onda, Hugo Lepage, Fabien Pointurier, Caroline Chartin, et al.. Tracking the dispersion of particle-borne radioactive contamination along coastal rivers of Fukushima Prefecture (2011-2014). ICOBTE 2015: 13th International Conference on the Biogeochemistry of Trace Elements, Jul 2015, Fukuoka, Japan. cea-02669093

**HAL Id: cea-02669093**

**<https://cea.hal.science/cea-02669093>**

Submitted on 31 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Tracking the dispersion of particle-borne radioactive contamination along coastal rivers of Fukushima Prefecture (2011–2014)

Olivier Evrard <sup>a\*</sup>, Yuichi Onda <sup>b</sup>, Hugo Lepage <sup>a</sup>, Fabien Pointurier <sup>c</sup>, Caroline Chartin <sup>d</sup>, J. Patrick Lacey <sup>a</sup>, Anne-Claire Pottin <sup>c</sup>, Amélie Hubert <sup>c</sup>, Irène Lefèvre <sup>a</sup>, Sophie Ayrault <sup>a</sup>, Philippe Bonté <sup>a</sup>

<sup>a</sup> *Laboratoire des Sciences du Climat et de l'Environnement (LSCE/IPSL) – Unité Mixte de Recherche 8212 (CEA-CNRS-UVSQ) – 91 198 Gif-sur-Yvette Cedex (France)*

<sup>b</sup> *Center for Research in Isotopes and Environmental Dynamics (CRIED), Tsukuba University, Tsukuba (Japan)*

<sup>c</sup> *CEA, DAM, DIF, 91297 Arpaçon (France)*

<sup>d</sup> *TECLIM, Université catholique de Louvain (Belgium)*

\* Corresponding author: [olivier.evrard@lsce.ipsl.fr](mailto:olivier.evrard@lsce.ipsl.fr)

The Fukushima Dai-ichi nuclear power plant (FDNPP) accident that occurred in March 2011 led to the formation of a 3000-km<sup>2</sup> radioactive pollution plume and resulted in the fallout consisting predominantly of radiocesium (<sup>137</sup>Cs and <sup>134</sup>Cs) on soils located up to 70 km to the northwest of the damaged site. This mountainous region is drained by several coastal rivers to the Pacific Ocean, and original sediment fingerprinting methods had to be developed to track the dispersion of contaminated sediment across these coastal catchments. Sediment fingerprinting based on the heterogeneous deposition of metastable silver-110 (<sup>110m</sup>Ag) across the investigated catchments was shown to provide relevant information on the dispersion of contaminated sediment. Based on the measurement of the <sup>110m</sup>Ag:<sup>137</sup>Cs activity ratio in soils and river sediment, we demonstrated the occurrence of a seasonal cycle of soil erosion during typhoons followed by sediment export in rivers during the subsequent spring snowmelt in 2011 and 2012. However, due to the rapid decay of <sup>110m</sup>Ag (half-life of 250 days), alternative methods had to be developed to continue tracking sediment from 2013 onwards. We therefore used local ground dose rate and <sup>241</sup>Pu/<sup>239</sup>Pu atom ratio measurements to further document this dispersion. We thereby demonstrated that, although overall sediment contamination decreased two years after the accident, decontamination of paddy fields led to the local supply of contamination to the rivers. Ongoing work is focused on the development of a sediment fingerprinting technique based on geochemical properties characterizing the different soil types found in the region. In future, we expect that the contaminated sediment stored in river channels and reservoirs will represent a significant management challenge for local authorities.