



**HAL**  
open science

## **Radiolytic approach for efficient, selective and catalyst-free CO<sub>2</sub> conversion at room temperature**

Changjiang Hu, Sarah Al Gharib, Yunlong Wang, Pingping Gan, Qiu hao Li, Sergey A Denisov, Sophie Le Caër, Jacqueline Belloni, Jun Ma, Mehran Mostafavi

► **To cite this version:**

Changjiang Hu, Sarah Al Gharib, Yunlong Wang, Pingping Gan, Qiu hao Li, et al.. Radiolytic approach for efficient, selective and catalyst-free CO<sub>2</sub> conversion at room temperature. *ChemPhysChem*, 2021, 22 (18), pp.1900-1906. 10.1002/cphc.202100378 . cea-03287511

**HAL Id: cea-03287511**

**<https://cea.hal.science/cea-03287511>**

Submitted on 15 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Radiolytic Approach for Efficient, Selective and Catalyst-free CO<sub>2</sub> conversion at room temperature

Changjiang Hu,<sup>a†</sup> Sarah Al Gharib,<sup>b†</sup> Yunlong Wang,<sup>a</sup> Pingping Gan,<sup>a</sup> Qiu hao Li,<sup>a</sup>

Sergey A. Denisov,<sup>b</sup> Sophie Le Caer,<sup>c</sup> Jacqueline Belloni,<sup>b</sup> Jun Ma,<sup>a\*</sup> Mehran Mostafavi<sup>b\*</sup>

*† Both Changjiang Hu and Sarah Al Gharib are considered as first authors.*

*[a] C.J. Hu, Y.L. Wang, P.P. Gan, Q.H. Li, Prof. J. Ma.*

*Department of Materials Science and Technology,*

*Nanjing University of Aeronautics and Astronautics, Nanjing 211106, China*

*E-mail: junma@nuaa.edu.cn*

*[b] Dr S. Al Gharib, Dr S. A. Denisov, Dr J. Belloni, and Pr M. Mostafavi  
Institut de Chimie Physique CNRS-Université Paris-saclay, Orsay, France*

*[c] Dr S. Le Caer NIMBE, UMR 3685 CEA, CNRS, Université Paris-Saclay, CEA Saclay  
91191 Gif-sur-Yvette Cedex, France*

**Abstract:** The present study proposes a new approach for direct CO<sub>2</sub> conversion using primary radicals from water irradiation. In order to ensure reduction of CO<sub>2</sub> into CO<sub>2</sub><sup>•-</sup> by all the hydrated electrons, we use formate ions to scavenge simultaneously the parent oxidizing radicals H<sup>•</sup> and OH<sup>•</sup> producing the same transient CO<sub>2</sub><sup>•-</sup> radicals. Conditions are optimized to obtain the highest conversion yield of CO<sub>2</sub>. The goal is achieved under mild conditions of room temperature, neutral pH and 1 atm of CO<sub>2</sub> pressure. All the available radicals are exploited for selectively converting CO<sub>2</sub> into oxalate that is accompanied by H<sub>2</sub> evolution. The mechanism presented accounts for the results and also sheds light on the data in the literature. The radiolytic approach is a mild and scalable route of direct CO<sub>2</sub> capture at the source in industry and the products, oxalate salt and H<sub>2</sub>, can be easily separated.

## Introduction

The excessive anthropometric carbon dioxide (CO<sub>2</sub>) emission, that is responsible for climate changes, has led to worldwide efforts for a drastic diminution of its atmosphere level. Beyond the progressive abandonment of fossil fuels and the storage and sequestration of CO<sub>2</sub> surplus, the alternative strategy has been actively developed for efficient and cost-effective CO<sub>2</sub> conversion into other eco-friendly and value-added chemicals.<sup>[1-3]</sup>

Due to CO<sub>2</sub> chemical stability, the existing methods of its conversion into less oxidized

chemical species typically require a source of energy (i.e., heat, light, or electricity) to drive one- or multiply electron transfer reactions. For example, the reduction of CO<sub>2</sub> by coal to form CO occurs at very high temperature (> 700 °C) due to a large positive reaction enthalpy ( $\Delta H^{\circ}_{298\text{K}} = 172 \text{ kJ mol}^{-1}$ ).<sup>[4]</sup> The use for CO<sub>2</sub> reduction electron-rich chemicals, e.g. CH<sub>4</sub> and even H<sub>2</sub>, also consumes significant amounts of energy, even in the presence of a catalyst.<sup>[5-7]</sup> The photocatalytic and electrocatalysis reduction of CO<sub>2</sub> using renewable energy resources likely accomplish the carbon neutral energy cycle while synthesizing fuels. However, their slow kinetics together with side reactions lead to a low efficiency and diverse products.<sup>[8,9]</sup> The newly -developed catalysis covering various matter-scales exhibits superior performance for CO<sub>2</sub> conversion,<sup>[10-12]</sup> but their ultimate practical viability is greatly hindered by several limiting factors such as high cost, lack of long-term stability, and upscale synthesis methodologies.<sup>[13]</sup> All these methods generally use multi-step processes based on complex chemistry, and essentially call for more direct and selective pathways.

The hydrated electron ( $e_{\text{aq}}^-$ ) is the most powerful reducing species with a standard reduction potential of  $-2.9 \text{ V}_{\text{NHE}}$  that is larger than the reduction potential of CO<sub>2</sub>/CO<sub>2</sub><sup>•-</sup> ( $-2.14 \text{ V}_{\text{NHE}}$  in water).<sup>[14]</sup> Therefore,  $e_{\text{aq}}^-$  can reduce aqueous CO<sub>2</sub> at room temperature without any catalyst. By taking advantage of the  $e_{\text{aq}}^-$  chemistry, the approaches of electrochemistry,<sup>[15]</sup> and more recently photoelectrochemistry<sup>[16-18]</sup> have been developed to reduce CO<sub>2</sub>, but with a low Faradaic efficiency.

Herein, we exploit high-energy ionizing radiation as a promising method for aqueous CO<sub>2</sub> reduction.<sup>[19]</sup> As the reducing radicals are directly formed in water at room temperature, this method is effective without requiring the use of any catalyst. However, up to now, the appropriated chemical conditions were not explored to enable maximizing CO<sub>2</sub> conversion yield. Indeed, the fast back-oxidation by the symbiotic OH<sup>•</sup> radicals has to be eliminated which remains a challenge for the CO<sub>2</sub> conversion efficiency.

This work aims to reduce the aqueous CO<sub>2</sub> into CO<sub>2</sub><sup>•-</sup>, the key precursor of radiolytic products, by utilizing all primary radiation-induced water radicals. Among the few possible candidates to create the fully reducing conditions in irradiated aqueous CO<sub>2</sub> solutions, formate ions were chosen as OH<sup>•</sup> and H<sup>•</sup> scavengers instead of methanol<sup>[20]</sup> or other organic chemicals,<sup>[21]</sup> because in this case, the reaction product is the same CO<sub>2</sub><sup>•-</sup> radical as in the reduction of CO<sub>2</sub> by  $e_{\text{aq}}^-$ . The oxidation reactions of formate by OH<sup>•</sup> and H<sup>•</sup> into CO<sub>2</sub><sup>•-</sup> and its second-order decay have already been extensively studied.<sup>[22-28]</sup> However, this process is hindered by H<sup>+</sup>-catalysis leading to CO<sub>2</sub><sup>•-</sup> disproportionation back to formate and CO<sub>2</sub>.<sup>[29]</sup> Therefore, the present radiolytic experiments of CO<sub>2</sub> reduction in presence of formate, under catalyst-free aqueous

environment, were first performed at 1 atmosphere at various pH and then extended to higher CO<sub>2</sub> pressure up to 30 atmospheres. Based on the stoichiometric radical yields values in water, we establish the reaction mechanism under various conditions in order to optimize the CO<sub>2</sub> conversion.

## **Experimental methods**

### **Materials**

The gases CO<sub>2</sub> (99.999%) and Ar (99.999%) were purchased from Air Liquide Industrial Gases Company or Nanjing Shangyuan Industrial gas plant. Sodium formate NaCOOH and formic acid HCOOH were purchased from Merck. Potassium oxalate NaC<sub>2</sub>O<sub>4</sub> was obtained from Fluka, NaOH from Sigma Aldrich Chemistry and H<sub>2</sub>SO<sub>4</sub> (98%) from Nanjing Chemical Reagent Co., Ltd). Aqueous solutions were prepared in deionized water (Millipore, resistivity 18.2 MΩ cm).

13 mL of the samples were saturated at 20 °C with CO<sub>2</sub> through the septum. The hydration of CO<sub>2</sub> into H<sub>2</sub>CO<sub>3</sub> is weak:  $K ([H_2CO_3]/[CO_2]) = 10^{-2}$ . The pH was controlled by adequate addition of NaOH or HCO<sub>2</sub>H before the saturation by CO<sub>2</sub>.

### ***γ-Irradiation and analysis***

γ-radiolysis was carried out at room temperature under CO<sub>2</sub> atmosphere using a panoramic <sup>60</sup>Co source. The dose rate, measured by the Fricke dosimeter, was 8 kGy h<sup>-1</sup> (1 Gy = 1 J kg<sup>-1</sup>), and the doses were increased up to 60 kGy.

In the case of the experiments performed under pressure, a stainless-steel vessel was used at the Nanjing University. We found very small amount of CH<sub>4</sub> at high pressure and one of the reasons of this production could be the reaction at the surface of stainless steel. However, as the CH<sub>4</sub> amount is very low we neglect it. Moreover, the mechanism and the yield were deduced from the experiments obtained at one atmosphere in glass vessels where we did not find CH<sub>4</sub>.

Irradiation was carried out with a <sup>60</sup>Co source. The dose rate in the vessel was determined to be 0.49 kGy h<sup>-1</sup>.

Oxalate ions are quantified thanks to their optical absorption spectra. Measurements were performed using a Hewlett-Packard spectrophotometer). The absorbance is measured at 250 nm ( $\epsilon = 28 \text{ mol}^{-1} \text{ L cm}^{-1}$ ) where the absorbance of formate anion is negligible (Figure SI-1). The sodium oxalate was also analyzed by high performance liquid chromatography (HPLC) (Figure SI-2), (Eclassical 3100 HPLC Dalian Elite Analytical Instruments) on a chromatographic

column Shodex Sugar SH1011 ( $8.0 \times 300 \text{ mm} \times 5.0 \text{ }\mu\text{m}$ ) and the eluent was an aqueous solution of  $\text{H}_2\text{SO}_4$  (0.025 mol/L) at a flow rate of 1 mL/min and  $50 \text{ }^\circ\text{C}$  with a UV absorption detection.

The gases produced ( $\text{H}_2$  and  $\text{CO}$ ) were measured by a gas chromatography set-up (GC9790 II co. Fuli instruments) equipped with a thermal conductivity detector (TCD) for  $\text{H}_2$  and a flame ionization detector (FID) for  $\text{CO}$  (Figures SI-3). To examine whether the  $\text{CO}$  arose from the ionization/excitation in the solution or in the headspace gaseous  $\text{CO}_2(\text{g})$ , control radiolytic experiments of  $\text{CO}_2(\text{g})$  alone were performed (Figure SI-4). The contribution of the emitted  $\text{CO}$  from the gaseous decomposition is low (for example at 1  $\text{CO}_2$  atm it is less than 2% of the total amount including the solution) (Figure SI-4), which is consistent with earlier radiolytic studies of gas-phase reactions that are often characterized by low product yields. The  $\text{CO}$  measurements were then systematically corrected from the  $\text{CO}$  formation arising from irradiated  $\text{CO}_2(\text{g})$  under the same conditions.


## Results and Discussion

### *pH effects on radiolytic product yields at 1 atm $\text{CO}_2$*

The amounts of oxalate,  $\text{CO}$  and  $\text{H}_2$  are measured versus irradiation dose and pH in aqueous solutions under 1 atm of  $\text{CO}_2$  (or  $3.8 \times 10^{-2} \text{ mol L}^{-1}$  of  $\text{CO}_2$ ). First, preliminary experiments performed without formate on the radiolysis of  $\text{CO}_2$ -saturated solutions showed that the oxalate yield is negligible, and the major product is  $\text{H}_2$ . The radiolytic yield of  $\text{H}_2$ ,  $G(\text{H}_2) \sim 0.4 \times 10^{-7} \text{ mol J}^{-1}$ , is measured from the slope of the species amounts versus the absorbed dose (Figure SI-3).

In the presence of sodium formate ( $5 \times 10^{-2} \text{ mol L}^{-1}$ ) as the  $\text{OH}^\bullet$  and  $\text{H}^\bullet$  radicals scavenger, the  $\text{H}_2$  radiolytic yield,  $G(\text{H}_2)$  decreases from  $2.75 \times 10^{-7} \text{ mol J}^{-1}$  at pH 1 to  $1.04 \times 10^{-7} \text{ mol J}^{-1}$  at pH 7. The  $G(\text{CO})$  value at 1 atm  $\text{CO}_2$  is very low at pH 7 and its maximum reaches  $0.17 \times 10^{-7} \text{ mol J}^{-1}$  at pH 2 and 3 (Figure 1 right).

The oxalate concentration increases at all pH linearly with the dose (Figure 1 left). The  $G(\text{C}_2\text{O}_4^{2-})$  value increases from  $0.05 \times 10^{-7} \text{ mol J}^{-1}$  at pH 3 to  $3.5 \times 10^{-7} \text{ mol J}^{-1}$  at pH 7 following an opposite trend compared to  $\text{H}_2$  (Figure 1 right). Below pH = 3 no oxalate is observed in  $\text{CO}_2$ -saturated formate solutions.



**Figure 1. (Left) Representative measurements for the concentrations of oxalate (top), CO (middle) H<sub>2</sub> (bottom), produced versus the irradiation dose at various pH. (Right) Yields of H<sub>2</sub>, CO and oxalate in aqueous solution containing formate ( $5 \times 10^{-2} \text{ mol L}^{-1}$ ) in the presence of CO<sub>2</sub> at 1 atm (the concentration of CO<sub>2</sub> in water is then  $3.8 \times 10^{-2} \text{ mol L}^{-1}$ ). Dotted line: Maximum oxalate yield, which is equal to  $\frac{1}{2} (G(e_{aq}^-) + G(H^*) + G(OH^*)) = 3.6 \times 10^{-7} \text{ mol J}^{-1}$  (see text)**

Around pH 7, according to the mechanism described below, the CO<sub>2</sub> consumption yield ( $G(-\text{CO}_2)$ ) corresponds to about the formate consumption yield ( $G(OH^*) + G(H^*)$ ), or to the formation yield of oxalate ( $\frac{1}{2}(G(OH^*) + G(H^*) + (G(e_{aq}^-)))$ ). Therefore, the CO<sub>2</sub> concentration after 60 kGy irradiation is estimated to be  $1.3 \times 10^{-2} \text{ mol L}^{-1}$ . That represents 66% of the initial concentration that have been converted. By comparison, the decrease of the formate concentration is only 17%. In addition, the yields of H<sub>2</sub> and oxalate remain constant in this CO<sub>2</sub> concentration range. We also showed that  $G(\text{C}_2\text{O}_4^{2-})$  is independent on  $[\text{HCO}_2^-]$  from  $5 \times 10^{-2}$  to  $10^{-1} \text{ mol L}^{-1}$ . Therefore, these observations at very high doses indicate that the radiolytic approach is effective for sustaining CO<sub>2</sub> conversion in large amounts and that formate plays a significant role in the mechanism promoting the practicable and feasibility of this strategy.


### CO<sub>2</sub> pressure effects on the radiolytic yields

The increase of the CO<sub>2</sub> pressure induces an increase of the CO<sub>2</sub> aqueous concentration, and thus an acidification of the solution. The final pH obtained after having imposed different values of CO<sub>2</sub> pressure can be calculated according to the various acid-base equilibria of the system for two initial pH (Figure SI-5).<sup>[30]</sup> Noteworthy, both HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> are present in solutions. However, the rate constants of the e<sub>aq</sub><sup>-</sup> scavenging by these species ( $k < 10^6 \text{ M}^{-1} \text{ s}^{-1}$ ) are 3 orders of magnitude lower than that of reaction with CO<sub>2</sub>. Therefore, HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> are omitted in the reaction mechanism.

In the absence of formate, in CO<sub>2</sub>-saturated solutions, the substantial increase of the CO<sub>2</sub> concentration by the pressurization leads to the rise of G(CO) from 0 to  $0.18 \times 10^{-7} \text{ mol J}^{-1}$  (30 atm, pH 5.3), but to a decrease of G(H<sub>2</sub>) from 0.4 to  $0.1 \times 10^{-7} \text{ mol J}^{-1}$ , respectively (Figure SI-3). The oxalate yield G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) is negligible. In the presence of formate, for final pH ranging from 3.9 to 3.2 for a CO<sub>2</sub> pressure varying from 1 to 30 atm, we observe a decrease of the H<sub>2</sub> yield and a marked increase of the CO yield without any oxalate formation (Figure 2 left).

For final pH around 7, the effect of CO<sub>2</sub> pressure on the yields of H<sub>2</sub> and CO presents similar trend as at pH 3. As shown above (Figure 1 Right), G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) measured at 1 atm of CO<sub>2</sub> is the highest and equal to  $3.5 \times 10^{-7} \text{ mol J}^{-1}$ . However, when the CO<sub>2</sub> pressure increases, G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) decreases down to a plateau value of  $1.2 \times 10^{-7} \text{ mol J}^{-1}$  at 30 atm (pH 5.5) (Figure 2 Right). Obviously, in spite of a marked increase of the CO<sub>2</sub> concentration, the increasing acidity caused by pressurization leads to a decrease of the oxalate yield.

It is worthy to note that CO<sub>2</sub> pressure effects on decreasing G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) and increasing G(CO) (Figure 2), are comparable to the pH effect observed at 1 atm pressure (Figure 1). However, the H<sub>2</sub> formation yield does decrease to lower values at higher CO<sub>2</sub> pressure (Figure 2) which is in contrast to the G(H<sub>2</sub>) increase when pH decreases at 1 CO<sub>2</sub> atm (Figure 1).




**Figure 2. Radiolytic formation yields of H<sub>2</sub> and CO versus CO<sub>2</sub> pressure from 1 to 30 atm (Left) in formic acid solution (pH 3). In these conditions, G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) is negligible at any pressure. (Right) in solution containing initially 0.5 mol L<sup>-1</sup> NaOH (final pH ranges from 7 to 5.5)**

### **Mechanism**

#### **Neutral medium, 1 atm CO<sub>2</sub>**

The generation of various species from water radiolysis at pH 7 is expressed as:



(in brackets are the yields in 10<sup>-7</sup> mol J<sup>-1</sup> unit).<sup>[31, 32]</sup> These yields correspond to a scavenging factor of 10<sup>7</sup> s<sup>-1</sup> (or 100 ns after the radical production and their mutual reactions in spurs). Note that the direct effect of radiation on CO<sub>2</sub> is negligible because of the low electron fraction (< 0.1) compared to water even at 30 atm pressure.


The reduction of CO<sub>2</sub> occurs through the reaction with e<sub>aq</sub><sup>-</sup> (reaction 2 in Table 1). This reaction has been found to be almost controlled by diffusion (k<sub>2</sub> = 8.2 × 10<sup>9</sup> M<sup>-1</sup>s<sup>-1</sup>),<sup>[20, 21, 33]</sup> and the activation energy of k<sub>2</sub> between 5 and 200 °C was measured to be 15.9 kJ mol<sup>-1</sup>.<sup>[20]</sup> The value of the rate constant k<sub>2</sub> warrants that all the solvated electrons reduce efficiently CO<sub>2</sub>. The second-order reaction of dimerization of radicals CO<sub>2</sub><sup>•-</sup> (reaction 3 in Table 1) into oxalate is in competition with the H<sup>+</sup>-catalysed disproportionation (reaction 4 in Table 1) which leads to the formation of the initial reactants.<sup>[22-29]</sup>

In the absence of formate as an OH<sup>•</sup> scavenger, the oxalate yield G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) = 0.02 × 10<sup>-7</sup> mol J<sup>-1</sup>, is much less than G(e<sub>aq</sub><sup>-</sup>)/2. This very low yield and inefficiency of the process are most likely ascribed to significant loss of CO<sub>2</sub><sup>•-</sup> caused by a secondary reaction with OH<sup>•</sup> radicals (reaction 5 in Table 1) faster than their dimerization (reaction (3)). However, the G(H<sub>2</sub>) value is close to the primary H<sub>2</sub> yield G(H<sub>2</sub>) = 0.42 × 10<sup>-7</sup> mol J<sup>-1</sup>.

**Table 1. Reactions and corresponding rate constants involved in the mechanism of  $C_2O_4^{2-}$ ,  $H_2$  and  $CO$  formation**

<b>Reactions</b>	<b>Rate constants (L mol<sup>-1</sup> s<sup>-1</sup>) at 20 °C</b>
$e_{aq}^- + CO_2 \rightarrow CO_2^{\cdot -}$	(2) $8.2 \times 10^9$ [20]
$CO_2^{\cdot -} + CO_2^{\cdot -} \rightarrow C_2O_4^{2-}$	$1.4 \times 10^9$ [29]
(3)	
$CO_2^{\cdot -} + CO_2^{\cdot -} + H^+ \rightarrow {}_2OC-CO_2H^{\cdot} \rightarrow CO_2 + HCO_2^{\cdot}$	(4) $1.4 \times 10^9$ [29]
$OH^{\cdot} + CO_2^{\cdot -} \rightarrow OH^- + CO_2$	
(5)	
$OH^{\cdot} + HCO_2^{\cdot} \rightarrow H_2O + CO_2^{\cdot -}$	(6) $4.1 \times 10^9$ [22]
$H^{\cdot} + HCO_2^{\cdot} \rightarrow H_2 + CO_2^{\cdot -}$	(7) $2.1 \times 10^8$ [31]
$e_{aq}^- + H^+ \rightarrow H^{\cdot}$	(8) $2.3 \times 10^{10}$ [31]
$H^{\cdot} + CO_2^{\cdot -} \rightarrow CO + OH^-$	(9)
$CO_2^{\cdot -} + CO_2 \rightleftharpoons [O_2C-CO_2]^{\cdot -}$	(10)
$[O_2C-CO_2]^{\cdot -} + [O_2C-CO_2]^{\cdot -} + H^+ \rightarrow HCO_2^{\cdot} + 3 CO_2$	(11)
$[O_2C-CO_2]^{\cdot -} + [O_2C-CO_2]^{\cdot -} \rightarrow C_2O_4^{2-} + 2 CO_2$	(12)
$CO_2 + e_{aq}^- \rightarrow CO_2^{\cdot -}$	(13)
$[O_2C-CO_2]^{\cdot -} + [O_2C-CO_2]^{\cdot -} + H^+ \rightarrow CO + HCO_3^- + 2 CO_2$	(14)

In the presence of formate, at neutral pH, the  $H_2$  yield corresponds to the sum of the primary yield  $G(H_2) = 0.42 \times 10^{-7} \text{ mol J}^{-1}$  in reaction (1) and of the scavenging yield of  $H^{\cdot}$  radicals by formate ( $G(H^{\cdot}) = 0.62 \times 10^{-7} \text{ mol J}^{-1}$ ) in reaction (7). The experimental value  $G(H_2) = 1.04 \times 10^{-7} \text{ mol J}^{-1}$  (Figure 1 right) is in fair agreement with this sum  $G(H_2) + G(H^{\cdot})$  (Figure 3 and Table 2). That means that in addition to primary radiolytic  $H_2$ , all the  $H^{\cdot}$  radicals are exploited to produce  $H_2$ .



**Figure 3. Scheme of the radiolysis of CO<sub>2</sub>-saturated solutions of sodium formate.** At pH 7 and 1 atm (in black), the CO<sub>2</sub><sup>•-</sup> dimerization and the CO<sub>2</sub> conversion into oxalate are optimized:  $G((C_2O_4^{2-})) = \frac{1}{2} (G(e_{aq}^-) + G(H^\bullet) + G(OH^\bullet)) = 3.5 \times 10^{-7} \text{ mol J}^{-1}$  and  $G(H_2) = (G(H^\bullet) + G(H_2)) = 1.04 \times 10^{-7} \text{ mol J}^{-1}$ . At pH < 7,  $G(C_2O_4^{2-})$  decreases because the H<sup>+</sup>-catalyzed disproportionation of CO<sub>2</sub><sup>•-</sup> is involved and becomes significant. At high CO<sub>2</sub> pressure (in red), CO<sub>2</sub> scavenges early the electrons before their hydration and the radicals CO<sub>2</sub><sup>•-</sup> react with H<sup>•</sup> radicals in spurs yielding CO. Simultaneously, the pH decrease favors the disproportionation, so that the oxalate yield becomes negligible. The arrow on the left indicates the time after the radiation-induced ionisation and excitation of the aqueous solution.

**Table 2. Summary of product yields and predominant mechanisms in the radiolysis of CO<sub>2</sub> solutions with sodium formate  $5 \times 10^{-2} \text{ mol L}^{-1}$ .** *dispr* and *dim* are the disproportionation and the dimerization of the precursor radical CO<sub>2</sub><sup>•-</sup>, respectively. Primary yields of species arising from water radiolysis are in italic.

Irradiated Systems	$G(H_2)$ ( $10^{-7} \text{ mol J}^{-1}$ )	$G(CO)$ ( $10^{-7} \text{ mol J}^{-1}$ )	$G((CO_2)_2^{2-})$ ( $10^{-7} \text{ mol J}^{-1}$ )
CO <sub>2</sub> 30 atm	0.16	0.7	0
pH 3	$\ll G(H_2) + G(H^\bullet)_{acidic}$	$= G(H_2O^*)$	<i>Only dispr</i>
CO <sub>2</sub> 30 atm	0.13	0.7	1.3
pH 5.3	$\ll G(H_2) + G(H^\bullet)_{acidic}$	$= G(H_2O^*)$	<i>Competition dim/dispr</i>

CO <sub>2</sub> 1 atm	2.0		0
pH 3	$< G(H_2) + G(H^*)_{acidic}$	0.14	<i>Only dispr</i>
CO <sub>2</sub> 1 atm	1.0		1.5
pH 5	$= G(H_2) + G(H^*)_{neutral}$	0.08	<i>Competition dim/dispr</i>
			3.5
CO <sub>2</sub> 1 atm	1.04		$= \frac{1}{2} (G(e_{aq}^-) + G(H) +$
pH 7	$= G(H_2) + G(H^*)_{neutral}$	0.08	$G(OH^*))$
			<i>Only dim</i>

Part of CO<sub>2</sub><sup>-</sup> produced in CO<sub>2</sub>-saturated sodium formate solutions is due to the e<sub>aq</sub><sup>-</sup> scavenging by CO<sub>2</sub> (reaction 2) and the other part is due to the scavenging of OH<sup>\*</sup> and H<sup>\*</sup> by formate (reactions 6 and 7). According to k<sub>6</sub> and k<sub>7</sub> values, and to the formate concentration, the scavenging half-time for reaction (6) is 4 ns and  $G_{4ns}(OH^*) \approx 3.3 \times 10^{-7} \text{ mol J}^{-1}$ .<sup>[34,35]</sup> Therefore, from reactions (2, 3, 5, 6 and 7), the maximum value of oxalate yield is  $G(C_2O_4^{2-})_{calc} = \frac{1}{2} (G(e_{aq}^-) + G(H^*) + G(OH^*)) = 3.6 \times 10^{-7} \text{ mol J}^{-1}$  (dotted line in Figure 1 Right), with  $G_{4ns}(e_{aq}^-) = G_{4ns}(OH^*) \approx 3.3 \times 10^{-7} \text{ mol J}^{-1}$  and  $G(H^*) \approx 0.62 \times 10^{-7} \text{ mol J}^{-1}$ . The experimental yield obtained at pH 7:  $G(C_2O_4^{2-}) = 3.5 \times 10^{-7} \text{ mol J}^{-1}$  is in excellent agreement with this maximum value. It is remarkable that under these mild conditions, the e<sub>aq</sub><sup>-</sup> scavenging, which yields the conversion of CO<sub>2</sub> at 1 atm into oxalate, is complete and that the disproportionation path (4) is inactive. Indeed, in formate and CO<sub>2</sub> solutions, efficient reductive conditions of the aqueous solutions are fulfilled because, as an OH<sup>\*</sup> and H<sup>\*</sup> scavenger, formate inhibits the recombination reaction (5).<sup>[36,37]</sup> The scheme of the mechanism is summarized in Figure 3. The e<sub>aq</sub><sup>-</sup> scavenging for selectively converting CO<sub>2</sub> into oxalate is completely exploited. Moreover, the H<sup>\*</sup> radical scavenging by formate yielding H<sub>2</sub>, a useful fuel, is also complete. The efficiency of the absorbed radiation energy is thus 70% for the oxalate formation (32% for the reducing conversion of CO<sub>2</sub> and 38% for the formate oxidation),<sup>[38]</sup> plus 14.5% for the hydrogen formation.

### ***Acidic medium***

The yield of H<sub>2</sub> increases in acid medium at CO<sub>2</sub> atmospheric pressure (Figure 1) because e<sub>aq</sub><sup>-</sup> are replaced by H<sup>\*</sup> radicals via reaction (8) and more H<sup>\*</sup> radicals are scavenged by formate (reaction (7)). Thus at pH 1 under 1 atm CO<sub>2</sub>, the yield would be  $G(H_2)_{calc} = G(H^*)_{acidic} + G(H_2) = (3.42 + 0.42) \times 10^{-7} \text{ mol J}^{-1}$ . However, the experimental value is lower:  $(G(H_2))_{exp} = 2.8 \times 10^{-7}$

mol J<sup>-1</sup>), showing that a small part of H<sup>•</sup> radicals are thus lost in another reaction, such as the CO<sub>2</sub><sup>•-</sup> radical reduction into CO (reaction (9)). This reaction may occur within the ionisation spurs before the homogenous dispersion of the radicals and explains the CO formation, although the experimental result is only G(CO) = 0.17 × 10<sup>-7</sup> mol J<sup>-1</sup>.

At 1 atm of CO<sub>2</sub>, the oxalate yield G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) decreases with the pH decrease (Figure 1). However, the simultaneous formation yield of the oxalate precursor CO<sub>2</sub><sup>•-</sup> (either formed by reactions 2 or 6 and 7) is pH-independent. In a previous mechanistic study<sup>[29]</sup> of CO<sub>2</sub><sup>•-</sup> arising from CO oxidation by OH<sup>•</sup> (without CO<sub>2</sub>), it was shown that at low pH the oxalate yield G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) was strongly inhibited, and in contrast G(CO<sub>2</sub>) increased, owing to the disproportionation of CO<sub>2</sub><sup>•-</sup> into CO<sub>2</sub> and formate (reaction 4). However, the rate constant of the second-order decay of CO<sub>2</sub><sup>•-</sup> (2k = 1.3 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup>) is also pH-independent. They concluded that the disproportionation (4) was catalysed by protons and favored by the acid form HCO<sub>2</sub><sup>•</sup> of the radical CO<sub>2</sub><sup>•-</sup> (pK<sub>a</sub> (HCO<sub>2</sub><sup>•</sup>/CO<sub>2</sub><sup>•-</sup>) ≈ 2.3). The protons would govern the disproportionation because of the presence of an intermediate head-to-tail adduct linking the carbon atom of one molecule to an oxygen atom of the other one.<sup>[29]</sup> Other values of CO<sub>2</sub><sup>•-</sup> pK<sub>a</sub> were proposed in the literature.<sup>[26, 27, 39]</sup> However, the value recently determined by time resolved Raman spectroscopy measurements was pK<sub>a</sub> = 3.4.<sup>[40]</sup>

We observe a more drastic influence of H<sup>+</sup> on the G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) decrease than in ref.<sup>[29]</sup> (Figure 1). The main difference here to that study is the saturation of the solution by CO<sub>2</sub>. The mechanism of the photoelectrochemical CO<sub>2</sub> reduction proposed by Tryk et al.<sup>[16]</sup> involves the reaction of the radical CO<sub>2</sub><sup>•-</sup> with CO<sub>2</sub> forming, at least partially, the transient dimer radical anion O<sub>2</sub>C-CO<sub>2</sub><sup>•-</sup> (reaction 10). Various structures and stabilities of this dimer radical anion have been ab initio calculated.<sup>[41]</sup> Here we assume also that the dimer radical anion O<sub>2</sub>C-CO<sub>2</sub><sup>•-</sup> is formed under CO<sub>2</sub> atmosphere (reaction (10)). The fate of the dimer radical anion is not well known.<sup>[42]</sup> However, we propose that the dissociative disproportionation of O<sub>2</sub>C-CO<sub>2</sub><sup>•-</sup> radical (reaction 11) would be favored vs the dimerization (reaction 12). As it was recently suggested, the disproportionation of O<sub>2</sub>C-CO<sub>2</sub><sup>•-</sup> radical may also lead to CO formation (reaction 14).<sup>[43]</sup> Considering that the oxalate yield at a given pH is less in CO<sub>2</sub>-saturated than in Ar-saturated formate solutions of ref 29, the carbon-carbon bonding in the dimer radical O<sub>2</sub>C-CO<sub>2</sub><sup>•-</sup> structure possibly inhibits a new addition and reorganization into the oxalate in contrast to reaction (4). Because the inflexion point of the oxalate yield increase is shifted to higher pH value than in conditions without CO<sub>2</sub>, the pK<sub>a</sub> (O<sub>2</sub>C-CO<sub>2</sub>H<sup>•</sup>/O<sub>2</sub>C-CO<sub>2</sub><sup>•-</sup>) value is tentatively estimated at about 5.5 (Figure 1). As a result, the influence of protons in the H<sup>+</sup>-catalysed disproportionation is stronger for O<sub>2</sub>C-CO<sub>2</sub><sup>•-</sup> (reaction 11) than for CO<sub>2</sub><sup>•-</sup> (reaction 4).

### ***Effects of high CO<sub>2</sub> pressure***

The pressure increase induces also a marked concentration increase of CO<sub>2</sub> that scavenges rapidly in the radiation spurs the quasi-free electrons ( $e_{\text{qf}}^-$ ) prior the hydration process (reaction (13)) as it was recently observed (Figure 3, red part).<sup>[30]</sup> In these conditions, the source of primary H<sup>•</sup> radicals (reacting with formate and producing H<sub>2</sub>, reaction 7) and of molecular hydrogen (reaction 1) progressively vanishes with the pressure increase. At pH = 3.9 to 3.2 for 1 to 30 atm respectively, we observe a decrease of the formation yield G(H<sub>2</sub>) to zero, and an increase of G(CO) (Figure 2). In contrast, the CO yield increases with the pressure because of the formation of CO via the reaction (9) between CO<sub>2</sub><sup>•-</sup> and the H<sup>•</sup> radicals issued from the dissociation of excited H<sub>2</sub>O\* (Figure 3, red part). The OH<sup>•</sup> radicals are still scavenged by formate and yield CO<sub>2</sub><sup>•-</sup>, but their H<sup>+</sup>-catalysed disproportionation totally inhibits the formation of oxalate (Figure 3). Thus, the formation yield of oxalate is negligible under acidic conditions at any pressure.

In formate solutions initially added with 0.5 mol L<sup>-1</sup> NaOH at increasing CO<sub>2</sub> pressure (final pH = 7 to 5.4 for 1 to 30 atm, respectively), the oxalate yield decreases from G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) = 3.5 × 10<sup>-7</sup> mol J<sup>-1</sup> at 1 atm to a plateau at 1.3 × 10<sup>-7</sup> mol J<sup>-1</sup> at pressures > 10 atm (Figure 2). However, even if the part of CO<sub>2</sub><sup>•-</sup> radicals, arising from the fast (reactions 2) and ultrafast (reaction 13) CO<sub>2</sub> reduction, increases versus the part arising from formate oxidation by OH<sup>•</sup> scavenging (reaction 6), the total yield G(CO<sub>2</sub><sup>•-</sup>) does not change. Above 10 atm (pH 5.4), CO<sub>2</sub><sup>•-</sup> radicals undergo only partly the dimerization into oxalate (reactions 3 and 12) because of the competition with the H<sup>+</sup>-catalysed disproportionation (reaction (11)) (Figure 3 and Table 2)). In fact, the value of the yield G(C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) = 1.3 × 10<sup>-7</sup> mol J<sup>-1</sup> at pressures > 10 atm (Figure 2) is close to the value at the same pH 5 at 1 atm (Figure 1). The effect of pH is thus more decisive than that of CO<sub>2</sub> concentration in the conversion into oxalate.

The mechanism proposed in the present study (Figure 3 and Table 2) could be also of interest for previous results. The  $e_{\text{aq}}^-$  chemistry contributing to scalable CO<sub>2</sub> reduction has recently attracted increasing attention for example in photoelectrocatalytic<sup>[17]</sup> or plasma<sup>[15]</sup> processes where the analogy with the initial reduction of CO<sub>2</sub> by  $e_{\text{aq}}^-$  in radiolysis was underlined. However, the main feature in these studies is that the symbiotic oxidizing species which are unavoidably formed together with  $e_{\text{aq}}^-$ , are responsible of back-reactions and low overall efficiencies of the reduction. In contrast, in the present radiolytic approach the oxidizing species are scavenged by formate, which is chosen to contribute moreover to a selective production of oxalate and H<sub>2</sub>. The mechanism in Figure 3 and Table 2 explains how the radiation radicals are

used and the efficiency is thus optimized. Furthermore, the radiation-induced reduction method is an achievable, promising and scalable route for industrial CO<sub>2</sub> conversion. This method is promoted by the current availability of cost-effective industrial electron accelerators and <sup>60</sup>Co - $\gamma$  sources. Though the aim of our work was not to give details on the economic aspect of this process and its possible valorization, it is known that the price of oxalate is higher than that of sacrificial formate. In addition, the valuable H<sub>2</sub> is produced during this process. Several parameters intervene in the calculation of the total cost of the CO<sub>2</sub> conversion. Eventually, our first estimation suggests that this simple process has a competitive energy cost compared to other chemical processes.

## Conclusion

We demonstrated that the versatility of radiolytic pathways can be harnessed to direct the conversion of aqueous CO<sub>2</sub>, 1 atm / HCO<sub>2</sub>Na solutions at pH 7 at room temperature in order to produce value-added chemicals (oxalate, H<sub>2</sub>) with optimized yields. All the radiolytic radicals are scavenged so that the radiation energy is significantly exploited for converting CO<sub>2</sub> into oxalate and H<sub>2</sub>. The method does not require the presence of any catalyst. The knowledge of radiolytic yield values of water radicals was exploited in order to unravel the reaction mechanisms of all the processes at stake in the systems. The mechanism sheds light also on other previous results. Moreover, we suggest that the ionizing radiation delivered by irradiation facilities may be directly and easily coupled with an industrial source of CO<sub>2</sub>. Therefore, the radiation-induced reducing radicals constitute a promising solution for the CO<sub>2</sub> conversion.

## Acknowledgements

This work was supported by National Natural Science Foundation of China (Nos. [11975122](#), [21906083](#)), National Natural Science Foundation of Jiangsu Province (BK. [2019030384](#)) and Fundamental Research Funds for the Central Universities (N<sup>o</sup>. NE2020006).

**Keywords:** CO<sub>2</sub> conversion • Oxalate • CO<sub>2</sub> radiolysis • CO<sub>2</sub> reduction

---

[1] R. Snoeckx, A. Bogaerts, *Chem. Soc. Rev.* **2010**, *46*, 5805-5863.

[2] J. Albo, M. Alvarez-Guerra, P. Castaño, A. Irabien, *Green Chem.* **2015**, *17*, 2304-2324.

[3] R. M. Cue'llar-Franca, A. J. Azapagic, *CO<sub>2</sub> Util.* **2015**, *9*, 82-102.

[4] P. Lahijani, Z. A. Zainal, M. Mohammadi, A.R. Mohamed, *Renew. Sust. Energ. Reviews.* **2015**, *41*, 615-632.

[5] J. Ma, N. Sun, X. Zhanga, N. Zhao, F. Xiao, W. Wei, Y. Sun, *Cat. Today*, **2019**, *148*, 221-231.

[6] A. Akhundi, A. Habibi-Yangjeh, M. Abitorabi, S. Rahim Pouran, *Cata. Rev. Sci. Eng.* **2019**, *61*, 595-628.

[7] B. Yao, T. Xiao, O. A. Makgae, X. Jie, S. Gonzalez-Cortes, S. Guan, A. I. Kirkland, J. R. Dilworth, H. A. Al-Megren, S. M. Alshihri, P. J. Dobson, G. P. Owen, J. M. Thomas, P. P. Edwards, *Nature Commun.* **2020**, *11*, 1-12.

[8] H. Ooka, M. C. Figueiredo, M. T. Koper, *Langmuir.* **2017**, *33*, 9307-9313.

- 
- [9] T. Burdyny, W. A. Smith, *Ene. Environ. Sci.* **2019**, *12*, 1442-1453.
- [10] W. Tu, Y. Zhou, Z. Zou, *Adv. Mater.* **2014**, *26*, 4607-4626.
- [11] Y. Y. Birdja, E. Pérez-Gallent, M. C. Figueiredo, A. J. Göttle, F. Calle-Vallejo, M. T. Koper, *Nature Energy*. **2019**, *4*, 732-745.
- [12] H. Rao, L. C. Schmidt, J. Bonin, M. Robert, *Nature*. **2017**, *548*, 74-77.
- [13] B. Hu, C. Guild, S. L. Suib, *J. CO<sub>2</sub> Util.* **2013**, *1*, 18-27.
- [14] D. A. Armstrong, R. E. Huie, W. H. Koppenol, S. V. Lymar, G. Mere'nyi, P. Neta, B. Ruscic, D. M. Stanbury, S. Steenkenand, P. Wardman, *Pure Appl. Chem.* **2015**, *87*, 1139-1150.
- [15] P. Rumbach, R. Xu, D.B. Go, *J. Electrochem. Soc.* **2016**, *163*, F1157-F1161.
- [16] D.A. Tryk, T. Yamamoto, M. Kokubun, K. Hirota, K. Hashimoto, M. Okawa, A. Fujishima, *Appl. Organometal. Chem.* **2001**, *15*, 113-120.
- [17] L. Zhang, Z. J. Zhao, T. Wang, J. L. Gong, *Chem. Soc. Rev.* **2018**, *47*, 5423-5443.
- [18] L. Zhang, D. Zhu, G.M. Nathanson, R.J. Hamers, *Angew. Chem. Int. Ed.* **2014**, *126*, 9904-9908.
- [19] M. M. Ramirez-Corredores, G. Gadikota, E. E. Huang, A. M. Gaffney, *Front. Energy Res.* **2020**, *8*, 108.
- [20] A. Lisovskaya, D.M. Bartels, *Radiat. Phys. Chem.* **2019**, *158*, 61-63.
- [21] N. Getoff, *Int. J. Hydrog. Energy.* **1994**, *19*, 667-672.
- [22] A. J. Elliot, D.R. McCracken, G.V. Buxton, N.D. Wood, *J. Chem. Soc. Faraday Trans.* **1990**, *86*, 1539-1547.
- [23] A. M. Lossack, E. Roduner, D.M. Bartels, *Phys. Chem. Chem. Phys.* **2001**, *3*, 2031-2037.
- [24] P. Neta, M. Simic, E. Hayon, *J. Phys. Chem.* **1969**, *73*, 4207-4213.
- [25] J. P. Keene, Y. Raef, A.J. Swallow, In Pulse Radiolysis (Eds.: Ebert, M., Keene, J.P., Swallow, A.J. Baxendale, H.J.), Academic Press, New York, **1965**, pp. 99-106.
- [26] A. Fojtik, G. Czapski, A. Henglein, *J. Phys. Chem.*, **1970**, *74*, 3204-3208.
- [27] G. V. Buxton, R.M. Sellers, *J. Chem. Soc. Faraday Trans. 1*, **1973**, *69*, 555-559.
- [28] M. Z. Lin, Y. Katsumura, Y. Muroya, H. He, T. Miyazaki, D. Hiroishi, *Radiat. Phys. Chem.*, **2008**, *86*, 1208-1212.
- [29] R. Flyunt, M. N. Schuchmann, C. von Sonntag, *Chem. Eur. J.* **2001**, *7*, 796-799.
- [30] A. S. Denisov, M. Mostafavi, *Phys. Chem. Chem. Phys.* **2021**, *23*, 5804-5808.
- [31] G. V. Buxton, C.L. Greenstock, W.P. Helman, A.B. Ross, *J. Phys. Chem. Ref. Data*, **1988**, *17*, 513-886.
- [32] G. V. Buxton, **2008**. An overview of the radiation chemistry of liquids. In: M. Spothem-Maurizot, M. Mostafavi, T. Douki, J. Belloni, (Eds.), Radiation Chemistry from Basics to Application in Material and Life Sciences. EDP Sciences-L'Actualité Chimique, Paris, pp. 3-16.
- [33] S. Gordon, E.J. Hart, M.S. Matheson, J. Rabani, J.K. Thomas, *Disc. Farad. Soc.* **1963**, *36*, 193-205.
- [34] A. K. El Omar, U. Schmidhammer, P. Jeunesse, J. P. Larbre, M. Lin, Y. Muroya, Y. Katsumura, P. Pernot, M. Mostafavi, *J. Phys. Chem. A.* **2011**, *115*, 12212-12216.
- [35] F. Wang, U. Schmidhammer, J. P. Larbre, Z. Z., Zong, J. L. Marignier, M. Mostafavi, *Phys. Chem. Chem. Phys.* **2018**, *20*, 15671-15679.
- [36] R.L. Willson, C.L. Greenstock, G.E. Adams, R. Wageman, L. M. Dorfman, *Int. J. Radiat. Phys. Chem.* **1971**, *3*, 211-220.
- [37] P. Neta, R.H. Schuler, *J. Phys. Chem.* **1972**, *76*, 2673-2679.
- [38] According to the ionization potential of water, each joule may ionize at maximum 10<sup>6</sup> mole of water.
- [39] A.S. Jeevarjan, I. Carmichael, R. W. Fessenden, *J. Phys. Chem.* **1990**, *94*, 1372-1376.
- [40] I. Janik, G. N. R. Tripathi, *J. Chem. Phys.* **2016**, *144*, 154307.
- [41] S. H. Fleischman, K. D. Jordan, *J. Phys. Chem.* **1987**, *91*, 1300-1302.
- [42] D. C. Grills, S. V. Lymar, *Phys. Chem. Chem. Phys.* **2018**, *20*, 10011-10017.
- [43] H. Sheng, M. Hwan Oh, W. T. Osowiecki, W. Kim, A. P. Alivisatos, H. Frei, *J. Am. Chem. Soc.* **2018**, *140*, 4363-4371.