HAL
open science

Self-generated reversed radial electric field in 3D global flux-driven fluid edge plasma turbulence simulations

Gloria L. Falchetto, Patrick Tamain, Hugo Bufferand, G. Ciraolo, Nicolas Fedorczak, P Ghendrih, Elias Laribi, Benjamin Luce, Eric Serre, Raffaele Tatali

To cite this version:

Gloria L. Falchetto, Patrick Tamain, Hugo Bufferand, G. Ciraolo, Nicolas Fedorczak, et al.. Self-generated reversed radial electric field in 3D global flux-driven fluid edge plasma turbulence simulations. EPS 2021 - 47th EPS Conference on Plasma Physics, Jun 2021, Virtual conference, France. 45A, pp.P1.1065, Europhysics conference abstracts - ISBN: 979-10-96389-13-1. cea-03301601

HAL Id: cea-03301601

https://cea.hal.science/cea-03301601

Submitted on 3 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Self-generated reversed radial electric field in 3D global flux-driven fluid edge plasma turbulence simulations

G.L. Falchetto[1], P. Tamain[1], H. Bufferand[1], G. Ciraolo[1], N. Fedorczak[1], Ph. Ghendrih[1], E. Laribi[1], B. Luce[2], E. Serre[2], R. Tatali[2]

[1] CEA, IRFM, F-13108 Saint Paul-lez-Durance, France.
[2]Aix-Marseille University, Laboratoire M2P2, Marseille, France

INTRODUCTION

The formation of a strongly reversed radial electric field in the pedestal region of a tokamak plasma is considered to play a key role in the generation of improved confinement regimes, as the related sheared flow is suspected to strongly impact turbulence, by stabilizing it [1] and leading to the formation of an edge transport barrier and to the transition to high-confinement "H-mode" [2]. However, the mechanisms underlying its formation and interplay with turbulent transport are not yet elucidated.

Here, the dynamics of a **self-generated reversed radial electric field, Er** and its interplay with edge/SOL turbulence are investigated via global flux-driven simulations in 3D geometry with the fluid electrostatic code TOKAM3X [3-6], in response to increasing injected heating power.

TOKAM3X 3D global flux-driven model *[Tamain et al., JCP 321, 2016]*

- Self-consistent **3D electrostatic drift-reduced Braginskii model**, 6 fields: $N, \Gamma\ (u_\parallel), E_e\ (T_e), E_i\ (T_i), W(\phi), J_\parallel$
- **Non-isothermal** turbulence *[Baudoin CPP 58 2018; Tatali NF 61, 2021]*
- **Flux-driven global** approach: no scale separation
- Bohm-Chodura boundary conditions in parallel direction at the limiter *[Stangeby, 2000]*
- Verified via MMS/PoPe/iPoPe (independent Projection on Proper elements) *[Cartier-Michaud et al., PoP 23, 2016; PoP 27, 2020]*
- Versatile geometry (slab/limiter/divertor/different plasma shapes/RMP) *[Galassi et al. NF 57, 2017; Nespoli et al. NF 59, 2019; Luce et al., PPCF 63, 2021]*
- Neutrals via coupling to EIRENE *[Fan et al. Nucl. Mat. Energy 18, 2019]*

Simulation set-up and parameters

Circular limited case, encompassing **closed field lines region (CFR) and Scrape-Off-Layer (SOL)** r/a=0.8-1.2 (including buffers)
COMPASS-like tokamak plasma parameters:

R/a	ρ_*	$\nu_*\left(\frac{\nu_{coll}}{\omega_c}\right)$
3.4	1/256	$5\cdot10^{-2}$

Scan on injected core heating:
n freely evolving



Mesh resolution on ½ torus : (ψ,θ,φ) 64 x 512 x 64
Injected heating increased in two steps from **baseline** simulation: [**P**, **Px1.5**, **Px2**]
Runs > 10^6 t (ω_{ci}^{-1}) up to quasi-stationary state ~ [5, 3.8, 1] ms

Self-consistent reversed E_r generation & profile steepening



- E_r shear max in TOKAM3X simulations (w/o recycling) systematically @ few Larmor radii outside separatrix, in near SOL region similar to *[Chankin PPCF 2017]*
- **Reversed E_r peak @separatrix, strongly increases with increasing injected power**
- **Global density profile steepening with increasing power** and formation of a **pedestal in temperature**





- E_r **oscillation** ~ f_{GAM} *[Sugama JPP 2006]*
  -similarity to LCO/I-phase? *to be further investigated*
  Cfr in AUG *[Cavedon NF 2017 Medvedeva PPCF 2017]*
  in WEST ***Vermare P4.1078 EPS 2021***



E_r shear interplay with turbulence: fluctuation level & λ_N



Slight reduction of fluctuation level in CFR
increase in near SOL

Density radial decay length λ_N decreases with power

	$\lambda_n\ (\rho_L)$
P	24.9
Px1.5	18.8
Px2	15.3

$$\partial_\psi \langle N\rangle_t = \langle N\rangle_t / \lambda_N$$

Particle and energy fluxes: from turbulent to diffusive

Flux-driven simulation, with constant particle flux incoming from the core ➔ radial particle flux constant.
The radial particle and electron energy fluxes can be decomposed as:

$$\bar{\Gamma}^r_N = \overline{Nu^r_{E\times B}} + \overline{Nu^r_{\nabla B}} + \bar{\Gamma}^r_{N,D_\perp}$$

ExB flux dominating — In SOL ∇B flux > 0 dominates — diffusive flux In CFR ↗ dominates

Baseline case
Power x 2 ↘



$$\bar{\Gamma}^r_{E_e} = \overline{NT_e u^r_{E\times B}} + \overline{NT_e u^r_{\nabla B}} + \bar{\Gamma}^r_{E_e,D_\perp}$$



Increased power affects the particle and energy fluxes:

- **transport from turbulent to diffusive in the CFR** : no evident impact of Er shear on the **turbulence stabilization**
- interplay with Er shear in the pedestal



Energy ExB flux dominantly **conducted** in SOL

Discussion and perspectives

- **Systematic spontaneous generation of a reversed radial electric field E_r** in the proximity of the separatrix is found in TOKAM3X electrostatic non-isothermal turbulence simulations in a limited circular plasma, encompassing CFR and SOL.
- ✓ **E_r well strongly increases with increasing injected power**, consistently with experimental observations.
- ✓ **Global steepening of the density profile** with increasing power.
- ✓ **Plasma temperature profiles steepen**, with the creation of a **strong gradient in the near SOL, reminiscent of the narrow λ_q feature**.
- **Interplay of the self-generated reversed E_r with turbulence**:
  - **particle and energy transport fluxes in the CFR from turbulent to diffusive,**
  - **dominantly conducted energy fluxes in the SOL**
  - **turbulence stabilization in the closed field lines region mainly resulting from reduced collisionality due to the increased T** *[Falchetto PRL 2004; Tatali NF 2021]*
- Upcoming, simulations in **more realistic edge plasma conditions**
  - particle source at the limiter
  - recycling neutrals
  - divertor geometry
  - neoclassical viscosity
  - realistic collisionality

➔ Recently released **SOLEDGE3X code [8]**
cfr Bufferand IAEA 2021

References [1] K. Burrell, Physics of Plasmas 6, 4418 (1999); [2] F. Wagner, et al. Phys. Rev. Lett. 49, 1408–1412 (1982).
[3] P. Tamain, et al., Journal Comput. Physics 321, 606-623 (2016); [4] D. Galassi, et al., Nuclear Fusion 57, 036029 (2017)
[5] R. Tatali et al., Nucl. Fusion 61, 056002 (2021); [6] B. Luce, et al., Plasma Phys. Control. Fusion 63 055017 (2021)
[7] A. V. Chankin, et al., Plasma Phys. Control. Fusion 59 (4), 045012 (2017)
[8] H. Bufferand et al., Nuclear Materials and Energy, 18 (2019); [9] H Bufferand et al., IAEA 2021, submitted to Nucl. Fus.

This work was supported by the EUROfusion - Theory and Advanced Simulation Coordination (E-TASC).
This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 and 2019-2020 under grant agreement No 633053.
The views and opinions expressed herein do not necessarily reflect those of the European Commission.