



## Photocatalytic deoxygenation of nitrogen oxide compounds

Marianne Kjellberg, Emmanuel Nicolas, Lucile Anthore-Dalion, Alexia Ohleier, Annamaria Quaranta, Thibault Cantat

### ► To cite this version:

Marianne Kjellberg, Emmanuel Nicolas, Lucile Anthore-Dalion, Alexia Ohleier, Annamaria Quaranta, et al.. Photocatalytic deoxygenation of nitrogen oxide compounds. GECOM-CONCOORD 2022, May 2022, Sévrier, France. cea-03662299

**HAL Id: cea-03662299**

**<https://cea.hal.science/cea-03662299>**

Submitted on 9 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Photocatalytic deoxygenation of nitrogen oxide compounds

M. Kjellberg,<sup>a\*</sup> E. Nicolas,<sup>a</sup> L. Anthore-Dalion,<sup>a</sup> A. Ohleier,<sup>a</sup> A. Quaranta,<sup>b</sup> T. Cantat<sup>a</sup>

<sup>a</sup> CEA Saclay – Laboratoire de Chimie Moléculaire et de Catalyse pour l'Energie


<sup>b</sup> CEA Saclay – Laboratoire des Mécanismes fondamentaux de la Bioénergétique

marianne.kjellberg@cea.fr

The accumulation of nitrogen oxides in the environment calls for new pathways to interconvert the various oxidation states of nitrogen, and especially their reduction. However, the large spectrum of reduction potentials covered by nitrogen oxides makes it difficult to find general systems capable of efficiently reducing various *N*-oxides.

Commonly known as 'laughing gas', nitrous oxide N<sub>2</sub>O is an ozone-depleting substance 298 times more powerful a greenhouse gas than CO<sub>2</sub>. It accumulates in the atmosphere due to its kinetic stability (120 years), thus contributing to 6% of anthropogenic greenhouse effect.<sup>1,2</sup> A solution would be the reduction of N<sub>2</sub>O back to N<sub>2</sub>, which would close the nitrogen cycle; however, few methods have been proposed due to the low reactivity of N<sub>2</sub>O. Herein we describe a new pathway to reduce N<sub>2</sub>O back to nitrogen at room temperature by means of photocatalysis. N<sub>2</sub>O being isoelectronic with CO<sub>2</sub>, inspiration came from previous work on the photoreduction of CO<sub>2</sub> to CO first disclosed by the group of Lehn.<sup>3</sup> Remarkably, [Re(bpy)(CO)<sub>3</sub>Cl] and its derivatives proved successful for the reduction of N<sub>2</sub>O, up to 86% yield.

Encouraged by this success, we extended this methodology to the reduction of other N–O bonds, leading to a selective and synthetically applicable photocatalytic version of the deoxygenation of pyridine *N*-oxides. The mechanism of the N–O bond deoxygenation was studied on this substrate using laser flash photolysis and spectroelectrochemistry. These mechanistic insights enabled to optimize the system by shortening the reaction time from 34 h to 2 h. This unlocked the path to the deoxygenation of a broader scope of organic, *N*-oxide compounds.



<sup>1</sup> D. Ussiri, R. Lal, *Soil Emission of Nitrous Oxide and its Mitigation*, **2013**, 243-275.

<sup>2</sup> M. Dameris, *Angew. Chem. Int. Ed. Engl.*, **2010**, 49, 489-491.

<sup>3</sup> J. Hawecker, J.-M. Lehn, R. Ziessel, *J. Chem. Soc., Chem. Commun.*, **1983**, 536-538.

<sup>4</sup> M. Kjellberg, A. Ohleier, P. Thuéry, E. Nicolas, L. Anthore-Dalion, T. Cantat, *Chem. Sci.* **2021**, 12, 10266-10272.