



## Reduction of Carboxamides to Imines Catalyzed by the Schwartz Reagent

Liam Donnelly, Jean Claude Berthet, Thibault Cantat

### ► To cite this version:

Liam Donnelly, Jean Claude Berthet, Thibault Cantat. Reduction of Carboxamides to Imines Catalyzed by the Schwartz Reagent. Symposium International sur la chimie durable (ISGC 2022), May 2022, La Rochelle, France. cea-03675182

HAL Id: cea-03675182

<https://cea.hal.science/cea-03675182>

Submitted on 23 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

N°854 / OC

TOPIC(s) : Homogenous, heterogenous and biocatalysis / Clean reactions

## Reduction of Carboxamides to Imines Catalyzed by the Schwartz Reagent

### AUTHORS


Liam DONNELLY / CEA SACLAY, IRAMIS/NIMBE/LCMCE BÂT 125, GIF-SUR-YVETTE CEDEX  
Jean-Claude BERTHET / CEA SACLAY, IRAMIS/NIMBE/LCMCE BÂT 125, GIF-SUR-YVETTE CEDEX  
Thibault CANTAT / CEA SACLAY, IRAMIS/NIMBE/LCMCE BÂT 125, GIF-SUR-YVETTE CEDEX

### PURPOSE OF THE ABSTRACT

Carboxamides are prevalent in pharmaceuticals, agrochemicals and natural products as well as a variety of functional materials. Therefore, methods to reduce these bonds could open new opportunities for late-stage functionalization and post-polymerization modification to access new chemicals that are difficult to synthesize by classical methods.[1] However, the carboxamide C=O bond is relatively inert to hydride addition, which means that forcing conditions are often required, leading to issues with functional group tolerance and chemoselectivity. Several catalytic protocols have been developed to enable the use of milder reducing agents; however, most of these protocols form amines as the final product and typically proceed by initial formation of the imine equivalent followed by further reduction.[2] The development of methodologies that avoid overreduction to the amine product and instead allow selective formation of the corresponding imine are rare but highly valuable and can complement traditional methods for imine synthesis.[3]

Ganem and co-workers showed that the commercially available and well-defined Schwartz reagent ( $\text{Cp}_2\text{Zr}(\text{H})\text{Cl}$ ) can reduce carboxamides to imines stoichiometrically with liberation of  $(\text{ClZrCp}_2)_2\text{O}$ .[4] Inspired by this work, we report the use of silanes as stoichiometric reductants to unlock a catalytic protocol for this transformation, using  $\text{Cp}_2\text{Zr}(\text{H})\text{Cl}$  as a catalyst for the deoxygenation of a range of secondary carboxamides. This catalytic system retains the excellent chemoselectivity observed for the original stoichiometric protocol. We have investigated the mechanism of this reaction through the isolation of potential reactive intermediates as well as developing one-pot processes for the further derivitization of the imine products. This work presents a rare example of Zr hydride catalysis, enabling the practical and efficient functionalization of a challenging class of substrates without recourse to highly sensitive reagents or rare and expensive metals.

## FIGURES



**FIGURE 1**

Scheme 1.

The Zr-catalysed reduction of carboxamides to imines.

**FIGURE 2**

## KEYWORDS

Homogeneous Catalysis | Amide | Hydrosilylation | Zirconium

## BIBLIOGRAPHY

- [1] Czerwinski, P. J.; Furman, B., *Front. chem.* 2021, 9, 1-10
- [2] Volkov, A.; Tinnis, F.; Slagbrand, T.; Trilloa, P.; Adolfsson, H., *Chem. Soc. Rev.*, 2016, 45, 6685-6697
- [3] (a) Pelletier, G.; Bechara, W. S.; Charette, A. B., *J. Am. Chem. Soc.* 2010, 132, 12817-12819 (b) Chen, C.; Brookhart, M., *J. Am. Chem. Soc.* 2012, 134, 11304-11307.
- [4] Schedler, D. J. A.; Li, J.; Ganem, B., *J. Org. Chem.* 1996, 61, 4115-4119.